

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

**EP 1 054 064 A1**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:  
22.11.2000 Bulletin 2000/47

(51) Int. Cl.<sup>7</sup>: **C12N 15/86**, C12N 15/34,  
C12N 5/10, A61K 48/00

(21) Application number: 00201738.2

(22) Date of filing: 16.05.2000

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU**  
**MC NL PT SE**  
Designated Extension States:  
**AL LT LV MK RO SI**

(30) Priority: 17.05.1999 EP 99201545

(71) Applicant: **Introgene B.V.**  
**2333 AL Leiden (NL)**

(72) Inventors:  
• **Bout, Abraham**  
**2751 AR Moerkapelle (NL)**  
• **Vogels, Ronald**  
**3461 HW Linschoten (NL)**

• **Havenga, Menzo Jans Emco**  
**2401 BV Alphen aan den Rijn (NL)**

(74) Representative:  
**Prins, Adrianus Willem et al**  
**Vereenigde,**  
**Nieuwe Parklaan 97**  
**2587 BN Den Haag (NL)**

Remarks:

The sequence listing, which is published as annex to the application documents, was filed after the date of filing. The applicant has declared that it does not include matter which goes beyond the content of the application as filed.

(54) **Adenovirus derived gene delivery vehicles comprising at least one element of adenovirus type 35**

(57) The serotypes differ in their natural tropism. The adenovirus serotypes 2, 4, 5 and 7 all have a natural affiliation towards lung epithelia and other respiratory tissues. In contrast, serotypes 40 and 41 have a natural affiliation towards the gastrointestinal tract. The serotypes described above, differ in at least capsid proteins (penton-base, hexon), proteins responsible for cell binding (fiber protein), and proteins involved in adenovirus replication. This difference in tropism and capsid protein among serotypes has led to the many research efforts aimed at redirecting the adenovirus tropism by modification of the capsid proteins

**EP 1 054 064 A1**

## Description

[0001] The present invention relates to the field of gene therapy, in particular gene therapy involving elements derived from viruses, more in particular elements of adenoviruses. Adenoviruses have been proposed as suitable vehicles to deliver genes to the host.

[0002] There are a number of features of adenoviruses that make them particularly useful for the development of gene-transfer vectors for human gene therapy:

The adenovirus genome is well characterized. It consists of a linear double-stranded DNA molecule of approximately 36000 base pairs. The adenovirus DNA contains identical Inverted Terminal Repeats (ITR) of approximately 90-140 base pairs with the exact length depending on the serotype. The viral origins of replication are within the ITRs exactly at the genome ends;

The biology of the adenoviruses is characterized in detail; the adenovirus is not associated with severe human pathology in immuno-competent individuals

The virus is extremely efficient in introducing its DNA into the host cell; the virus can infect a wide variety of cells and has a broad host-range;

The virus can be produced at high virus titers in large quantities;

The virus can be rendered replication defective by deletion of the early-region 1 (E1) of the viral genome (Brody et al, 1994). Most adenoviral vectors currently used in gene therapy have a deletion in the E1 region, where desired genetic information can be introduced.

[0003] Based on these features, preferred methods for in vivo gene transfer into human target cells, make use of adenoviral vectors as gene delivery vehicles.

[0004] However, there are still drawbacks associated with the therapeutic use of adenoviral vectors in humans. A major drawback is the existence of widespread pre-existing immunity among the population against adenoviruses. Exposure to wild-type adenoviruses is very common in humans, as has been documented extensively [reviewed in Wadell, 1984]. This exposure has resulted in immune responses against most types of adenoviruses, not alone against adenoviruses to which individuals have actually been exposed, but also against adenoviruses which have similar (neutralizing) epitopes.

[0005] This phenomenon of pre-existing antibodies in humans, in combination with a strong secondary humoral and cellular immune response against the virus, can seriously affect gene transfer using recombinant adenoviral vectors.

[0006] To date, six different subgroups of human adenoviruses have been proposed which in total encompasses 51 distinct adenovirus serotypes (see table 1). A serotype is defined on the basis of its immunological distinctiveness as determined by quantitative neutralization with animal antisera (horse, rabbit). If neutralization shows a certain degree of cross-reaction between two viruses, distinctiveness of serotype is assumed if A) the hemagglutinins are unrelated, as shown by lack of cross-reaction on hemagglutination-inhibition, or B) substantial biophysical/ biochemical differences in DNA exist (Francki et al, 1991). The nine serotypes identified last (42-51) were isolated for the first time from HIV-infected patients (Hierholzer et al 1988; Schnurr et al 1993;). For reasons not well understood, most of such immune-compromised patients shed adenoviruses that were rarely or never isolated from immune-competent individuals (Hierholzer et al 1988, 1992; Khoo et al, 1995, De Jong et al, 1998).

[0007] The vast majority of individuals have had previous exposure to adenoviruses, especially the well investigated adenovirus serotypes 5 and type 2 (Ad5 and Ad2) or immunologically related serotypes. Importantly, these two serotypes are also the most extensively studied for use in human gene therapy. from other (adeno) viruses, as long as one replaces an element which could lead to immunity against such a gene delivery vehicle by an element of adenovirus 35 or a functional homologue thereof, which has less of such a drawback and which preferably avoids such a drawback. In the present invention a gene delivery vehicle is any vehicle that is capable of delivering a nucleic acid of interest to a host cell. It must, according to the invention comprise an element of adenovirus 35 or a functional equivalent of such an element, which must have a beneficial effect regarding the immune response against such a vehicle. Basically all other elements making up the vehicle can be any elements known in the art or developed in the art, as long as together they are capable of delivering said nucleic acid of interest. In principle the person skilled in the art can use and/or produce any adenoviral products or production systems that can or have been applied in the adenoviral field.

[0008] Typically the products of the invention can be made in the packaging cells useable for e.g. adenovirus 5, typically the vectors based on adenovirus 35 can be produced and/or used in the same manner as those of other adenoviruses e.g. adenovirus 2 and/or 5. A good overview of the possibilities of minimal vectors, packaging systems, intracellular amplification, vector and plasmid based systems can be found in applicant's copending applications (PCT/NL99/00235 and PCT/NL96/00244) incorporated herein by reference. Non-viral delivery systems can also be provided with elements according to the invention as can viral delivery systems.



[0009] Both kinds of systems are well known in the art in many different set-ups and do therefor not need any further elaboration here. A review on the many different systems and their properties can be found in Robbins and Ghivizani (1998) and in Prince (1998) incorporated herein by reference.

[0010] Gene delivery vehicles typically contain a nucleic acid of interest. A nucleic acid of interest can be a gene or a functional part of a gene (wherein a gene is any nucleic acid which can be expressed) or a precursor of a gene or a transcribed gene on any nucleic acid level (DNA and/or RNA: double or single stranded). Genes of interest are well known in the art and typically include those encoding therapeutic proteins such as TPA, EPO, cytokines, antibodies or derivatives thereof, etc. An overview of therapeutic proteins to be applied in gene therapy are listed below.

Immune-stimulatory factors like tumor-specific antigens, cytokines, etc.;

Anti-angiogenic factors non-limiting examples endostatin, angiostatin, ATF-BPTI CDT-6, dominant negative VEGF-mutants, etc.;

Angiogenic factors non-limiting example VEGF, Fibroblast growth factors, Nitric oxide synthases, C-type natriuretic peptide, etc.;

Inflammation inhibiting proteins like soluble CD40, FasL, IL-12, IL-10, IL-4, IL-13 and excreted single chain antibodies to CD4, CD5, CD7, CD52, IL-2, IL-1, IL-6, TNF, etc. or excreted single chain antibodies to the T-cell receptor on the auto-reactive T-cells. Also, dominant negative mutants of PML may be used to inhibit the immune response.

[0011] Furthermore, antagonists of inflammation promoting cytokines may be used, for example IL-1RA(receptor antagonist) and soluble receptors like sIL-1RI, sIL-1RII, sTNFRI and sTNFRII. Growth and/or immune response inhibiting genes such as ceNOS, Bcl3, cactus and I $\kappa$ B $\alpha$ ,  $\beta$  or  $\gamma$  and apoptosis inducing proteins like the VP3 protein of chicken anemia virus may also be used. Furthermore, suicide genes like HSV-TK, cytosine deaminase, nitroreductase and linamerase may be used.

[0012] A nucleic acid of interest may also be a nucleic acid which can hybridise with a nucleic acid sequence present in the host cell thereby inhibiting expression or transcription or translation of said nucleic acid. It may also block through cosuppression. In short a nucleic acid of interest is any nucleic acid that one may wish to provide a cell with in order to induce a response by that cell, which response may be production of a protein, inhibition of such production, apoptosis, necrosis, proliferation, differentiation etc.

[0013] The present invention is the first to disclose adenovirus 35 or a functional homologue thereof for therapeutic use, therefor the invention also provides an adenovirus serotype 35 or a functional homologue thereof or a chimaeric virus derived therefrom, or a gene delivery vehicle based on said virus its homologue or its chimaera for use as a pharmaceutical. The serotype of the present invention, adenovirus type 35, is in itself known in the art. It is an uncommon group B adenovirus that was isolated from patients with acquired immunodeficiency syndrome and other immunodeficiency disorders (Flomenberg *et al.*, 1987; De Jong *et al.*, 1983). Ad 35 has been shown to differ from the more fully characterized subgroup C (including Ad2 and Ad5) with respect to pathogenic properties (Basler *et al.*, 1996). It has been suggested that this difference may be correlated with differences in the E3 region of the Ad35 genome (Basler *et al.*, 1996). The DNA of Ad35 has been partially cloned and mapped (Kang *et al.*, 1989a and b; Valderrama-Leon *et al.*, 1985).

[0014] B type adenovirus serotypes such as 34 and 35 have a different E3 region than other serotypes. Typically this region is involved in suppressing immune response to adenoviral products. Thus the invention provides a gene delivery vehicle according to the invention whereby said elements involved in avoiding or diminishing immune response comprise adenovirus 35 E3 expression products or the genes encoding them or functional equivalents of either or both.

[0015] Another part of adenoviruses involved in immune responses is the capsid, in particular the penton and/or the hexon proteins. Thus the invention also provides a gene delivery vehicle according to the invention whereby the elements comprise at least one adenovirus 35 capsid protein or functional part thereof, such as fiber, penton and/or hexon proteins or a gene encoding at least one of them. It is not necessary that a whole protein relevant for immune response is of adenovirus 35 (or a functional homologue thereof) origin. It is very well possible to insert a part of an adenovirus fiber, penton or hexon protein into another fiber, penton or hexon. Thus chimaeric proteins are obtained.

[0016] It is also possible to have a penton of a certain adenovirus, a hexon from another and a fiber or an E3 region from yet another adenovirus. According to the invention at least one of the proteins or genes encoding them should comprise an element from adenovirus 35 or a functional homologue thereof, whereby said element has an effect on the immune response of the host. Thus the invention provides a gene delivery according to the invention, which is a chimaera of adenovirus 35 with at least one other adenovirus. In this way one can also modify the resulting virus in other aspects than the immune response alone. One can enhance its efficiency of infection with elements responsible therefor; one can enhance its replication on a packaging cell, or one can change its tropism.

[0017] Thus the invention e.g. provides a gene delivery vehicle according to the invention which has a different tropism than adenovirus 35. Of course the tropism should be altered preferably such that the gene delivery vehicle is delivered preferentially to a subset of the host's cells, i.e. the target cells. Changes in tropism and other changes which can

also be applied in the present invention of adenoviral or other gene delivery vehicles are disclosed in applicant's copending applications (nos. 98204482.8, 99200624.7 and 98202297.2) incorporated herein by reference. Of course the present application also provides any and all building blocks necessary and/or useful to get to the gene delivery vehicles and/or the chimaeras, etc. of the present invention. This includes packaging cells such as PER.C6 (ECACC deposit number 96022940) or cells based thereon, but adapted for Ad35 or a functional homologue thereof; it also includes any nucleic acids encoding functional parts of adenovirus 35 or a functional homologue thereof, such as helper constructs and packaging constructs, as well as vectors comprising genes of interest and e.g. an ITR, etc.

[0018] Typically applicant's application (PCT/NL96/00244) incorporated herein by reference, discloses elements necessary and useful for arriving at the invented gene delivery vehicles. Thus the invention also provides a nucleic acid encoding at least a functional part of a gene delivery vehicle according to the invention, or a virus, homologue or chimaera thereof according to the invention.

[0019] According to the invention, such elements, which encode functions that will end up in the resulting gene delivery vehicle must comprise or be encoded by a nucleic acid encoding at least one of the adenovirus serotype 35 elements or a functional equivalent thereof, responsible for avoiding or diminishing neutralising activity against adenoviral elements by the host to which the gene is to be delivered.

[0020] Typically the gene of interest would be present on the same nucleic acid which means that such a nucleic acid has such a gene or that it has a site for introducing a gene of interest therein.

[0021] Typically such a nucleic acid also comprises at least one ITR and if it is a nucleic acid to be packaged also a packaging signal. However, as mentioned before all necessary and useful elements and/or building blocks for the present invention can be found in applicant's application (PCT/NL96/00244). A set of further improvements in the field of producing adenoviral gene delivery vehicles is applicant's plasmid system disclosed in PCT/NL99/00235 mentioned herein before. This system works in one embodiment as a homologous recombination of an adapter plasmid and a longer plasmid, together comprising all elements of the nucleic acid to be incorporated in the gene delivery vehicle. These methods can also be applied to the presently invented gene delivery vehicles and their building elements.

[0022] Thus the invention also provides a nucleic acid according to the invention further comprising a region of nucleotides designed or useable for homologous recombination, preferably as part of at least one set of two nucleic acids comprising a nucleic acid according to the invention, whereby said set of nucleic acids is capable of a single homologous recombination event with each other, which leads to a nucleic acid encoding a functional gene delivery vehicle.

[0023] Both empty packaging cells (in which the vector to be packaged to make a gene delivery vehicle according to the invention still has to be introduced or produced) as well as cells comprising a vector according to the invention to be packaged are provided. Thus the invention also encompasses a cell comprising a nucleic acid according to the invention or a set of nucleic acids according to the invention, preferably a cell which complements the necessary elements for adenoviral replication which are absent from the nucleic acid according to be packaged, or from a set of nucleic acids according to the invention. In the present invention it has been found that E1-deleted adenovirus 35 vectors, are not capable of replication on cells that provide adenovirus 5 proteins *in trans*. The invention therefore further provides a cell capable of providing adenovirus 35 E1 proteins in trans. Such a cell is typically a human cell derived from the retina or the kidney. Embryonal cells such as amniocytes, have been shown to be particularly suited for the generation of an E1 complementing cell line. Such cells are therefor preferred in the present invention. Serotype specific complementation by E1 proteins can be due to one or more protein(s) encoded by the E1 region. It is therefor essential that at least the serotype specific protein is provided in trans in the complementing cell line. The non-serotype specific E1 proteins essential for effective complementation of an E1-deleted adenovirus can be derived from other adenovirus serotypes. Preferably, at least an E1 protein from the E1B region of adenovirus 35 is provided in trans to complement E1-deleted adenovirus 35 based vectors.

[0024] In one embodiment nucleic acid encoding the one or more serotype specific E1-proteins is introduced into the PER.C6 cell or a cell originating from a PER.C6 cell (ECACC deposit number 96022940), or a similar packaging cell complementing with elements from Ad 35 or a functional homologue thereof.

[0025] As already alluded to the invention also encompasses a method for producing a gene delivery vehicle according to the invention, comprising expressing a nucleic acid according to the invention in a cell according to the invention and harvesting the resulting gene delivery vehicle. The above refers to the filling of the empty packaging cell with the relevant nucleic acids. The format of the filled cell is of course also part of the present invention, which provides a method for producing a gene delivery vehicle according to the invention, comprising culturing a filled packaging cell (producer cell) according to the invention in a suitable culture medium and harvesting the resulting gene delivery vehicle.

[0026] The resulting gene delivery vehicles obtainable by any method according to the invention are of course also part of the present invention, particularly also a gene delivery vehicle according to the invention, which is derived from a chimaera of an adenovirus and an integrating virus.

[0027] It is well known that adenoviral gene delivery vehicles do not integrate into the host genome normally. For

long term expression of genes in a host cell it is therefor preferred to prepare chimaeras which do have that capability. Such chimaeras have been disclosed in our copending application PCT/NL98/00731 incorporated herein by reference. A very good example of such a chimaera of an adenovirus and an integrating virus wherein said integrating virus is an adeno associated virus. As discussed hereinbefore other useful chimaeras, which can also be combined with the above are chimaeras (be it in swapping whole proteins or parts thereof or both) which have altered tropism. A very good example thereof is a chimaera of Ad 35 and Ad 16, possibly with elements from for instance Ad 2 or Ad 5, wherein the tropism determining part of Ad 16 or a functional equivalent thereof is used to direct the gene delivery vehicle to synoviocytes and/or smooth muscle cells (see our copending applications nos. 98204482.8 and 99200624.7) incorporated herein by reference). Dendritic cells (DC) and hemopoietic stem cells (HSC) are not easily transduced with Ad2 or Ad5 derived gene delivery vehicles. The present invention provides gene delivery vehicles that posses increased transduction capacity of DC and HSC cells. Such gene delivery vehicles at least comprises the tissue tropism determining part of an Ad35 adenovirus. The invention therefore further provides the use of a tissue tropism determining part of an adenovirus 35 capsid for transducing dendritic cells and/or hemopoietic stem cells. Other B-type adenoviruses are also suited. A tissue tropism determining part comprises at least the knob and/or the shaft of a fiber protein. Of course it is very well possible for a person skilled in the art to determine the amino acid sequences responsible for the tissue tropism in the fiber protein. Such knowledge can be used to devise chimearic proteins comprising such amino acid sequences. Such chimaeric proteins are therefor also part of the invention. DC cells are very efficient antigen presenting cells. By introducing the gene delivery vehicle into such cells the immune system of the host can be triggered to toward specific antigens. Such antigens can be encoded by nucleic acid delivered to the DC or by the proteins of the gene delivery vehicle it self. The present invention therefor also provides a gene delivery vehicle with the capacity to evade to host immune system as a vaccine. The vector being capable to evade the immune system long enough to efficiently find it target cells and at the same time capable of delivering specific antigens to antigen presenting cells thereby allowing the induction and/or stimulation of an efficient immune responses toward the specific antigen(s). To further modulate the immune response, the gene delivery vehicle may comprise proteins and/or nucleic encoding such proteins capable of modulating an immune response. Non-limiting examples of such proteins are found among the interleukins, the adhesion molecules, the co-stimulatory proteins, the interferons etc. The invention therefore further provides a vaccine comprising a gene delivery vehicle of the invention. The invention further provides an adenovirus vector with the capacity to efficiently transduce DC and/or HSC, the vehicle comprising at least a tissue tropism determing part of serotype 35 adenvirus. The invention further provides the use of such delivery vehicles for the transduction of HSC and/or DC cells. Similar tissue tropisms are found among other adenoviruses of serotype B, particularly in serotype 11 and are also part of the invention. Of course it is also possible to provide other gene delivery vehicles with the tissue tropism determining part thereby providing such delivery vehicles with an enhanced DC and/or HSC transduction capacity. Such gene delivery vehicles are therefor also part of the invention.

[0028] The gene delivery vehicles according to the invention can be used to deliver genes or nucleic acids of interest to host cells. This will typically be a pharmaceutical use. Such a use is included in the present invention. Compositions suitable for such a use are also part of the present invention. The amount of gene delivery vehicle that needs to be present per dose or per infection (m.o.i) will depend on the condition to be treated, the route of administration (typically parenteral) the subject and the efficiency of infection, etc. Dose finding studies are well known in the art and those already performed with other (adenoviral) gene delivery vehicles can typically be used as guides to find suitable doses of the gene delivery vehicles according to the invention. Typically this is also where one can find suitable excipients, suitable means of administration, suitable means of preventing infection with the vehicle where it is not desired, etc. Thus the invention also provides a pharmaceutical formulation comprising a gene delivery vehicle according to the invention and a suitable excipient, as well as a pharmaceutical formulation comprising an adenovirus, a chimaera thereof, or a functional homologue thereof according to the invention and a suitable excipient.

#### Detailed description

[0029] As described above, the most extensively studied serotypes of adenovirus are not ideally suitable for delivering additional genetic material to host cells. This is partly due to the pre-existing immunity among the population against these serotypes. This presence of pre-existing antibodies in humans, in combination with a strong secondary humoral and cellular immune response against the virus will affect adenoviral gene therapy.

[0030] The present invention provides the use of at least elements of a serotype and functional homologues thereof of adenovirus which are very suitable as gene therapy vectors.

[0031] The present invention also discloses an automated high-throughput screening of all known adenovirus serotypes against sera from many individuals. Surprisingly, no neutralizing ability was found in any of the sera that were evaluated against one particular serotype, adenovirus 35 (Ad35). This makes the serotype of the present invention extremely useful as a vector system for gene therapy in man.

[0032] Such vector system is capable of efficiently transferring genetic material to a human cell without the inherent



problem of pre-existing immunity.

[0033] Typically, a virus is produced using an adenoviral vector (typically a plasmid, a cosmid or baculovirus vector). Such vectors are of course also part of the present invention. The invention also provides adenovirus derived vectors that have been rendered replication defective by deletion or inactivation of the E1 region. Of course, also a gene of interest can be inserted at for instance the site of E1 of the original adenovirus from which the vector is derived.

[0034] In all aspects of the invention the adenoviruses may contain deletions in the E1 region and insertions of heterologous genes linked either or not to a promoter. Furthermore, the adenoviruses may contain deletions in the E2, E3 or E4 regions and insertions of heterologous genes linked to a promoter. In these cases, E2 and/or E4 complementing cell lines are required to generate recombinant adenoviruses.

[0035] One may choose to use the Ad35 serotype itself for the preparation of recombinant adenoviruses to be used in gene therapy. Alternatively, one may choose to use elements derived from the serotype of the present invention in such recombinant adenoviruses. One may for instance develop a chimaeric adenovirus that combines desirable properties from different serotypes. Some serotypes have a somewhat limited host range, but have the benefit of being less immunogenic, some are the other way round. Some have a problem of being of a limited virulence, but have a broad host range and/or a reduced immunogenicity. Such chimaeric adenoviruses are known in the art, and they are intended to be within the scope of the present invention. Thus in one embodiment the invention provides a chimaeric adenovirus comprising at least a part of the adenovirus genome of the present serotype, providing it with absence of pre-existing immunity, and at least a part of the adenovirus genome from another adenovirus serotype resulting in a chimaeric adenovirus. In this manner the chimaeric adenovirus produced is such that it combines the absence of pre-existing immunity of the serotype of the present invention, to other characteristics of another serotype. Such characteristics may be temperature stability, assembly, anchoring, redirected infection, production yield, redirected or improved infection, stability of the DNA in the target cell, etc.

[0036] A packaging cell will generally be needed in order to produce sufficient amount of adenoviruses. For the production of recombinant adenoviruses for gene therapy purposes, several cell lines are available. These include but are not limited to the known cell lines PER.C6 (ECACC deposit number 96022940), 911, 293, and E1 A549.

[0037] An important feature of the present invention is the means to produce the adenovirus. Typically, one does not want an adenovirus batch for clinical applications to contain replication competent adenovirus. In general therefore, it is desired to omit a number of genes (but at least one) from the adenoviral genome on the adenoviral vector and to supply these genes in the genome of the cell in which the vector is brought to produce chimaeric adenovirus. Such a cell is usually called a packaging cell. The invention thus also provides a packaging cell for producing an adenovirus (a gene delivery vehicle) according to the invention, comprising in trans all elements necessary for adenovirus production not present on the adenoviral vector according to the invention. Typically vector and packaging cell have to be adapted to one another in that they have all the necessary elements, but that they do not have overlapping elements which lead to replication competent virus by recombination.

[0038] Thus the invention also provides a kit of parts comprising a packaging cell according to the invention and a recombinant vector according to the invention whereby there is essentially no sequence overlap leading to recombination resulting in the production of replication competent adenovirus between said cell and said vector.

[0039] Thus the invention provides methods for producing adenovirus, which upon application will escape pre-existing humoral immunity, comprising providing a vector with elements derived from an adenovirus serotype against which virtually no natural immunity exists and transfecting said vector in a packaging cell according to the invention and allowing for production of viral particles.

[0040] In one aspect this invention describes the use of the adenovirus serotype of the present invention to overcome natural existing or induced, neutralising host activity towards adenoviruses administered in vivo for therapeutic applications. The need for a new serotype is stressed by observations that 1) repeated systemic delivery of recombinant adenovirus serotype 5 is unsuccessful due to formation of high titers of neutralising antibodies against the recombinant adenovirus serotype 5 (Schulick et al, 1997), and 2) pre-existing or humoral immunity is widespread in the population.

[0041] In another aspect this invention provides the use of gene delivery vehicles of the invention or the use of adenovirus serotype 35 for vaccination purposes. Such use prevents at least in part undesired immune responses of the host. Non-limiting examples of undesired immune responses are evoking an immune response against the gene delivery vehicle or adenovirus serotype 35 and/or boosting of an immune response against the gene delivery vehicle or adenovirus serotype 35.

[0042] In another aspect of the invention, alternating use is made of Ad vectors belonging to different subgroups. This aspect of the invention therefore circumvents the inability to repeat the administration of an adenovirus for gene therapy purposes.

## Example 1

A high throughput assay for the detection of neutralising activity in human serum

- 5 [0043] To enable screening of a large amount of human sera for the presence of neutralising antibodies against all adenovirus serotypes, an automated 96-wells assay was developed.

*Human sera*

- 10 [0044] A panel of 100 individuals was selected. Volunteers (50% male, 50% female) were healthy individuals between 20 and 60 years old with no restriction for race. All volunteers signed an informed consent form. People professionally involved in adenovirus research were excluded.

- [0045] Approximately 60 ml blood was drawn in dry tubes. Within two hours after sampling, the blood was centrifuged at 2500 rpm for 10 minutes. Approximately 30 ml serum was transferred to polypropylene tubes and stored frozen at -20°C until further use.

- 15 [0046] Serum was thawed and heat-inactivated at 56°C for 10 minutes and then aliquotted to prevent repeated cycles of freeze/thawing. Part was used to make five steps of twofold dilutions in medium (DMEM, Gibco BRL) in a quantity enough to fill out approximately 70 96-well plates. Aliquots of undiluted and diluted sera were pipetted in deep well plates (96-well format) and using a programmed platemate dispensed in 100 µl aliquots into 96-well plates. This way the plates were loaded with eight different sera in duplo (100 µl/well) according to the scheme below:

25

S1/ 2	S1/ 4	S1/ 8	S1/ 16	S1/ 32	S5/ 2	S5/ 4	S5/ 8	S5/ 16	S5/ 32	-	-
S1/ 2	S1/ 4	S1/ 8	S1/ 16	S1/ 32	S5/ 2	S5/ 4	S5/ 8	S5/ 16	S5/ 32	-	-
S2/ 2	S2/ 4	S2/ 8	S2/ 16	S2/ 32	S6/ 2	S6/ 4	S6/ 8	S6/ 16	S6/ 32	-	-
S2/ 2	S2/ 4	S2/ 8	S2/ 16	S2/ 32	S6/ 2	S6/ 4	S6/ 8	S6/ 16	S6/ 32	-	-
S3/ 2	S3/ 4	S3/ 8	S3/ 16	S3/ 32	S7/ 2	S7/ 4	S7/ 8	S7/ 16	S7/ 32	-	-
S3/ 2	S3/ 4	S3/ 8	S3/ 16	S3/ 32	S7/ 2	S7/ 4	S7/ 8	S7/ 16	S7/ 32	-	-
S4/ 2	S4/ 4	S3/ 8	S3/ 16	S3/ 32	S8/ 2	S8/ 4	S8/ 8	S8/ 16	S8/ 32	-	-
S4/ 2	S4/ 4	S3/ 8	S3/ 16	S3/ 32	S8/ 2	S8/ 4	S8/ 8	S8/ 16	S8/ 32	-	-

35

- [0047] Where S1/2 to S8/2 in columns 1 and 6 represent 1x diluted sera and Sx/4, Sx/8, Sx/16 and Sx/32 the two-fold serial dilutions. The last plates also contained four wells filled with 100µl foetal calf serum as a negative control. Plates were kept at -20°C until further use.

*Preparation of human adenovirus stocks*

- 40 [0048] Prototypes of all known human adenoviruses were inoculated on T25 flasks seeded with PER.C6 cells (ECACC deposit number 96022940) (Fallaux *et al.*, 1998) and harvested upon full CPE. After freeze/thawing 1-2 ml of the crude lysates was used to inoculate a T80 flask with PER.C6 cells (ECACC deposit number 96022940) and virus was harvested at full CPE. The timeframe between inoculation and occurrence of CPE as well as the amount of virus needed to re-infect a new culture, differed between serotypes. Adenovirus stocks were prepared by freeze/thawing and used to inoculate 3-4 T175 cm<sup>2</sup> three-layer flasks with PER.C6 cells (ECACC deposit number 96022940). Upon occurrence of CPE, cells were harvested by tapping the flask, pelleted and virus was isolated and purified by a two step CsCl gradient as follows. Cell pellets were dissolved in 50 ml 10 mM NaPO<sub>4</sub> buffer (pH 7.2) and frozen at -20°C. After thawing at 37°C, 5.6 ml sodium deoxycholate (5% w/v) was added. The solution was mixed gently and incubated for 5-15 minutes at 37°C to completely lyse the cells. After homogenizing the solution, 1875 µl 1M MgCl<sub>2</sub> was added. After the addition of 375 µl DNase (10 mg/ml) the solution was incubated for 30 minutes at 37°C. Cell debris was removed by centrifugation at 1880xg for 30 minutes at PT without brake. The supernatant was subsequently purified from proteins by extraction with freon (3x). The cleared supernatant was loaded on a 1M Tris/HCl buffered cesiumchloride blockgradient (range: 1.2/1.4 gr/ml) and centrifugated at 21000 rpm for 2.5 hours at 10°C. The virus band is isolated after which a second purification using a 1M Tris/HCl buffered continuous gradient of 1.33 gr/ml of cesiumchloride was performed. The virus was then centrifuged for 17 hours at 55000 rpm at 10°C. The virus band is isolated and sucrose (50 % w/v)

is added to a final concentration of 1%. Excess cesiumchloride is removed by dialysis (three times 1 hr at RT) in dialysis slides (Slide-a-lizer, cut off 10000 kDa, Pierce, USA) against 1.5 ltr PBS supplemented with  $\text{CaCl}_2$  (0.9 mM),  $\text{MgCl}_2$  (0.5mM) and an increasing concentration of sucrose (1, 2, 5%). After dialysis, the virus is removed from the slide-a-lizer after which it is aliquoted in portions of 25 and 100  $\mu\text{l}$  upon which the virus is stored at  $-85^\circ\text{C}$ .

5 [0049] To determine the number of virus particles per milliliter, 50  $\mu\text{l}$  of the virus batch is run on a high-pressure liquid chromatograph (HPLC) as described by Shabram et al (1997).

[0050] Viruses were eluted using an NaCl gradient ranging from 0 to 600 mM. As depicted in table I, the NaCl concentration by which the viruses were eluted differed significantly among serotypes.

10 [0051] Most human adenoviruses replicated well on PER.C6 cells ((ECACC deposit number 96022940) with a few exceptions. Adenovirus type 8 and 40 were grown on 911-E4 cells (He et al., 1998). Purified stocks contained between  $5 \times 10^{10}$  and  $5 \times 10^{12}$  virus particles/ml (VP/ml; see table I).

#### *Titration of purified human adenovirus stocks*

15 [0052] Adenoviruses were titrated on PER.C6 cells (ECACC deposit number 96022940) to determine the amount of virus necessary to obtain full CPE in five days, the length of the neutralisation assay. Hereto, 100  $\mu\text{l}$  medium was dispensed into each well of 96-well plates. 25  $\mu\text{l}$  of adenovirus stocks prediluted  $10^4$ ,  $10^5$ ,  $10^6$  or  $10^7$  times were added to column 2 of a 96-well plate and mixed by pipetting up and down 10 times. Then 25  $\mu\text{l}$  was brought from column 2 to column 3 and again mixed. This was repeated until column 11 after which 25  $\mu\text{l}$  from column 11 was discarded. This way serial dilutions in steps of 5 were obtained starting off from a prediluted stock. Then  $3 \times 10^4$  PER.C6 cells (ECACC deposit number 96022940) were added in a 100  $\mu\text{l}$  volume and the plates were incubated at  $37^\circ\text{C}$ , 5%  $\text{CO}_2$  for five or six days. CPE was monitored microscopically. The method of Reed and Muensch was used to calculate the cell culture inhibiting dose 50% (CCID50).

20 [0053] In parallel identical plates were set up that were analysed using the MTT assay (Promega). In this assay living cells are quantified by colorimetric staining. Hereto, 20  $\mu\text{l}$  MTT (7.5 mgr/ml in PBS) was added to the wells and incubated at  $37^\circ\text{C}$ , 5%  $\text{CO}_2$  for two hours. The supernatant was removed and 100  $\mu\text{l}$  of a 20:1 isopropanol/triton-X100 solution was added to the wells. The plates were put on a 96-wells shaker for 3-5 minutes to solubilise precipitated staining.

30 [0054] Absorbance was measured at 540 nm and at 690 nm (background). By this assay wells with proceeding CPE or full CPE can be distinguished.

#### *Neutralisation assay*

35 [0055] 96-well plates with diluted human serum samples were thawed at  $37^\circ\text{C}$ , 5%  $\text{CO}_2$ . Adenovirus stocks diluted to 200 CCID50 per 50  $\mu\text{l}$  were prepared and 50  $\mu\text{l}$  aliquots were added to columns 1-11 of the plates with serum. Plates were incubated for 1 hour at  $37^\circ\text{C}$ , 5%  $\text{CO}_2$ . Then 50  $\mu\text{l}$  PER.C6 cells (ECACC deposit number 96022940) at  $6 \times 10^5/\text{ml}$  were dispensed in all wells and incubated for 1 day at  $37^\circ\text{C}$ , 5%  $\text{CO}_2$ . Supernatant was removed using fresh pipet tips for each row and 200  $\mu\text{l}$  fresh medium was added to all wells to avoid toxic effects of the serum. Plates were incubated for another 4 days at  $37^\circ\text{C}$ , 5%  $\text{CO}_2$ . In addition, parallel control plates were set up in duplo with diluted positive control sera generated in rabbits and specific for each serotype to be tested in rows A and B and with negative control serum (FCS) in rows C and D. Also, in each of the rows E-H a titration was performed as described above with steps of five times dilutions starting with 200 CCID50 of each virus to be tested. On day 5 one of the control plates was analysed microscopically and with the MTT assay. The experimental titer was calculated from the control titration plate observed microscopically.

45 [0056] If CPE was found to be complete, i.e. the first dilution in the control titration experiment analysed by MTT shows clear cell death, all assay plates were processed. If not, the assay was allowed to proceed for one or more days until full CPE was apparent after which all plates were processed. In most cases the assay was terminated at day 5. For Ad1, 5, 33, 39, 42 and 43 the assay was left for six days and for Ad2 for eight days.

50 [0057] A serum sample is regarded to be non-neutralising when at the highest serum concentration a maximum protection is seen of 40% compared to the controls without serum.

[0058] The results of the analysis of 44 prototype adenoviruses against serum from 100 healthy volunteers is shown in figure 1. As expected the percentage of serum samples that contained neutralising antibodies to Ad2 and Ad5 was very high. This was also true for most of the lower numbered adenoviruses. Surprisingly, none of the serum samples contained neutralising antibodies to adenovirus serotype 35.

55 [0059] Also, the number of individuals with neutralising antibody titers to the serotypes 26, 34 and 48 was very low.

[0060] Therefor, recombinant E1-deleted adenoviruses based on Ad35 or one of the other above mentioned serotypes have an important advantage compared to recombinant vectors based on Ad5 with respect to clearance of the viruses by neutralising antibodies.

[0061] Also, Ad5-based vectors that have (parts of) the capsid proteins involved in immunogenic response of the host replaced by the corresponding (parts of) the capsid proteins of Ad35 or one of the other serotypes will be less, or even not, neutralised by the vast majority of human sera.

5 [0062] As can be seen in table I the VP/CCID50 ratio calculated from the virus particles per ml and the CCID50 obtained for each virus in the experiments was highly variable and ranged from 0.4 to 5 log. This is probably caused by different infection efficiencies of PER.C6 cells (ECACC deposit number 96022940) and by differences in replication efficiency of the viruses. Furthermore, differences in batch qualities may play a role. A high VP/CCID50 ratio means that more virus was put in the wells to obtain CPE in 5 days. As a consequence the outcome of the neutralisation study might be biased since more (inactive) virus particles could shield the antibodies. To check whether this phenomenon  
10 had taken place, the VP/CCID50 ratio was plotted against the percentage of serum samples found positive in the assay (Figure 2). The graph clearly shows that there is no negative correlation between the amount of viruses in the assay and neutralisation in serum.

## Example 2

15

### Generation of Ad5 plasmid vectors for the production of recombinant viruses and easy manipulation of adenoviral genes

#### *pBr/Ad.Bam-rITR (ECACC deposit P97082122)*

20

[0063] In order to facilitate blunt end cloning of the ITR sequences, wild-type human adenovirus type 5 (Ad5) DNA was treated with Klenow enzyme in the presence of excess dNTPs. After inactivation of the Klenow enzyme and purification by phenol/chloroform extraction followed by ethanol precipitation, the DNA was digested with BamHI. This DNA preparation was used without further purification in a ligation reaction with pBr322 derived vector DNA prepared as follows: pBr322 DNA was digested with EcoRV and BamHI, dephosphorylated by treatment with TSAP enzyme (Life Technologies) and purified on LMP agarose gel (SeaPlaque GTG). After transformation into competent *E.coli* DH5 $\alpha$  (Life Techn.) and analysis of ampicillin resistant colonies, one clone was selected that showed a digestion pattern as  
25 expected for an insert extending from the BamHI site in Ad5 to the right ITR.

[0064] Sequence analysis of the cloning border at the right ITR revealed that the most 3' G residue of the ITR was  
30 missing, the remainder of the ITR was found to be correct. Said missing G residue is complemented by the other ITR during replication.

#### *pBr/Ad.Sal-rITR (ECACC deposit P97082119)*

35 [0065] pBr/Ad.Bam-rITR was digested with BamHI and Sall. The vector fragment including the adenovirus insert was isolated in LMP agarose (SeaPlaque GTG) and ligated to a 4.8 kb Sall-BamHI fragment obtained from wt Ad5 DNA and purified with the GeneClean II kit (Bio 101, Inc.). One clone was chosen and the integrity of the Ad5 sequences was determined by restriction enzyme analysis. Clone pBr/Ad.Sal-rITR contains adeno type 5 sequences from the Sall site at bp 16746 up to and including the rITR (missing the most 3' G residue).

40

#### *pBr/Ad.Cla-Bam (ECACC deposit P97082117)*

[0066] wt Adeno type 5 DNA was digested with ClaI and BamHI, and the 20.6 kb fragment was isolated from gel by electro-elution. pBr322 was digested with the same enzymes and purified from agarose gel by GeneClean. Both fragments were ligated and transformed into competent DH5 $\alpha$ . The resulting clone pBr/Ad.Cla-Bam was analysed by  
45 restriction enzyme digestion and shown to contain an insert with adenovirus sequences from bp 919 to 21566.

#### *pBr/Ad.AfIII-Bam (ECACC deposit P97082114)*

50 [0067] Clone pBr/Ad.Cla-Bam was linearized with EcoRI (in pBr322) and partially digested with AfIII. After heat inactivation of AfIII for 20' at 65°C the fragment ends were filled in with Klenow enzyme. The DNA was then ligated to a blunt double stranded oligo linker containing a PacI site (5'-AATTGTCTTAATTAACCGCTTAA-3'). This linker was made by annealing the following two oligonucleotides: 5'-AATTGTCTTAATTAACCGC-3' and 5'-AATTGCGGTTAATTAAGAC-3', followed by blunting with Klenow enzyme. After precipitation of the ligated DNA to change buffer, the ligations were  
55 digested with an excess PacI enzyme to remove concatameres of the oligo. The 22016 bp partial fragment containing Ad5 sequences from bp 3534 up to 21566 and the vector sequences, was isolated in LMP agarose (SeaPlaque GTG), religated and transformed into competent DH5 $\alpha$ . One clone that was found to contain the PacI site and that had retained the large adeno fragment was selected and sequenced at the 5' end to verify correct insertion of the PacI linker



in the (lost) AflII site.

*pBr/Ad.Bam-r.ITRpac#2 (ECACC deposit P97082120) and pBr/Ad.Bam-r.ITRpac#8 (ECACC deposit P97082121)*

5 [0068] To allow insertion of a PacI site near the ITR of Ad5 in clone pBr/Ad.Bam-r.ITR about 190 nucleotides were removed between the ClaI site in the pBr322 backbone and the start of the ITR sequences. This was done as follows: pBr/Ad.Bam-r.ITR was digested with ClaI and treated with nuclease Bal31 for varying lengths of time (2', 5', 10' and 15'). The extend of nucleotide removal was followed by separate reactions on pBr322 DNA (also digested at the ClaI site), using identical buffers and conditions. Bal31 enzyme was inactivated by incubation at 75 °C for 10', the DNA was pre-

10 cipitated and resuspended in a smaller volume TE buffer.

[0069] To ensure blunt ends, DNAs were further treated with T4 DNA polymerase in the presence of excess dNTPs. After digestion of the (control) pBr322 DNA with SalI, satisfactory degradation (~150 bp) was observed in the samples created for 10' or 15'. The 10' or 15' treated pBr/Ad.Bam-r.ITR samples were then ligated to the above described blunted PacI linkers (See pBr/Ad.AflII-Bam). Ligations were purified by precipitation, digested with excess PacI and separated

15 from the linkers on an LMP agarose gel. After religation, DNAs were transformed into competent DH5α and colonies analyzed. Ten clones were selected that showed a deletion of approximately the desired length and these were further analyzed by T-track sequencing (T7 sequencing kit, Pharmacia Biotech). Two clones were found with the PacI linker inserted just downstream of the r.ITR. After digestion with PacI, clone #2 has 28 bp and clone #8 has 27 bp attached to the ITR.

20

*pWE/Ad.AflII-r.ITR (ECACC deposit P97082116)*

[0070] Cosmid vector pWE15 (Clontech) was used to clone larger Ad5 inserts. First, a linker containing a unique PacI site was inserted in the EcoRI sites of pWE15 creating pWE.pac. To this end, the double stranded PacI oligo as

25

described for pBr/Ad.AflII-BamHI was used but now with its EcoRI protruding ends. The following fragments were then isolated by electro-elution from agarose gel: pWE.pac digested with PacI, pBr/AflII-Bam digested with PacI and BamHI and pBr/Ad.Bam-r.ITR#2 digested with BamHI and PacI. These fragments were ligated together and packaged using λ phage packaging extracts (Stratagene) according to the manufacturer's protocol. After infection into host bacteria, colonies were grown on plates and analyzed for presence of the complete insert. pWE/Ad.AflII-r.ITR contains all adenovirus

30

*pBr/Ad.IITR-Sal(9.4) (ECACC deposit P97082115)*

[0071] Adeno 5 wt DNA was treated with Klenow enzyme in the presence of excess dNTPs and subsequently

35

digested with SalI. Two of the resulting fragments, designated left ITR-Sal(9.4) and Sal(16.7)-right ITR, respectively, were isolated in LMP agarose (Seaplaque GTG). pBr322 DNA was digested with EcoRV and SalI and treated with phosphatase (Life Technologies). The vector fragment was isolated using the GeneClean method (BIO 101, Inc.) and ligated to the Ad5 SalI fragments. Only the ligation with the 9.4 kb fragment gave colonies with an insert. After analysis and sequencing of the cloning border a clone was chosen that contained the full ITR sequence and extended to the SalI

40

*pBr/Ad.IITR-Sal(16.7) (ECACC deposit P97082118)* pBr/Ad.IITR-Sal(9.4) is digested with SalI and dephosphorylated (TSAP, Life Technologies). To extend this clone upto the third SalI site in Ad5, pBr/Ad.Cla-Bam was linearized with BamHI and partially digested with SalI. A 7.3 kb SalI fragment containing adenovirus sequences from 9462-16746 was

45

*pWE/Ad.AflII-EcoRI*

[0072] pWE.pac was digested with ClaI and 5' protruding ends were filled using Klenow enzyme. The DNA was then digested with PacI and isolated from agarose gel. pWE/AflII-r.ITR was digested with EcoRI and after treatment with Klenow enzyme digested with PacI. The large 24 kb fragment containing the adenoviral sequences was isolated from agarose gel and ligated to the ClaI-digested and blunted pWE.pac vector using the Ligation Express™ kit from Clontech. After transformation of Ultracompetent XL10-Gold cells from Stratagene, clones were identified that contained the expected insert. pWE/AflII-EcoRI contains Ad5 sequences from bp 3534-27336.

55

Generation of *pWE/Ad.AflII-r.ITRsp*

[0073] The 3' ITR an the vector pWE/Ad.AflII-r.ITR does not include the terminal G-nucleotide. Furthermore, the



PacI site is located almost 30 bp from the right ITR. Both these characteristics may decrease the efficiency of virus generation due to inefficient initiation of replication at the 3' ITR. Note that during virus generation the left ITR in the adapter plasmid is intact and enables replication of the virus DNA after homologous recombination.

[0074] To improve the efficiency of initiation of replication at the 3' ITR, the pWE/Ad.AfIII-rITR was modified as follows: construct pBr/Ad.Bam-rITRpac#2 was first digested with PacI and then partially digested with AvrII and the 17.8 kb vector containing fragment was isolated and dephosphorylated using SAP enzyme (Boehringer Mannheim). This fragment lacks the adenosequences from nucleotide 35464 to the 3'ITR. Using DNA from pWE/Ad.AfIII-rITR as template and the primers ITR-EPH:

10 5'-CGG AAT TCT TAA TTA AGT TAA CAT CAT CAA TAA TAT ACC-3'  
and  
Ad101: 5'-TGA TTC ACA TCG GTC AGT GC-3'

15 a 630 bp PCR fragment was generated corresponding to the 3' Ad5 sequences. This PCR fragment was subsequently cloned in the vector pCR2.1 (Invitrogen) and clones containing the PCR fragment were isolated and sequenced to check correct amplification of the DNA. The PCR clone was then digested with PacI and AvrII and the 0.5 kb adeno insert was ligated to the PacI/ partial AvrII digested pBr/Ad.Bam-rITRpac#2 fragment generating pBr/Ad.Bam-rITRsp. Next this construct was used to generate a cosmid clone (as described above) that has an insert corresponding to the adenosequences 3534 to 35938. This clone was named pWE/AfIII-rITRsp.

20 Generation of *pWE/Ad.AfIII-rITRΔE2A*:

[0075] Deletion of the E2A coding sequences from pWE/Ad.AfIII-rITR (ECACC deposit P97082116) has been accomplished as follows.

25 [0076] The adenoviral sequences flanking the E2A coding region at the left and the right site were amplified from the plasmid pBr/Ad.Sal.rITR (ECACC deposit P97082119) in a PCR reaction with the Expand PCR system (Boehringer) according to the manufacturers protocol. The following primers were used: Right flanking sequences (corresponding Ad5 nucleotides 24033 to 25180):

30 ΔE2A.SnaBI: 5'-GGC GTA CGT AGC CCT GTC GAA AG-3'  
ΔE2A.DBP-start: 5'-CCA ATG CAT TCG AAG TAC TTC CTT CTC CTA TAG GC-3'

[0077] The amplified DNA fragment was digested with SnaBI and NsiI (NsiI site is generated in the primer ΔE2A.DBP-start, underlined).

35 [0078] Left flanking sequences (corresponding Ad5 nucleotides 21557 to 22442):

ΔE2A.DBP-stop: 5'-CCA ATG CAT ACG GCG CAG ACG G-3'  
ΔE2A.BamHI: 5'-GAG GTG GAT CCC ATG GAC GAG-3'

40 [0079] The amplified DNA was digested with BamHI and NsiI (NsiI site is generated in the primer ΔE2A.DBP-stop, underlined).

[0080] Subsequently, the digested DNA fragments were ligated into SnaBI/BamHI digested pBr/Ad.Sal-rITR. Sequencing confirmed the exact replacement of the DBP coding region with a unique NsiI site in plasmid pBr/Ad.Sal-rITRΔE2A. The unique NsiI site can be used to introduce an expression cassette for a gene to be transduced by the recombinant vector.

45 [0081] The deletion of the E2A coding sequences was performed such that the splice acceptor sites of the 100K encoding L4-gene at position 24048 in the top strand was left intact. In addition, the poly adenylation signals of the original E2A-RNA and L3-RNAs at the left hand site of the E2A coding sequences were left intact. This ensures proper expression of the L3-genes and the gene encoding the 100K L4-protein during the adenovirus life cycle.

50 [0082] Next, the plasmid pWE/Ad.AfIII-rITRΔE2A was generated. The plasmid pBr/Ad.Sal-rITRΔE2A was digested with BamHI and SpeI. The 3.9-Kb fragment in which the E2A coding region was replaced by the unique NsiI site was isolated. The pWE/Ad.AfIII-rITR was digested with BamHI and SpeI. The 35 Kb DNA fragment, from which the BamHI/SpeI fragment containing the E2A coding sequence was removed, was isolated. The fragments were ligated and packaged using λ phage-packaging extracts according to the manufacturer protocol (Stratagene), yielding the plasmid pWE/Ad.AfIII-rITRΔE2A.

55 [0083] This cosmid clone can be used to generate adenoviral vectors that are deleted for E2A by cotransfection of PacI digested DNA together with digested adapter plasmids onto packaging cells that express functional E2A gene product.

*Construction of adapter plasmids*

[0084] The absence of sequence overlap between the recombinant adenovirus and E1 sequences in the packaging cell line is essential for safe, RCA-free generation and propagation of new recombinant viruses. The adapter plasmid pMLPI.TK (described in PCT/NL96/00244) is an example of an adapter plasmid designed for use according to the invention in combination with the improved packaging cell lines of the invention. This plasmid was used as the starting material to make a new vector in which nucleic acid molecules comprising specific promoter and gene sequences can be easily exchanged.

[0085] First, a PCR fragment was generated from pZipΔMo+PyF101(N<sup>-</sup>) template DNA (described in PCT/NL96/00195) with the following primers: LTR-1: 5'-CTG TAC GTA CCA GTG CAC TGG CCT AGG CAT GGA AAA ATA CAT AAC TG-3' and LTR-2: 5'-GCG GAT CCT TCG AAC CAT GGT AAG CTT GGT ACC GCT AGC GTT AAC CGG GCG ACT CAG TCA ATC G-3'. Pwo DNA polymerase (Boehringer Mannheim) was used according to manufacturers protocol with the following temperature cycles: once 5' at 95°C; 3' at 55°C; and 1' at 72°C, and 30 cycles of 1' at 95°C, 1' at 60°C, 1' at 72°C, followed by once 10' at 72°C. The PCR product was then digested with BamHI and ligated into pMLP10 (Levrero *et al.*, 1991) vector digested with PvuII and BamHI, thereby generating vector pLTR10. This vector contains adenoviral sequences from bp 1 up to bp 454 followed by a promoter consisting of a part of the Mo-MuLV LTR having its wild-type enhancer sequences replaced by the enhancer from a mutant polyoma virus (PyF101). The promoter fragment was designated L420. Next, the coding region of the murine HSA gene was inserted. pLTR10 was digested with BstBI followed by Klenow treatment and digestion with NcoI. The HSA gene was obtained by PCR amplification on pUC18-HSA (Kay *et al.*, 1990) using the following primers: HSA1, 5'-GCG CCA CCA TGG GCA GAG CGA TGG TGG C-3' and HSA2, 5'-GTT AGA TCT AAG CTT GTC GAC ATC GAT CTA CTA ACA GTA GAG ATG TAG AA-3'. The 269 bp amplified fragment was subcloned in a shuttle vector using the NcoI and BglII sites. Sequencing confirmed incorporation of the correct coding sequence of the HSA gene, but with an extra TAG insertion directly following the TAG stop codon. The coding region of the HSA gene, including the TAG duplication was then excised as a NcoI(sticky)-Sall(blunt) fragment and cloned into the 3.5 kb NcoI(sticky)/BstBI(blunt) fragment from pLTR10, resulting in pLTR-HSA10.

[0086] Finally, pLTR-HSA10 was digested with EcoRI and BamHI after which the fragment containing the left ITR, packaging signal, L420 promoter and HSA gene was inserted into vector pMLPI.TK digested with the same enzymes and thereby replacing the promoter and gene sequences. This resulted in the new adapter plasmid pAd/L420-HSA that contains convenient recognition sites for various restriction enzymes around the promoter and gene sequences. SnaBI and AvrII can be combined with HpaI, NheI, KpnI, HindIII to exchange promoter sequences, while the latter sites can be combined with the ClaI or BamHI sites 3' from HSA coding region to replace genes in this construct.

[0087] Another adapter plasmid that was designed to allow easy exchange of nucleic acid molecules was made by replacing the promoter, gene and poly A sequences in pAd/L420-HSA with the CMV promoter, a multiple cloning site, an intron and a poly-A signal. For this purpose, pAd/L420-HSA was digested with AvrII and BglII followed by treatment with Klenow to obtain blunt ends. The 5.1 kb fragment with pBr322 vector and adenoviral sequences was isolated and ligated to a blunt 1570 bp fragment from pcDNA1/amp (Invitrogen) obtained by digestion with HhaI and AvrII followed by treatment with T4 DNA polymerase. This adapter plasmid was named pAd5/CLIP. To enable removal of vector sequences from the left ITR in pAd5/Clip, this plasmid was partially digested with EcoRI and the linear fragment was isolated. An oligo of the sequence 5' TTAAGTCGAC-3' was annealed to itself resulting in a linker with a Sall site and EcoRI overhang. The linker was ligated to the partially digested pAd5/Clip vector and clones were selected that had the linker inserted in the EcoRI site 23 bp upstream of the left adenovirus ITR in pAd5/Clip resulting in pAd5/Clipsal. Likewise, the EcoRI site in pAd5/Clip has been changed to a PacI site by insertion of a linker of the sequence 5'-AATT-GTCTTAATTAACCGCAATT-3'. The pAd5/Clip vector was partially digested with EcoRI, dephosphorylated and ligated to the PacI linker with EcoRI overhang. The ligation mixture was digested with PacI to remove concatamers, isolated from agarose gel and religated. The resulting vector was named pAd5/Clippac. These changes enable more flexibility to liberate the left ITR from the plasmid vector sequences.

[0088] The vector pAd5/L420-HSA was also modified to create a Sall or PacI site upstream of the left ITR. Hereto pAd5/L420-HSA was digested with EcoRI and ligated to the above described PacI linker. The ligation mixture was digested with PacI and religated after isolation of the linear DNA from agarose gel to remove concatamerised linkers. This resulted in adapter plasmid pAd5/L420-HSApac. This construct was used to generate pAd5/L420-HSAsal as follows: pAd5/L420-HSApac was digested with Scal and BsrGI and the vector fragment was ligated to the 0.3 kb fragment isolated after digestion of pAd5/Clipsal with the same enzymes.

*Generation of adapter plasmids pAdMire and pAdApt*

[0089] To create an adapter plasmid that only contains a polylinker sequence and no promoter or polyA sequences, pAd5/L420-HSApac was digested with AvrII and BglII. The vector fragment was ligated to a linker oligonucleotide

digested with the same restriction enzymes. The linker was made by annealing oligos of the following sequence:

PLL-1: 5'- GCC ATC CCT AGG AAG CTT GGT ACC GGT GAA TTC GCT AGC GTT AAC GGA TCC TCT AGA CGA GAT CTG G-3' and

PLL-2: 5'- CCA GAT CTC GTC TAG AGG ATC CGT TAA CGC TAG CGA ATT CAC CGG TAC CAA GCT TCC TAG GGA TGG C-3'.

[0090] The annealed linkers were digested with AvrII and BglII and separated from small ends by column purification (Qiaquick nucleotide removal kit) according to manufacturers recommendations. The linker was then ligated to the AvrII/BglII digested pAd5/L420-HSApac fragment. A clone, named AdMire, was selected that had the linker incorporated and was sequenced to check the integrity of the insert.

[0091] Adapter plasmid AdMire enables easy insertion of complete expression cassettes.

[0092] An adapter plasmid containing the human CMV promoter that mediates high expression levels in human cells was constructed as follows: pAd5/L420-HSApac was digested with AvrII and 5' protruding ends were filled in using Klenow enzyme. A second digestion with HindIII resulted in removal of the L420 promoter sequences. The vector fragment was isolated and ligated to a PCR fragment containing the CMV promoter sequence. This PCR fragment was obtained after amplification of CMV sequences from pCMVLacI (Stratagene) with the following primers:

CMVplus: 5'-GATCGGTACCACTGCAGTGGTCAATATTGGCCATTAGCC-3' and

CMVminA: 5'-GATCAAGCTTCCAATGCACCGTTCCCGGC-3'.

[0093] The PCR fragment was first digested with PstI (underlined in CMVplus) after which the 3'-protruding ends were removed by treatment with T4 DNA polymerase. Then the DNA was digested with HindIII (underlined in CMVminA) and ligated into the above described pAd5/L420-HSApac vector fragment digested with AvrII and HindIII. The resulting plasmid was named pAd5/CMV-HSApac. This plasmid was then digested with HindIII and BamHI and the vector fragment was isolated and ligated to the polylinker sequence obtained after digestion of AdMire with HindIII and BglII. The resulting plasmid was named pAdApt. Adapter plasmid pAdApt contains nucleotides - 735 to +95 of the human CMV promoter (Boshart et al., 1985). A second version of this adapter plasmid containing a Sall site in place of the PacI site upstream of the left ITR was made by inserting the 0.7 kb Scal-BsrGI fragment from pAd5/Clipsal into pAdApt digested with Scal and partially digested with BsrGI. This clone was named pAdApt.sal.

#### *Generation of recombinant adenoviruses based on Ad5*

[0094] RCA free recombinant adenoviruses can be generated very efficiently using the above described adapter plasmids and the pWe/Ad.AfIII-rITR or pWE/Ad.AfIII-rITRsp constructs.

[0095] Generally, the adapter plasmid containing the desired transgene in the desired expression cassette is digested with suitable enzymes to liberate the insert from vector sequences at the 3' and/or at the 5' end. The adenoviral complementation plasmids pWE/Ad.AfIII-rITR or pWE/Ad.AfIII-rITRsp are digested with PacI to liberate the adeno sequences from the vector plasmids. As a non-limiting example the generation of AdApt-LacZ is described. Adapter plasmid pAdApt-LacZ was generated as follows. The E.coli LacZ gene was amplified from the plasmid pMLP.nlsLacZ (EP 95-202 213) by PCR with the primers 5'-GGGGTGGCCAGGGTACCTCTAGGCTTTTGCAA-3' and 5'-GGGGGGATCCATAAACAAGTTCAGAATCC-3'. The PCR reaction was performed with Ex Tag (Takara) according to the suppliers protocol at the following amplification program: 5 minutes 94°C, 1 cycle; 45 seconds 94°C and 30 seconds 60°C and 2 minutes 72°C, 5 cycles; 45 seconds 94°C and 30 seconds 65°C and 2 minutes 72°C, 25 cycles; 10 minutes 72°C, 1 cycle. The PCR product was subsequently digested with KpnI and BamHI and the digested DNA fragment was ligated into KpnI/BamHI digested pcDNA3 (Invitrogen), giving rise to pcDNA3.nlsLacZ. Construct pcDNA3.nlsLacZ was then digested with KpnI and BamHI and the 3 kb LacZ fragment was isolated from gel using the gene clean spin kit (Bio 101, Inc.).

[0096] pAdApt was also digested with KpnI and BamHI and the linear vector fragment was isolated from gel as above. Both isolated fragments were ligated and one clone containing the LacZ insert was selected. Construct pAdApt-LacZ was digested with Sall, purified by the gene clean spin kit and subsequently digested with PacI. pWE/Ad.AfIII-rITRsp was digested with PacI. Both digestion mixtures were treated for 30' by 65 °C to inactivate the enzymes. Samples were put on gel to estimate the concentration.  $2.5 \times 10^6$  PER.C6 cells (ECACC deposit number 96022940) were seeded in T25 flasks in DMEM with 10% FCS and 10mM MgCl. The next day four microgram of each plasmid was transfected into PER.C6 cells (ECACC deposit number 96022940) using lipofectamine transfection reagent (Life Technologies Inc.) according to instructions of the manufacturer. The next day the medium was replaced by fresh culture medium and cells were further cultured at 37° C, 10% CO<sub>2</sub>. Again 24 hrs. later cells were trypsinised, seeded into T80 flasks and cultured at 37°C, 10% CO<sub>2</sub>. Full CPE was obtained 6 days after seeding in the T80 flask. Cells were har-

vested in the medium and subjected to one freeze/thaw cycle. The crude lysate obtained this way was used to plaque purify the mixture of viruses. Ten plaques were picked, expanded in a 24 well plate and tested for LacZ expression following infection of A549 cells. Viruses from all ten plaques expressed LacZ.

### 5 Example 3

#### Generation of chimeric recombinant adenoviruses

##### *Generation of hexon chimeric Ad5-based adenoviruses*

10

[0097] Neutralising antibodies in human serum are mainly directed to the hexon protein and to a lesser extent to the penton protein. Hexon proteins from different serotypes show highly variable regions present in loops that are predicted to be exposed at the outside of the virus (Athappilly et al., 1994; J. Mol. Biol. 242, 430-455). Most type specific epitopes have been mapped to these highly variable regions (Toogood et al., 1989; J. Gen Virol. 70, 3203-3214). Thus replacement of (part of) the hexon sequences with corresponding sequences from a different serotype is an effective strategy to circumvent (pre-existing) neutralising antibodies to Ad5. Hexon coding sequences of adenovirus serotype 5 are located between nucleotides 18841 and 21697.

[0098] To facilitate easy exchange of hexon coding sequences from alternative adenovirus serotypes into the adenovirus serotype 5 backbone, first a shuttle vector was generated.

20

[0099] This subclone, coded pBr/Ad.Eco-PmeI, was generated by first digesting plasmid pBr322 with EcoRI and EcoRV and inserting the 14 kb PmeI-EcoRI fragment from pWE/Ad.AfIII-Eco. In this shuttle vector a deletion was made of a 1430 bp SmaI fragment by digestion with SmaI and religation to give pBr/Ad.Eco-PmeI ΔSmaI. The removed fragment contains unique SpeI and MunI sites. From pBr/Ad.Eco-PmeIΔSmaI the adenovirus serotype 5 DNA encoding hexon was deleted.

25

[0100] Hereto, the hexon flanking sequences were PCR amplified and linked together thereby generating unique restriction sites replacing the hexon coding region. For these PCR reactions four different oligonucleotides were required: Δhex1-Δhex4.

Δhex1: 5'- CCT GGT GCT CCC AAC AGC-3'

30

Δhex2: 5'- CCG GAT CCA CTA GTG GAA AGC GGG CGC GCG-3'

Δhex3: 5'- CCG GAT CCA ATT GAG NAG CAA GCA ACA TCA ACA AC-3'

Δhex4: 5'- GAG AAG GGC ATG GAG GCT G-3'

35

[0101] The amplified DNA product of ± 1100 bp obtained with oligonucleotides Δhex1 and Δhex2 was digested with BamHI and FseI. The amplified DNA product of ± 1600 bp obtained with oligonucleotides Δhex3 and Δhex4 was digested with BamHI and SbfI. These digested PCR fragments were subsequently purified from agarose gel and in a tri-part ligation reaction using T4 ligase enzyme linked to pBr/Ad.Eco-PmeI ΔSmaI digested with FseI and SbfI. The resulting construct was coded pBr/Ad.Eco-PmeIΔHexon. This construct was sequenced in part to confirm the correct nucleotide sequence and the presence of unique restriction sites MunI and SpeI.

40

[0102] pBr/Ad.Eco-PmeIΔHexon serves as a shuttle vector to introduce heterologous hexon sequences amplified from virus DNA from different serotypes using primers that introduce the unique restriction sites MunI and SpeI at the 5' and 3' ends of the hexon sequences respectively. To generate Ad5-based vectors that contain hexon sequences from the serotypes to which healthy individuals have no, or very low, titers of NAB the hexon sequences of Ad35, Ad34, Ad26 and Ad48 were amplified using the following primers:

45

Hex-up2: 5'-GACTAGTCAAGATGGCYACCCCHTCGATGATG-3' and

Hex-do2: 5'-GCTGGCCAATTGTTATGKGTGCGTTRCCGGC-3'.

50

[0103] These primers were designed using the sequences of published hexon coding regions (for example hexon sequences of Ad2, Ad3, Ad4, Ad5, Ad7, Ad16, Ad40 and Ad41 can be obtained at Genbank). Degenerated nucleotides were incorporated at positions that show variation between serotypes.

[0104] PCR products were digested with SpeI and MunI and cloned into the pBr/Ad.Eco-PmeIΔHexon construct digested with the same enzymes.

55

[0105] The hexon modified sequences were subsequently introduced in the construct pWE/Ad.AfIII-rITR by exchange of the Ascl fragment generating pWE/Ad.AfIII-rITRHexXX where XX stands for the serotype used to amplify hexon sequences.

[0106] The pWE/Ad.AfIII-rITRHexXX constructs are then used to make viruses in the same manner as described above for Ad5 recombinant viruses.



*Generation of penton chimeric Ad5-based recombinant viruses*

[0107] The adenovirus type 5 penton gene is located between sequences 14156 and 15869. Penton base is the adenovirus capsid protein that mediates internalisation of the virus into the target cell. At least some serotypes (type C and B) have been shown to achieve this by interaction of an RGD sequence in penton with integrins on the cell surface.

[0108] However, type F adenoviruses do not have an RGD sequence and for most viruses of the A and D group the penton sequence is not known. Therefore, penton may be involved in target cell specificity. Furthermore, as a capsid protein, the penton protein is involved in the immunogenicity of the adenovirus (Gahery-Segard *et al.*, 1998). Therefore, replacement of Ad5 penton sequences with penton sequences from serotypes to which no or low titers of NAB exist in addition to replacement of the hexon sequences will prevent clearance of the adenoviral vector more efficiently than replacement of hexon alone. Replacement of penton sequences may also affect infection specificity.

[0109] To be able to introduce heterologous penton sequences in Ad5 we made use of the plasmid-based system described above.

[0110] First a shuttle vector for penton sequences was made by insertion of the 7.2 kb *NheI*-*EcoRV* fragment from construct pWE/Ad.AfIII-*EcoRI* into pBr322 digested with the same enzymes. The resulting vector was named pBr/XN. From this plasmid Ad5 penton sequences were deleted and replaced by unique restriction sites that are then used to introduce new penton sequences from other serotypes. Hereto, the left flanking sequences of penton in pBr/XN were PCR amplified using the following primers:

DP5-F: 5'- CTG TTG CTG CTG CTA ATA GC-3' and  
DP5-R: 5'- CGC GGA TCC TGT ACA ACT AAG GGG AAT ACA AG-3'

[0111] DP5-R has an *Bam*HI site (underlined) for ligation to the right flanking sequence and also introduces a unique *Bsr*GI site (bold face) at the 5'-end of the former Ad5 penton region.

[0112] The right flanking sequence was amplified using:

DP3-F: 5'-CGC GGA TCC CTT AAG GCA AGC ATG TCC ATC CTT-3' and  
DP3-3R: 5'- AAA ACA CGT TTT ACG CGT CGA CCT TTC-3'

[0113] DP3-F has an *Bam*HI site (underlined) for ligation to the left flanking sequence and also introduces a unique *Afl*II site (bold face) at the 3'-end of the former Ad5 penton region.

[0114] The two resulting PCR fragments were digested with *Bam*HI and ligated together. Then this ligation mixture was digested with *Avr*II and *Bgl*II. pBr/XN was also digested with *Avr*II and *Bgl*II and the vector fragment was ligated to the digested ligated PCR fragments. The resulting clone was named pBr/Ad.Δpenton. Penton coding sequences from Ad35, Ad34, Ad26 and Ad48 were PCR amplified such that the 5' and 3' ends contained the *Bsr*GI and *Afl*II sites respectively. Hereto, the following primers were used:

For Ad34 and Ad35:

P3-for: 5'-GCT CGA TGT ACA ATG AGG AGA CGA GCC GTG CTA-3'  
P3-rev: 5'-GCT CGA CTT AAG TTA GAA AGT GCG GCT TGA AAG-3'

For Ad26 and Ad48:

P17F: 5'-GCT CGA TGT ACA ATG AGG CGT GCG GTG GTG TCT TC-3'  
P17R: 5'-GCT CGA CTT AAG TTA GAA GGT GCG ACT GGA AAG C-3'

[0115] Amplified pcr products were digested with *Bfr*I and *Bsr*GI and cloned into pBr/Ad.Δpenton digested with the same enzymes.

[0116] Introduction of these heterologous penton sequences in pBr/Ad.Δpenton generated constructs named pBr/Ad.pentonXX where XX represents the number of the serotype corresponding to the serotype used to amplify the inserted penton sequences. Subsequently the new penton sequences were introduced in the a pWE/Ad.AfIII-*r*ITR vector having a modified hexon. For example penton sequences from Ad35 were introduced in the construct pWE/Ad.AfIII-*r*ITRHex35 by exchange of the common *Fse*I fragment. Other combinations of penton and hexon sequences were also made. Viruses with modified hexon and penton sequences were made as described above using cotransfection with an adapter plasmid on PER.C6 cells (ECACC deposit number 96022940). In addition, penton sequences were introduced in the pWE/Ad.AfIII-*r*ITR construct. The latter constructs contain only a modified penton and viruses generated from these constructs will be used to study the contribution of penton sequences to the neutralisation of adenoviruses and

also for analysis of possible changes in infection efficiency and specificity.

#### Generation of fiber chimeric Ad5-based viruses

[0117] Adenovirus infection is mediated by two capsid proteins fiber and penton. Binding of the virus to the cells is achieved by interaction of the protruding fiber protein with a receptor on the cell surface. Internalisation then takes place after interaction of the penton protein with integrins on the cell surface. At least some adenovirus from subgroup C and B have been shown to use a different receptor for cell binding and therefore have different infection efficiencies on different cell types. Thus it is possible to change the infection spectrum of adenoviruses by changing the fiber in the capsid. The fiber coding sequence of adenovirus serotype 5 is located between nucleotides 31042 and 32787. To remove the adenovirus serotype 5 DNA encoding fiber we started with construct pBr/Ad.Bam-rITR. First a NdeI site was removed from this construct. For this purpose, pBr322 plasmid DNA was digested with NdeI after which protruding ends were filled using Klenow enzym. This pBr322 plasmid was then religated, digested with NdeI and transformed into *E.coli* DH5 $\alpha$ . The obtained pBr/ $\Delta$ NdeI plasmid was digested with ScaI and Sall and the resulting 3198 bp vector fragment was ligated to the 15349 bp ScaI-Sall fragment derived from pBr/Ad.BamrITR, resulting in plasmid pBr/Ad.Bam-rITR $\Delta$ NdeI which hence contained a unique NdeI site. Next a PCR was performed with oligonucleotides

NY-up:

5'- CGA CAT ATG TAG ATG CAT TAG TTT GTG TTA TGT TTC AAC GTG-3'

and

NY-down:

5'-GGA GAC CAC TGC CAT GTT-3'

[0118] During amplification, both a NdeI (bold face) and a NsiI restriction site (underlined) were introduced to facilitate cloning of the amplified fiber DNAs. Amplification consisted of 25 cycles of each 45 sec. at 94°C, 1 min. at 60°C, and 45 sec. at 72°C. The PCR reaction contained 25 pmol of oligonucleotides NY-up or NY-down, 2mM dNTP, PCR buffer with 1.5 mM MgCl<sub>2</sub>, and 1 unit of Elongase heat stable polymerase (Gibco, The Netherlands). One-tenth of the PCR product was run on an agarose gel which demonstrated that the expected DNA fragment of  $\pm$  2200 bp was amplified. This PCR fragment was subsequently purified using GeneClean kit system (Bio101 Inc.). Then, both the construct pBr/Ad.Bam-rITR $\Delta$ NdeI as well as the PCR product were digested with restriction enzymes NdeI and SbfI. The PCR fragment was subsequently cloned using T4 ligase enzyme into the NdeI and SbfI digested pBr/Ad.Bam-rITR $\Delta$ NdeI, generating pBr/Ad.BamR $\Delta$ Fib.

[0119] This plasmid allows insertion of any PCR amplified fiber sequence through the unique NdeI and NsiI sites that are inserted in place of the removed fiber sequence. Viruses can be generated by a double homologous recombination in packaging cells described in patent No. PCT/NL96/00244 using an adapter plasmid, construct pBr/Ad.AfIII-EcoRI digested with PacI and EcoRI and a pBr/Ad.BamR $\Delta$ Fib construct in which heterologous fiber sequences have been inserted. To increase the efficiency of virus generation, the construct pBr/Ad.BamR $\Delta$ Fib was modified to generate a PacI site flanking the right ITR. Hereto, pBr/Ad.BamR $\Delta$ Fib was digested with AvrII and the 5 kb adenofragment was isolated and introduced into the vector pBr/Ad.Bam-rITR.pac#8 described above replacing the corresponding AvrII fragment. The resulting construct was named pBr/Ad.BamR $\Delta$ Fib.pac.

[0120] Once a heterologous fiber sequence is introduced in pBr/Ad.BamR $\Delta$ Fib.pac, the fiber modified right hand adenovirus clone is introduced into a large cosmid clone as described above for pWE/Ad.AfIII-rITR. Such a large cosmid clone allows generation of adenovirus by only one homologous recombination. Ad5-based viruses with modified fibers have been made and described (nos. 98204482.8 and 99200624.7). In addition, hexon and penton sequences from serotypes from

[0121] this invention are combined with the desired fiber sequences to generate viruses which infect the target cell of choice very efficiently. For example smooth muscle cells, endothelial cells or synoviocytes all from human origin are very well infected with Ad5 based viruses with a fiber from subgroup B viruses especially adenovirus type 16.

[0122] The above described examples in which specific sequences can be deleted from the Ad5 backbone in the plasmids and replaced by corresponding sequences from other serotypes clearly show the flexibility of the system. It is evident that by the methods described above any combination of capsid gene from different serotypes can be made. Thus, chimeric recombinant Ad5-based adenoviruses are designed with desired hexon and penton sequences making the virus less sensitive for neutralisation and with desired fiber sequences allowing efficient infection in specific target tissues.

## Example 4

Construction of a plasmid-based system to generate Ad35 recombinant viruses

5 [0123] Partial restriction maps of Ad35 have been published previously (Valderrama-Leon *et al.*, 1985; Kang *et al.*, 1989; Li *et al.* 1991). An example of a functional plasmid-based system to generate recombinant adenoviruses based on Ad35 consists of the following elements:

- 10 1. An adapter plasmid comprising a left ITR and packaging sequences derived from Ad35 and at least one restriction site for insertion of an heterologous expression cassette and lacking E1 sequences. Furthermore, the adapter plasmid contains Ad35 sequences 3' from the E1B coding region including the pIX promoter and coding sequences sufficient to mediate homologous recombination of the adapter plasmid with a second nucleotide.
2. A second nucleotide comprising sequences homologous to the adapter plasmid and Ad35 sequences necessary for the replication and packaging of the recombinant virus, that is early, intermediate and late genes that are not
- 15 present in the packaging cell.
3. A packaging cell providing at least functional E1 proteins capable of complementing the E1 function of Ad35.

[0124] Ad35 DNA was isolated from a purified virus batch as follows. To 100  $\mu$ l of virus stock (Ad35:  $3.26 \times 10^{12}$  VP/ml) 10  $\mu$ l 10x DNase buffer (130 mM Tris-HCl pH7.5; 1,2 M  $\text{CaCl}_2$ ; 50mM  $\text{MgCl}_2$ ) was added. After addition of 10  $\mu$ l 10mgr/ml DNase I (Roche Diagnostics) the mixture was incubated for 1 hr. at 37°C. Following addition of 2.5  $\mu$ l 0.5M EDTA, 3.2  $\mu$ l 20% SDS and 1.5  $\mu$ l ProteinaseK (Roche Diagnostics; 20mgr/ml) samples were incubated at 50°C for 1 hr. Next, the viral DNA was isolated using the GeneClean spin kit (Bio101 Inc.) according to the manufacturers instructions. DNA was eluted from the spin column with 25  $\mu$ l sterile MilliQ water.

[0125] In the following sizes of DNA fragments and fragment numbering will be used according to Kang *et al.* (1989). Ad35 DNA was digested with EcoRI and the three fragments (approximately 22.3 (A), 7.3 (B) and 6 kb (C)) were isolated from gel using the GeneClean kit (Bio101, Inc.). pBr322 was digested with EcoRI or with EcoRI and EcoRV and digested fragments were isolated from gel and dephosphorylated with Tsap enzyme (Gibco BRL). Next, the 6 kb Ad35 C fragment was ligated to the pBr322xEcoRI fragment and the ITR-containing Ad35 fragment (EcoRI-B) was ligated to the pBr322xEcoRI/EcoRV fragment. Ligations were incubated at 16°C overnight and transformed into DH5 $\alpha$  competent bacteria (Life Techn.). Minipreps of obtained colonies were analysed for correct insertion of the Ad35 fragments by restriction analysis. Both the 6 kb and the 7.3 kb Ad35 fragment were found to be correctly inserted in pBr322. The 6kb fragment was isolated in both orientations pBr/Ad35-Eco6.0<sup>+</sup> and pBr/Ad35-Eco6.0<sup>-</sup> whereby the + stands for 5' to 3' orientation relative to pBr322. The clone with the 7.3 kb Ad35 B insert, named pBr/Ad35-Eco7.3 was partially sequenced to check correct ligation of the 3' ITR. It was found that the ITR had the sequence 5'- CATCATCAAT...-3' in the lower strand. Then pBr/Ad35-Eco7.3 was extended to the 5' end by insertion of the 6kb Ad35 fragment. Hereto, pBr/Ad35-Eco7.3 was digested with EcoRI and dephosphorylated. The fragment was isolated from gel and ligated to the 6kb Ad35 EcoRI fragment. After transformation clones were tested for correct orientation of the insert and one clone was selected, named pBr/Ad35-Eco13.3.

[0126] This clone is then extended with the ~5.4 kb Sall D fragment obtained after digestion of wt Ad35 with Sall. Hereto, the Sall site in the pBr322 backbone is removed by partial digestion of pBr/Ad35-Eco13.3 with Sall, filling in of the sticky ends by Klenow treatment and religation. One clone is selected that contains a single Sall site in the adenoviral insert. This clone, named pBr $\Delta$ sall/Ad35-Eco13.3 is then linearised with AatII which is present in the pBr322 backbone and ligated to a Sall linker with AatII complementary ends. The DNA is then digested with excess Sall and the linear fragment is isolated and ligated to the 5.4 kb Sall-D fragment from Ad35. One clone is selected that contains the Sall fragment inserted in the correct orientation in pBr/Ad35-Eco13.3. The resulting clone, pBr/Ad35.Sal2-rITR contains the 3' ~17 kb of Ad35 including the right ITR. To enable liberation of the right ITR from the vector sequences at the time of virus generation, a NotI site flanking the right ITR is introduced by PCR.

[0127] The Ad35 EcoRI-A fragment of 22.3 kb was also cloned in pBr322xEcoRI/EcoRV. One clone, named pBr/Ad35-EcoA3', was selected that apparently had a deletion of approximately 7kb of the 5' end. It did contain the Sall site at 9.4 kb in Ad35 wt DNA and approximately 1.5 kb of sequences upstream. Using this Sall site and the unique NdeI site in the pBr322 backbone this clone is extended to the 5' end by insertion of an approximately 5 kb Ad35 fragment 5' from the first Sall in Ad35 in such a way that a NotI restriction site is created at the 5' end of the Ad35 by insertion of a linker.

[0128] This clone, named pBr/Ad35.pIX-EcoA does not contain the left end sequences (ITR, packaging sequences and E1) and at the 3' end it has approximately 3.5 kb overlap with clone pBr/Ad35.Sal2-rITR.

[0129] To create an adapter plasmid, Ad35 was digested with Sall and the left end B fragment of ~9.4 kb was isolated. pBr322 was digested with EcoRV and Sall, isolated from gel and dephosphorylated with Tsap enzyme. Both fragments are ligated and clones with correct insertion and correct sequence of the left ITR are selected. To enable

liberation of the left ITR from the vector sequences at the time of virus generation, a NotI site flanking the left ITR is introduced by PCR. From this clone the E1 sequences are deleted and replaced by a polylinker sequence using PCR. The polylinker sequence is used to introduce an expression cassette for a gene of choice.

[0130] Recombinant Ad35 clones are generated by transfection of PER.C6 cells with the adapter plasmid, pBr/Ad35.pIX-EcoA and pBr/Ad35.Sal2-rITR as shown in figure 3. Homologous recombination gives rise to recombinant viruses.

#### Example 5

*The prevalence of neutralizing activity (NA) to Ad35 is low in human sera from different geographic locations*

[0131] In example 1 we have described the analysis of neutralizing activity (NA) in human sera from one location in Belgium. Strikingly, of a panel of 44 adenovirus serotypes tested, one serotype, Ad35, was not neutralized in any of the 100 sera assayed. In addition, a few serotypes, Ad26, Ad34 and Ad48 were found to be neutralized in 8%, or less, of the sera tested. This analysis was further extended to other serotypes of adenovirus not previously tested and, using a selection of serotypes from the first screen, was also extended to sera from different geographic locations.

[0132] Hereto, adenoviruses were propagated, purified and tested for neutralization in the CPE-inhibition assay as described in example 1. Using the sera from the same batch as in example 1, adenovirus serotypes 7B, 11, 14, 18 and 44/1876 were tested for neutralization. These viruses were found to be neutralized in respectively 59, 13, 30, 98 and 54 % of the sera. Thus, of this series Ad11 is neutralized with a relatively low frequency.

[0133] Since it is known that the frequency of isolation of adenovirus serotypes from human tissue as well as the prevalence of NA to adenovirus serotypes may differ on different geographic locations, we further tested a selection of the adenovirus serotypes against sera from different places. Human sera were obtained from two additional places in Europe (Bristol, UK and Leiden, The Netherlands) and from two places in the United States (Stanford, CA and Great Neck, NY). Adenoviruses that were found to be neutralized in 20% or less of the sera in the first screen, as well as Ad2, Ad5, Ad27, Ad30, Ad38, Ad43, were tested for neutralization in sera from the UK. The results of these experiments are presented in Figure 4.

[0134] Adenovirus serotypes 2 and 5 were again neutralized in a high percentage of human sera. Furthermore, some of the serotypes that were neutralized in a low percentage of sera in the first screen are neutralized in a higher percentage of sera from the UK, e.g. Ad26 (7% vs. 30%), Ad28 (13% vs. 50%), Ad34 (5% vs. 27%) and Ad48 (8% vs. 32%). Neutralizing activity against Ad11 and Ad49 that were found in a relatively low percentage of sera in the first screen, are found in an even lower percentage of sera in this second screen (13% vs. 5% and 20% vs. 11% respectively). Serotype Ad35 that was not neutralized in any of the sera in the first screen, was now found to be neutralized in a low percentage (8%) of sera from the UK. The prevalence of NA in human sera from the UK is the lowest to serotypes Ad11 and Ad35.

[0135] For further analysis, sera were obtained from two locations in the US (Stanford, CA and Great Neck, NY) and from the Netherlands (Leiden). Figure 5 presents an overview of data obtained with these sera and the previous data. Not all viruses were tested in all sera, except for Ad5, Ad11 and Ad35. The overall conclusion from this comprehensive screen of human sera is that the prevalence of neutralizing activity to Ad35 is the lowest of all serotypes throughout the western countries: on average 7% of the human sera contain neutralizing activity (5 different locations).

[0136] Another B-group adenovirus, Ad11 is also neutralized in a low percentage of human sera (average 11% in sera from 5 different locations). Adenovirus type 5 is neutralized in 56% of the human sera obtained from 5 different locations.

[0137] Although not tested in all sera, D-group serotype 49 is also neutralized with relatively low frequency in samples from Europe and from one location of the US (average 14%).

[0138] In the above described neutralization experiments a serum is judged non-neutralizing when in the well with the highest serum concentration the maximum protection of CPE is 40% compared to the controls without serum. The protection is calculated as follows:

$$\% \text{ protection} = \frac{\text{OD corresponding well} - \text{OD virus control}}{\text{OD non-infected control} - \text{OD virus control}} \times 100\%$$

[0139] As described in example 1, the serum is plated in five different dilutions ranging from 4x to 64x diluted.

[0140] Therefore, it is possible to distinguish between low titers (i.e. neutralization only in the highest serum concentrations) and high titers of NA (i.e. also neutralization in wells with the lowest serum concentration). Of the human sera used in our screen that were found to contain neutralizing activity to Ad5, 70% turned out to have high titers whereas of the sera that contained NA to Ad35, only 15% had high titers. Of the sera that were positive for NA to Ad11 only 8% had high titers.

[0141] For Ad49 this was 5%. Therefore, not only is the frequency of NA to Ad35, Ad11 and Ad49 much lower as



compared to Ad5, but of the sera that do contain NA to these viruses, the vast majority has low titers. Adenoviral vectors based on Ad11, Ad35 or Ad49 have therefore a clear advantage over Ad5 based vectors when used as gene therapy vehicles or vaccination vectors *in vivo* or in any application where infection efficiency is hampered by neutralizing activity.

5 [0142] In the following examples the construction of a vector system for the generation of safe, RCA-free Ad35-based vectors is described.

#### Example 6

##### 10 *Sequence of the human adenovirus type 35*

[0143] Ad35 viruses were propagated on PER.C6 cells and DNA was isolated as described in example 4. The total sequence was generated by Qiagen Sequence Services (Qiagen GmbH, Germany). Total viral DNA was sheared by sonification and the ends of the DNA were made blunt by T4 DNA polymerase.

15 [0144] Sheared blunt fragments were size fractionated on agarose gels and gel slices corresponding to DNA fragments of 1.8 to 2.2 kb were obtained. DNA was purified from the gel slices by the QIAquick gel extraction protocol and subcloned into a shotgun library of pUC19 plasmid cloning vectors. An array of clones in 96-wells plates covering the target DNA 8 (+/-2) times was used to generate the total sequence. Sequencing was performed on Perkin-Elmer 9700 thermo cyclers using BigDyeTerminator chemistry and AmpliTaq FS DNA polymerase followed by purification of  
20 sequencing reactions using QIAGEN DyeEx 96 technology. Sequencing reaction products were then subjected to automated separation and detection of fragments on ABI 377 XL 96 lane sequencers. Initial sequence contig sequence and gaps were filled in by primer walking reads on the target DNA or by direct sequencing of PCR products. The ends of the virus turned out to be absent in the shotgun library, most probably due to cloning difficulties resulting from the amino acids of pTP that remain bound to the ITR sequences after proteinase K digestion of the viral DNA.

25 [0145] Additional sequence runs on viral DNA solved most of the sequence in those regions, however it was difficult to obtain a clear sequence of the most terminal nucleotides. At the 5' end the sequence obtained was 5'-CCAATAATATACCT ...-3' while at the 3' end the obtained sequence was 5'-...AGGTATATTATTGATGATGGG-3'. Most human adenoviruses have a terminal sequence 5'-CATCATCAATAATATACC-3'. In addition, a clone representing the 3' end of the Ad35 DNA obtained after cloning the terminal 7 kb Ad35 EcoRI fragment into pBr322 (see example 4) also  
30 turned out to have the typical CATCATCAATAAT... sequence. Therefore, Ad35 may have the typical end sequence and the differences obtained in sequencing directly on the viral DNA are due to artefacts correlated with run-off sequence runs and the presence of residual amino acids of pTP.

[0146] The total sequence of Ad35 with corrected terminal sequences is given in Figure 6. Based sequence homology with Ad5 (genbank # M72360) and Ad7 (partial sequence Genbank # X03000) and on the location of open reading  
35 frames, the organization of the virus is identical to the general organization of most human adenoviruses, especially the subgroup B viruses. The total length of the genome is 34794 basepairs.

#### Example 7

##### 40 *Construction of a plasmid-based vector system to generate recombinant Ad35-based viruses.*

[0147] A functional plasmid-based vector system to generate recombinant adenoviral vectors comprises the following components:

- 45 1. An adapter plasmid comprising a left ITR and packaging sequences derived from Ad35 and at least one restriction site for insertion of an heterologous expression cassette and lacking E1 sequences. Furthermore, the adapter plasmid contains Ad35 sequences 3' from the E1B coding region including the pIX promoter and coding sequences enough to mediate homologous recombination of the adapter plasmid with a second nucleic acid molecule.
- 50 2. A second nucleic acid molecule, comprising sequences homologous to the adapter plasmid, and Ad35 sequences necessary for the replication and packaging of the recombinant virus, that is early, intermediate and late genes that are not present in the packaging cell.
3. A packaging cell providing at least functional E1 proteins capable of complementing the E1 function of Ad35.

[0148] Other methods for the generation of recombinant adenoviruses on complementing packaging cells are  
55 known in the art and may be applied to Ad35 viruses without departing from the invention. As an example, the construction of a plasmid based system, as outlined above, is described in detail below.

## 1) Construction of Ad35 adapter plasmids.

[0149] Hereto, the adapter plasmid pAdApt (Figure 7; described in example 2) was first modified to obtain adapter plasmids that contain extended polylinkers and that have convenient unique restriction sites flanking the left ITR and the adenovirus sequence at the 3' end to enable liberation of the adenovirus insert from plasmid vector sequences. Construction of these plasmids is described below in detail: Adapter plasmid pAdApt (Example 2) was digested with Sall and treated with Shrimp Alkaline Phosphatase to reduce religation. A linker, composed of the following two phosphorylated and annealed oligo's: ExSalPacF 5' - TCG ATG GCA AAC AGC TAT TAT GGG TAT TAT GGG TTC GAA TTA ATT AA- 3'; and ExSalPacR 5' - TCG ATT AAT TAA TTC GAA CCC ATA ATA CCC ATA ATA GCT GTT TGC CA- 3'; was directly ligated into the digested construct, thereby replacing the Sall restriction site by Pi-Pspl, Swal and PacI. This construct was named pADAPT+ExSalPac linker. Furthermore, part of the left ITR of pAdApt was amplified by PCR using the following primers: PCLIPMSF: 5'- CCC CAA TTG GTC GAC CAT CAT CAA TAA TAT ACC TTA TTT TGG -3' and pCLIPBSRG1: 5'- GCG AAA ATT GTC ACT TCC TGT G - 3'. The amplified fragment was digested with MunI and BsrGI and cloned into pAd5/Clip (see Example 2), which was partially digested with EcoRI and after purification digested with BsrGI, thereby re-inserting the left ITR and packaging signal. After restriction enzyme analysis, the construct was digested with Scal and SgrAI and an 800 bp fragment was isolated from gel and ligated into Scal/SgrAI digested pADAPT+ExSalPac linker. The resulting construct, named pIPspSalAdapt, was digested with Sall, dephosphorylated, and ligated to the phosphorylated ExSalPacF/ExSalPacR doublestranded linker mentioned above. A clone in which the PacI site was closest to the ITR was identified by restriction analysis and sequences were confirmed by sequence analysis. This novel pAdApt construct, termed pIPspAdapt (Figure 8) thus harbors two ExSalPac linkers containing recognition sequences for PacI, Pi-Pspl and BstBI, which surround the adenoviral part of the adenoviral adapter construct, and which can be used to linearize the plasmid DNA prior to cotransfection with adenoviral helper fragments.

[0150] In order to further increase transgene cloning permutations a number of polylinker variants were constructed based on pIPspAdapt. For this purpose pIPspAdapt was first digested with EcoRI and dephosphorylated. A linker composed of the following two phosphorylated and annealed oligo's:

Ecolinker+: 5' -AAT TCG GCG CGC CGT CGA CGA TAT CGA TAG CGG

CCG C -3' and Ecolinker-: 5' -AAT TGC GGC CGC TAT CGA TAT CGT CGA CGG CGC GCC G -3' was ligated into this construct, thereby creating restriction sites for Ascl, Sall, EcoRV, ClaI and NotI. Both orientations of this linker were obtained and sequences were confirmed by restriction analysis and sequence analysis. The plasmid containing the polylinker in the order 5' HindIII, KpnI, AgeI, EcoRI, Ascl, Sall, EcoRV, ClaI, NotI, NheI, HpaI, BamHI and XbaI was termed pIPspAdapt1 (Figure 9) while the plasmid containing the polylinker in the order HindIII, KpnI, AgeI, NotI, ClaI, EcoRV, Sall, Ascl, EcoRI, NheI, HpaI, BamHI and XbaI was termed pIPspAdapt2.

[0151] To facilitate the cloning of other sense or antisense constructs, a linker composed of the following two oligonucleotides was designed, to reverse the polylinker of pIPspAdapt: HindXba+ 5'-AGC TCT AGA GGA TCC GTT AAC GCT AGC GAA TTC ACC GGT ACC AAG CTT A-3'; HindXba- 5'-CTA GTA AGC TTG GTA CCG GTG AAT TCG CTA GCG TTA ACG GAT CCT CTA G-3'.

[0152] This linker was ligated into HindIII/XbaI digested pIPspAdapt and the correct construct was isolated.

[0153] Confirmation was done by restriction enzyme analysis and sequencing. This new construct, pIPspAdaptA, was digested with EcoRI and the above mentioned Ecolinker was ligated into this construct. Both orientations of this linker were obtained, resulting in pIPspAdapt3 (Figure 10), which contains the polylinker in the order XbaI, BamHI, HpaI, NheI, EcoRI, Ascl, Sall, EcoRV, ClaI, NotI, AgeI, KpnI and HindIII. All sequences were confirmed by restriction enzyme analysis and sequencing.

[0154] Adapter plasmids based on Ad35 were then constructed as follows:

[0155] The left ITR and packaging sequence corresponding to Ad35 wt sequences nucleotides 1 to 464 (Figure 6) were amplified by PCR on wtAd35 DNA using the following primers:

Primer 35F1:

5'-CGG AAT TCT TAA TTA ATC GAC ATC ATC AAT AAT ATA CCT TAT AG-3'

Primer 35R2:

5'-GGT GGT CCT AGG CTG ACA CCT ACG TAA AAA CAG-3'

[0156] Amplification introduces a PacI site at the 5' end and an AvrII site at the 3' end of the sequence.

[0157] For the amplification Platinum Pfx DNA polymerase enzyme (LTI) was used according to manufacturers instructions but with primers at 0.6  $\mu$ M and with DMSO added to a final concentration of 3%. Amplification program was as follows: 2 min. at 94°C, (30 sec. 94°C, 30 sec. at 56°C, 1 min. at 68°C) for 30 cycles, followed by 10 min. at 68°C.

[0158] The PCR product was purified using a pcr purification kit (LTI) according to the manufacturers instructions and digested with PacI and AvrII. The digested fragment was then purified from gel using the gene clean kit (Bio 101, Inc.).

5 [0159] The Ad5-based adapter plasmid pIPspAdApt-3 (Figure 10) was digested with AvrII and then partially with PacI and the 5762 bp fragment was isolated in an LMP agarose gel slice and ligated with the abovementioned PCR fragment digested with the same enzymes and transformed into electrocompetent DH10B cells (LTI). The resulting clone is named pIPspAdApt3-Ad35IITR.

[0160] In parallel, a second piece of Ad35 DNA was amplified using the following primers:

10 35F3: 5'- TGG TGG AGA TCT GGT GAG TAT TGG GAA AAC-3'  
35R4: 5'- CGG AAT TCT TAA TTA AGG GAA ATG CAA ATC TGT GAG G-3'

[0161] The sequence of this fragment corresponds to nucl. 3401 to 4669 of wtAd35 (Figure 6) and contains 1.3 kb of sequences starting directly 3' from the E1B 55k coding sequence.

15 [0162] Amplification and purification was done as described above for the fragment containing the left ITR and packaging sequence. The PCR fragment was then digested with PacI and subcloned into pNEB193 vector (New England Biolabs) digested with SmaI and PacI. The integrity of the sequence of the resulting clone was checked by sequence analysis.

20 [0163] pNEB/Ad35pF3R4 was then digested with BglII and PacI and the Ad35 insert was isolated from gel using the QIAExII kit (Qiagen). pIPspAdApt3-Ad35IITR was digested with BglII and then partially with PacI. The 3624 bp fragment (containing vector sequences, the Ad35 ITR and packaging sequences as well as the CMV promoter, multiple cloning region and polyA signal), was also isolated using the QIAExII kit (Qiagen).

25 [0164] Both fragments were ligated and transformed into competent DH10B cells (LTI). The resulting clone, pAdApt35IP3 (Figure 11), has the expression cassette from pIPspAdApt3 but contains the Ad35 left ITR and packaging sequences and a second fragment corresponding to nucl. 3401 to 4669 from Ad35. A second version of the Ad35 adapter plasmid having the multiple cloning site in the opposite orientation was made as follows:

30 pIPspAdapt1 (Figure 9) was digested with NdeI and BglII and the 0.7 kbp band containing part of the CMV promoter, the MCS and SV40 polyA was isolated and inserted in the corresponding sites of pAdApt35IP3 generating pAdApt35IP1 (Figure 12).

pAdApt35.LacZ and pAdApt35.Luc adapter plasmids were then generated by inserting the transgenes from pcDNA.LacZ (digested with KpnI and BamHI) and pAdApt.Luc (digested with HindIII and BamHI) into the corresponding sites in pAdApt35IP1. The generation of pcDNA.LacZ and pAdApt.Luc is described in WO99/55132.

## 35 2) Construction of cosmid pWE.Ad35.pXI-rITR

[0165] Figure 13 presents the various steps undertaken to construct the cosmid clone containing Ad35 sequences from bp 3401 to 34794 (end of the right ITR) that are described in detail below.

[0166] A first PCR fragment (pIX-NdeI) was generated using the following primer set:

40 35F5: 5'-CGG AAT TCG CGG CCG CGG TGA GTA TTG GGA AAA C -3'  
35R6: 5'-CGC CAG ATC GTC TAC AGA ACA G-3'

[0167] DNA polymerase Pwo (Roche) was used according to manufacturers instructions, however, with an endconcentration of 0.6  $\mu$ M of both primers and using 50 ngr wt Ad35 DNA as template.

[0168] Amplification was done as follows: 2 min. at 94 °C, 30 cycles of 30 sec. at 94 °C, 30 sec. at 65 °C and 1 min. 45 sec. at 72 °C, followed by 8 min. at 68 °C. To enable cloning in the TA cloning vector PCR2.1, a last incubation with 1 unit superTaq polymerase (HT Biotechnology LTD) for 10 min. at 72 °C was performed.

50 [0169] The 3370 bp amplified fragment contains Ad35 sequences from bp 3401 to 6772 with a NotI site added to the 5' end.

[0170] Fragments were purified using the PCR purification kit (LTI).

[0171] A second PCR fragment (NdeI-rITR) was generated using the following primers:

55 35F7: 5'-GAA TGC TGG CTT CAG TTG TAA TC -3'  
35R8: 5'- CGG AAT TCG CGG CCG CAT TTA AAT CAT CAT CAA TAA TAT ACC-3'

[0172] Amplification was done with pfx DNA polymerase (LTI) according to manufacturer's instructions but with 0.6  $\mu$ M of both primers and 3% DMSO using 10 ngr. of wtAd35 DNA as template. The program was as follows:

3 min. at 94 °C and 5 cycles of 30 sec. at 94 °C, 45 sec. at 40 °C, 2 min.45 sec. at 68 °C followed by 25 cycles of 30 sec. at 94 °C, 30 sec. at 60 °C, 2 min.45 sec. at 68 °C. To enable cloning in the TA-cloning vector PCR2.1, a last incubation with 1 unit superTag polymerase for 10 min. at 72 °C was performed. The 1.6 kb amplified fragment ranging from nucl. 33178 to the end of the right ITR of Ad35, was purified using the PCR purification kit ( LTI).

5 [0173] Both purified PCR fragments were ligated into the PCR2.1 vector of the TA-cloning kit (Invitrogen) and transformed into STBL-2 competent cells (LTI). Clones containing the expected insert were sequenced to confirm correct amplification. Next, both fragments were excised from the vector by digestion with NotI and NdeI and purified from gel using the geneClean kit (BIO 101, Inc.). Cosmid vector pWE15 (Clontech) was digested with NotI, dephosphorylated and also purified from gel. These three fragments were ligated and transformed into STBL2 competent cells (LTI). One  
10 of the correct clones that contained both PCR fragments was then digested with NdeI and the linear fragment was purified from gel using the geneClean kit. Ad35 wtDNA was digested with NdeI and the 26.6 kb fragment was purified from LMP gel using agarase enzyme (Roche) according to the manufacturers instructions. These fragments were ligated together and packaged using  $\lambda$  phage packaging extracts (Stratagene) according to the manufacturer's protocol. After infection into STBL-2 cells, colonies were grown on plates and analyzed for presence of the complete insert. One clone  
15 with the large fragment inserted in the correct orientation and having the correct restriction patterns after independent digestions with three enzymes (NcoI, PvuII and ScaI) was selected. This clone is named pWE.Ad35.pIX-rITR. It contains the Ad35 sequences from bp 3401 to the end and is flanked by NotI sites (Figure 14).

### 3) Generation of Ad35 based recombinant viruses on PER.C6.

20 [0174] Wild type Ad35 virus can be grown on PER.C6 packaging cells to very high titers. However, whether the Ad5-E1 region that is present in PER.C6 is able to complement E1-deleted Ad35 recombinant viruses is unknown. To test this, PER.C6 cells were cotransfected with the above described adapter plasmid pAdApt35.LacZ and the large backbone fragment pWE.Ad35.pIX-rITR. First, pAdApt35.LacZ was digested with PacI and pWE.Ad35.pIX-rITR was  
25 digested with NotI. Without further purification 4  $\mu$ g of each construct was mixed with DMEM (LTI) and transfected into PER.C6 cells, seeded at a density of  $5 \times 10^6$  cells in a T25 flask the day before, using Lipofectamin (LTI) according to the manufacturers instructions. As a positive control, 6  $\mu$ g of PacI digested pWE.Ad35.pIX-rITR DNA was cotransfected with a 6.7 kb NheI fragment isolated from Ad35 wt DNA containing the left end of the viral genome including the E1 region.

30 [0175] The next day medium (DMEM with 10% FBS and 10mM  $MgCl_2$ ) was refreshed and cells were further incubated. At day 2 following the transfection, cells were trypsinized and transferred to T80 flasks. The positive control flask showed CPE at five days following the transfection, showing that the pWE.Ad35.pIX-rITR construct is functional at least in the presence of Ad35-E1 proteins. The transfection with the Ad35 LacZ adapter plasmid and pWE.Ad35.pIX-rITR did not give rise to CPE. These cells were harvested in the medium at day 10 and freeze/thawed once to release virus from  
35 the cells. 4 ml of the harvested material was added to a T80 flask with PER.C6 cells (at 80% confluency) and incubated for another five days. This harvest/re-infection was repeated for two times but there was no evidence for virus associated CPE.

[0176] From this experiment it seems that the Ad5-E1 proteins are not, or not well enough, capable of complementing Ad35 recombinant viruses, however, it may be that the sequence overlap of the adapter plasmid and the  
40 pWE.Ad35.pIX-rITR backbone plasmid is not large enough to efficiently recombine and give rise to a recombinant virus genome. The positive control transfection was done with a 6.7 kb left end fragment and therefore the sequence overlap was about 3.5 kb. The adapter plasmid and the pWE.Ad35.pIX-rITR fragment have a sequence overlap of 1.3 kb. To check whether the sequence overlap of 1.3 kb is too small for efficient homologous recombination, a cotransfection was done with PacI digested pWE.Ad35.pIX-rITR and a PCR fragment of Ad35 wtDNA generated with the above mentioned  
45 35F1 and 35R4 using the same procedures as described before. The PCR fragment thus contains left end sequences up to bp 4669 and therefore has the same overlap sequences with pWE.Ad35.pIX-rITR as the adapter plasmid pAdApt35.LacZ but has Ad35 E1 sequences.

[0177] Following PCR column purification, the DNA was digested with SalI to remove possible intact template sequences. A transfection with the digested PCR product alone served as a negative control. Four days after the transfection, CPE occurred in the cells transfected with the PCR product and the Ad35 pIX-rITR fragment, and not in the negative control.  
50

[0178] This shows that 1.3 kb overlapping sequences is sufficient to generate viruses in the presence of Ad35 E1 proteins.

[0179] From these experiments we conclude that the presence of at least one of the Ad35.E1 proteins is necessary  
55 to generate recombinant Ad35 based vectors from plasmid DNA on Ad5 complementing cell lines.



## Example 8

## 1) Construction of Ad35.E1 expression plasmids

5 [0180] Since Ad5-E1 proteins in PER.C6 are not capable of complementing Ad35 recombinant viruses efficiently, Ad35 E1 proteins have to be expressed in Ad5 complementing cells (e.g. PER.C6) or a new packaging cell line expressing Ad35 E1 proteins has to be made, starting from either diploid primary human cells or established cell lines not expressing adenovirus E1 proteins. To address the first possibility, the Ad35 E1 region was cloned in expression plasmids as described below.

10 [0181] First, the Ad35 E1 region from bp 468 to bp 3400 was amplified from wtAd35 DNA using the following primer set:

35F11: 5'-GGG GTA CCG AAT TCT CGC TAG GGT ATT TAT ACC-3'

35F10: 5'-GCT CTA GAC CTG CAG GTT AGT CAG TTT CTT CTC CAC TG-3'

15

[0182] This PCR introduces a KpnI and EcoRI site at the 5'-end and a SbfI and XbaI site at the 3' end.

[0183] Amplification on 5 ngr. template DNA was done with Pwo DNA polymerase (Roche) using manufacturers instructions, however, with both primers at a final concentration of 0.6  $\mu$ M. The program was as follows: 2 min. at 94 °C, 5 cycles of 30 sec. at 94 °C, 30 sec. at 56 °C and 2 min. at 72 °C, followed by 25 cycles of 30 sec. at 94 °C, 30 sec. at 20 60 °C and 2 min. at 72 °C, followed by 10 min. at 72 °C. PCR product was purified by a PCR purification kit (LTI) and digested with KpnI and XbaI. The digested PCR fragment was then ligated to the expression vector pRSVhbvNeo (see below), also digested with KpnI and XbaI. Ligations were transformed into competent STBL-2 cells (LTI) according to manufacturers instructions and colonies were analysed for the correct insertion of Ad35E1 sequences into the polylinker in between the RSV promoter and HBV polyA.

25 [0184] The resulting clone was named pRSV.Ad35-E1 (Figure 15). The Ad35 sequences in pRSV.Ad35-E1 were checked by sequence analysis.

[0185] pRSVhbvNeo was generated as follows: pRc-RSV (Invitrogen) was digested with PvuII, dephosphorylated with TSAP enzyme (LTI) and the 3 kb vector fragment was isolated in low melting point agarose (LMP). Plasma pPG-KneopA (Figure 16; described in WO96/35798, was digested with SspI completely to linearise the plasmid and facilitate 30 partial digestion with PvuII. Following the partial digestion with PvuII, the resulting fragments were separated on a LMP agarose gel and the 2245 bp PvuII fragment, containing the PGK promoter, neomycine resistance gene and HBVpolyA, was isolated. Both isolated fragments were ligated to give the expression vector pRSV-pNeo that now has the original SV40prom-neo-SV40polyA expression cassette replaced by a PGKprom-neo-HBVpolyA cassette (Figure 17). This plasmid was further modified to replace the BGHpA with the HBVpA as follows: pRSVpNeo was linearised with Scal and further digested with XbaI. The 1145 bp fragment, containing part of the Amp gene and the RSV promoter 35 sequences and polylinker sequence, was isolated from gel using the GeneClean kit (Bio Inc. 101).

[0186] Next pRSVpNeo was linearised with Scal and further digested with EcoRI partially and the 3704 bp fragment containing the PGKneo cassette and the vector sequences were isolated from gel as above. A third fragment, containing the HBV polyA sequence flanked by XbaI and EcoRI at the 5' and 3' end respectively, was then generated 40 by PCR amplification on pRSVpNeo using the following primer set:

HBV-F: 5'- GGC TCT AGA GAT CCT TCG CGG GAC GTC -3' and

HBV-R: 5'- GGC GAA TTC ACT GCC TTC CAC CAA GC -3'.

45 [0187] Amplification was done with Elongase enzyme (LTI) according to the manufacturers instructions with the following conditions: 30 seconds at 94 °C, then 5 cycles of 45 seconds at 94 °C, 1 minute at 42 °C and 1 minute 68 °C, followed by 30 cycles of 45 seconds at 94 °C, 1 minute at 65 °C and 1 minute at 68 °C, followed by 10 minutes at 68 °C. The 625 bp PCR fragment was then purified using the Qiaquick PCR purification kit, digested with EcoRI and XbaI and purified from gel using the GeneClean kit. The three isolated fragments were ligated and transformed into DH5 $\alpha$  com- 50 petent cells (LTI) to give the construct pRSVhbvNeo (Figure 18). In this construct the transcription regulatory regions of the RSV expression cassette and the neomycine selection marker are modified to reduce overlap with adenoviral vectors that often contain CMV and SV40 transcription regulatory sequences.

2) Generation of Ad35 recombinant viruses on PER.C6 cells cotransfected with an Ad35-E1 expression construct.

55

[0188] PER.C6 cells were seeded at a density of  $5 \times 10^6$  cells in a T25 flask and the next day transfected with a DNA mixture containing:

- 1 µg pAdApt35.LacZ digested with PacI
- 5 µg pRSV.Ad35E1 undigested
- 2 µg pWE.Ad35.pIX-rITR digested with NotI

5 [0189] Transfection was done using Lipofectamine according to the manufacturers instructions. Five hours after addition of the transfection mixture to the cells, medium was removed and replaced by fresh medium. After two days cells were transferred to T80 flasks and further cultured. One week post-transfection 1 ml of the medium was added to A549 cells and the following day cells were stained for LacZ expression. Blue cells were clearly visible after two hours of staining indicating that recombinant LacZ expressing viruses were produced. The cells were further cultured but no  
10 clear appearance of CPE was noted. However, after 12 days clumps of cells appeared in the monolayer and 18 days following transfection cells were detached. Cells and medium were then harvested, freeze-thawed once and 1 ml of the crude lysate was used to infect PER.C6 cells in a 6-well plate. Two days after infection cells were stained for LacZ activity. After two hours 15% of the cells were stained blue. To test for the presence of wt and / or replicating competent viruses, A549 cells were infected with these viruses and further cultured. No signs of CPE were found indicating the  
15 absence of replication competent viruses.

[0190] These experiments show that recombinant AdApt35.LacZ viruses were made on PER.C6 cells cotransfected with an Ad35-E1 expression construct.

### 3) Ad35 recombinant viruses escape neutralization in human serum containing neutralizing activity to Ad5 viruses.

20 [0191] The AdApt35.LacZ viruses were then used to investigate infection in the presence of serum that contains neutralizing activity to Ad5 viruses. Purified Ad5-based LacZ virus served as a positive control for NA. Hereto, PER.C6 cells were seeded in a 24-wells plate at a density of  $2 \times 10^5$  cells/well. The next day a human serum sample with high neutralizing activity to Ad5 was diluted in culture medium in five steps of five times dilutions. 0.5 ml of diluted serum was  
25 then mixed with  $4 \times 10^6$  virus particles AdApt5.LacZ virus in 0.5 ml medium and after 30 minutes of incubation at 37 °C, 0.5 ml of the mixture was added to PER.C6 cells in duplicate. For the AdApt35.LacZ viruses, 0.5 ml of the diluted serum samples were mixed with 0.5 ml crude lysate containing AdApt35.LacZ virus and after incubation 0.5 ml of this mixture was added to PER.C6 cells in duplo.

[0192] Virus samples incubated in medium without serum was used as a positive control for infection. After two  
30 hours of infection at 37 °C, medium was added to reach a final volume of 1 ml and cells were further incubated. Two days after infection cells were stained for LacZ activity. The results are shown in Table II. From these results it is clear that whereas AdApt5.LacZ viruses are efficiently neutralized, AdApt35.LacZ viruses remain infectious irrespective of the presence of human serum. This proves that recombinant Ad35-based viruses escape neutralization in human sera that contain NA to Ad5-based viruses.

### Example 9:

#### *An Ad5/fiber35 chimeric vector with cell type specificity for hemopoietic CD34 Lin stem cells*

40 [0193] In example 3 we have described the generation of a library of Ad5 based adenoviruses harboring fiber proteins of other serotypes. As a non-limiting example for the use of this library we here describe the identification of fiber-modified adenoviruses that show improved infection of hemopoietic stem cells.

[0194] Cells isolated from human bone marrow, umbilical cord blood, or mobilized peripheral blood carrying the flow cytometric phenotype of being positive for the CD34 antigen and negative for the early differentiation markers CD33, CD38, and CD71 (lin) are commonly referred to as hemopoietic stem cells (HSC). Genetic modification of these  
45 cells is of major interest since all hemopoietic lineages are derived from these cells and therefore the HSC is a target cell for the treatment of many acquired or congenital human hemopoietic disorders. Examples of diseases that are amendable for genetic modification of HSC, but not limited to, include Hurlers disease, Hunters disease, Sanfilippus disease, Morquios disease, Gaucher disease, Farbers disease, Niemann-Pick disease, Krabbe disease, Metachromatic Leucodystrophy, I-cell disease, severe immunodeficiency syndrome, Jak-3 deficiency, Fucosidose deficiency, thalassemia, and erythropoietic porphyria. Besides these hemopoietic disorders also strategies to prevent or treat acquired immunodeficiency syndrome (AIDS) and hemopoietic cancers are based on the genetic modification of HSCs or cells derived from the HSCs such as CD4 positive T lymphocytes in case of AIDS. The examples listed above thus aim at  
50 introducing DNA into the HSC in order to complement on a genetic level for a gene and protein deficiency. In case of strategies for AIDS or cancer, the DNA to be introduced into the HSC can be anti-viral genes or suicide genes.

[0195] Besides the examples listed above, there are several other areas in which efficient transduction of HSCs using adenoviral vectors plays an important role. For instance in the field of tissue engineering. In this area it is important to drive differentiation of HSCs to specific lineages. Some, non-limiting, examples are *ex vivo* bone formation, car-

tilage formation, skin formation, as well as the generation of T-cell precursors or endothelial cell precursors. The generation of bone, cartilage or skin in bioreactors can be used for transplantation after bone fractures or spinal cord lesions or severe burn injuries.

[0196] Naturally, transduced cells can also directly be re-infused into a patient. The formation of large numbers of endothelial cell precursor from HSCs is of interest since these endothelial precursors cells can home, after re-infusion, to sites of cardiovascular injury such as ischemia. Likewise, the formation of large numbers of T-cells from HSCs is of interest since these T-cell precursors can be primed, *ex vivo*, to eradicate certain targets in the human body after reinfusion of the primed T-cells. Preferred targets in the human body can be tumours or virus infected cells.

[0197] From the examples described above, it can be concluded that efficient gene delivery to HSCs is a major interest for the field of gene therapy. Therefore, alteration of the adenovirus serotype 5 host cell range to be able to target HSCs *in vitro* as well as *in vivo* is a major interest of the invention. To identify a chimeric adenovirus with preferred infection characteristics for human HSCs, we generated a library of Ad5 based viruses carrying the fiber molecule from alternative serotypes (serotypes 8, 9, 13, 16, 17, 32, 35, 45, 40-L, 51). The generation of this fiber modified library is described in example 3. Ad5 was taken along as a reference. A small panel of this library was tested on human TF-1 (erythroidleukemia, ATCC CRL-2003) whereas all chimaeric viruses generated were tested on human primary stroma cells and human HSCs. Human TF-1 cell were routinely maintained in DMEM supplemented with 10% FCS and 50 ng/ml IL-3 (Sandoz, Basel, Switzerland). Human primary fibroblast-like stroma, isolated from a bone marrow aspirate, is routinely maintained in DMEM/ 10% FCS. Stroma was seeded at a concentration of  $1 \times 10^5$  cells per well of 24-well plates. 24 hours after seeding cells were exposed for 2 hours to 1000 virus particles per cell of Ad5, Ad5.Fib16, Ad5.Fib17, Ad5.Fib35, Ad5.Fib40-L, or Ad5.Fib51 all carrying the green fluorescent protein (GFP) as a marker. After 2 hours cells were washed with PBS and reseeded in medium without addition of virus. TF-1 cells were seeded at a concentration of  $2 \times 10^5$  cells per well of 24-well plates and were also exposed for 2 hours to 1000 virus particles of the different chimeric adenoviruses. Virus was removed by washing the cells after the 2 hours exposure. Both cell types were harvested 48 hours after virus exposure and analysed for GFP expression using a flow cytometer. The results on TF-1 cells, shown in figure 19, demonstrates that chimeric adenoviruses carrying a fiber from serotypes 16, 35, or 51 (all derived from adenovirus subgroup B) have preferred infection characteristics as compared to Ad5 (subgroup C), Ad5.Fib17 (subgroup D), or Ad5.Fib40-L (subgroup F). Primary human stroma was tested since these cells are commonly used as a "feeder" cell to allow proliferation and maintenance of HSCs under *ex vivo* culture conditions. In contrast to the transduction of TF-1 cells, none of the fiber chimeric adenoviruses were able to efficiently transduce human primary stroma (Figure 20). Reasonable infection of human fibroblast-like primary stroma was observed only with Ad5 despite the observation that none of the known receptor molecules are expressed on these cells (see table III). The absence of infection of human stroma using the chimeric viruses is advantageous since in a co-culture setting, the chimeric adenovirus will not be absorbed primarily by the stroma "feeder" cells.

[0198] To test the transduction capacity of the fiber chimaeric viruses, a pool of umbilical cord blood (3 individuals) was used for the isolation of stem cells. CD34<sup>+</sup> cells were isolated from mononuclear cell preparation using a MACS laboratory separation system (Miltenyi Biotec) using the protocol supplied by the manufacturer. Of the CD34<sup>+</sup> cells,  $2 \times 10^5$  were seeded in a volume of 150  $\mu$ l DMEM (no serum; Gibco, Gaithersburg, MD) and 10  $\mu$ l of chimeric adenovirus (to give a final virus particles/cell ratio of 1000) was added. The chimeric adenoviruses tested were Ad5, Ad5.Fib16, Ad5.Fib35, Ad5.Fib17, Ad5.Fib51 all containing Green fluorescent protein (GFP) as a marker. Cells were incubated for 2 hours in a humidified atmosphere of 10% CO<sub>2</sub> at 37°C.

[0199] Thereafter, cells were washed once with 500  $\mu$ l DMEM and resuspended in 500  $\mu$ l of StemPro-34 SF medium (Life Technologies, Grand Island, NY).

[0200] Cells were then cultured for 5 days in 24-well plates (Greiner, Frickenhausen, Germany) on irradiated (20 Gy) pre-established human bone marrow stroma (ref 1), in a humidified atmosphere of 10% CO<sub>2</sub> at 37°C. After 5 days, the entire cell population was collected by trypsinization with 100  $\mu$ l 0.25% Trypsin-EDTA (Gibco). The number of cells before and after 5 days of culture was determined using a hemacytometer. The number of CD34<sup>+</sup> and CD34<sup>+</sup>CD33,38,71 cells in each sample was calculated from the total number of cells recovered and the frequency of the CD34<sup>+</sup>CD33,38,71<sup>+</sup> cells in the whole population as determined by FACS analysis. The transduction efficiency was determined by FACS analysis while monitoring in distinct sub populations the frequency of GFP expressing cells as well as the intensity of GFP per individual cell. The results of this experiment, shown in figure 21, demonstrates that adenovirus serotype 5 or the chimeric adenovirus Ad5.Fib17 does not infect CD34<sup>+</sup>Lin<sup>-</sup> cells as witnessed by the absence of GFP expression. In contrast, with the chimeric viruses carrying the fiber molecule of serotypes 16, 51, or 35 high percentages of GFP positive cells are scored in this cell population. Specificity for CD34<sup>+</sup>Lin<sup>-</sup> is demonstrated since little GFP expression is observed in CD34<sup>+</sup> cells that are also expressing CD33, CD38, and CD71. Subfractioning of the CD34<sup>+</sup>Lin<sup>-</sup> cells (Figure 22) showed that the percentage of cells positive for GFP declines using Ad5.Fib16, Ad5.Fib35, or Ad5.Fib51 when the cells become more and more positive for the early differentiation markers CD33 (myeloid), CD71 (erythroid), and CD38 (common early differentiation marker).

[0201] These results thus demonstrate the specificity of the chimeric adenoviruses Ad5.Fib16, Ad5.Fib35, and



Ad5.Fib51 for HSCs. Figure 23 shows an alignment of the Ad5 fiber with the chimeric B-group fiber proteins derived from Ad16, 35 and 51. By determining the number of cells recovered after the transduction procedure the toxicity of adenovirus can be determined. The recovery of the amount of CD34<sup>+</sup> cells as well as the amount of CD34<sup>+</sup>Lin<sup>-</sup> (Figure 24) demonstrates that a 2 hour exposure to 1000 adenovirus particles did not have an effect on the number of cells recovered.

#### Example 10

##### *An Ad5/fiber35 chimeric vector with cell type specificity for Dendritic cells*

[0202] Dendritic cells are antigen presenting cells (APC), specialized to initiate a primary immune response and able to boost a memory type of immune response. Dependent on their stage of development, DC display different functions: immature DC are very efficient in the uptake and processing of antigens for presentation by Major Histocompatibility Complex (MHC) class I and class II molecules, whereas mature DC, being less effective in antigen capture and processing, perform much better at stimulating naive and memory CD4<sup>+</sup> and CD8<sup>+</sup> T cells, due to the high expression of MHC molecules and co-stimulatory molecules at their cell surface. The immature DCs mature *in vivo* after uptake of antigen, travel to the T-cell areas in the lymphoid organs, and prime T-cell activation.

[0203] Since DCs are the cells responsible for triggering an immune response there has been a long standing interest in loading DCs with immunostimulatory proteins, peptides or the genes encoding these proteins to trigger the immune system. The applications for this strategy are in the field of cancer treatment as well as in the field of vaccination. So far, anti-cancer strategies have focussed primarily on *ex vivo* loading of DCs with antigen (protein or peptide). These studies have revealed that this procedure resulted in induction of cytotoxic T cell activity. The antigens used to load the cells are generally identified as being tumor specific. Some, non-limiting, examples of such antigens are GP100, mage, or Mart-1 for melanoma.

[0204] Besides treatment of cancer many other potential human diseases are currently being prevented through vaccination. In the vaccination strategy, a "crippled" pathogen is presented to the immune system via the action of the antigen presenting cells, i.e. the immature DCs. Well-known examples of disease prevention via vaccination strategies include Hepatitis A, B, and C, influenza, rabies, yellow fever, measles. Besides these well-known vaccination programs, research programs for treatment of malaria, ebola, river blindness, HIV and many other diseases are being developed.

[0205] Many of the above mentioned pathogens are considered to dangerous for the generation of a "crippled" pathogen vaccine. This latter thus calls for the isolation and characterization of proteins of each pathogen which is able to mount a "full blown" immune response thus resulting in complete protection upon challenge with wild type pathogen.

[0206] For this strategy of loading DCs with immunostimulatory proteins or peptides to become therapeutically feasible

[0207] At least two distinct criteria have to be met 1) the isolation of large numbers of DCs which can be isolated, manipulated, and reinfused into a patient, making the procedure autologous. To date, it is possible to obtain such large quantities of immature DCs from cultured peripheral blood monocytes from any given donor. 2) a vector which can transduce DCs efficiently such that the DNA encoding for an immunostimulatory protein can be delivered. The latter is extremely important since it has become clear that the time required for DCs to travel to the lymphoid organs is such that most proteins or peptides are already released from the DCs resulting in incomplete immune priming. Because DCs are terminally differentiated and thus non-dividing cells, recombinant adenoviral vectors are being considered for delivering the DNA encoding for antigens to DCs. Ideally this adenovirus should have a high affinity for dendritic cells but also should not be recognized by neutralizing antibodies of the host such that *in vivo* transduction of DCs can be accomplished. This latter would omit the need for *ex vivo* manipulations of DCs but would result in a medical procedure identical to the vaccination programs which are currently in place, i.e. intramuscular or subcutaneous injection predominantly. Thus, DC transduced by adenoviral vectors encoding an immunogenic protein may be ideally suited to serve as natural adjuvants for immunotherapy and vaccination

[0208] From the above described examples, it can be concluded that efficient gene delivery to DCs is a major interest for the field of gene therapy. Therefore, alteration of the adenovirus serotype 5 host cell range to be able to target DCs *in vitro* as well as *in vivo* is a major interest of the invention. To identify a chimeric adenovirus with preferred infection characteristics for human DCs, we generated a library of Ad5 based viruses carrying the fiber molecule from alternative serotypes (serotypes 8, 9, 13, 16, 17, 32, 35, 45, 40-L, 51). Ad5 was taken along as a reference.

[0209] We evaluated the susceptibility of human monocyte derived immature and mature DC to recombinant chimeric adenoviruses expressing different fibers.

[0210] Human PBMC from healthy donors were isolated through Ficoll-Hypaque density centrifugation. Monocytes were isolated from PBMC by enrichment for CD14<sup>+</sup> cells using staining with FITC labeled anti-human CD 14 monoclonal antibody (Becton Dickinson), anti FITC microbeads and MACS separation columns (Miltenyi Biotec).

[0211] This procedure usually results in a population of cells that are < 90 % CD14<sup>+</sup> as analysed by FACS. Cells



were placed in culture using RPMI-1640 medium (Gibco) containing 10% Foetal Bovine Serum (Gibco), 200 ng/ml rhu GM-CSF (R&D/ITK diagnostics, 100 ng/ml rhu IL-4 (R&D/ITK diagnostics) and cultured for 7 days with feeding of the cultures with fresh medium containing cytokines on alternate days. The immature DC resulting from this procedure after 7 days express a phenotype CD83<sup>-</sup>, CD14<sup>low</sup> or CD14<sup>-</sup>, HLA-DR<sup>+</sup>, as was demonstrated by FACS analysis. Immature DC are matured by culturing the cells in medium containing 100 ng/ml TNF- $\alpha$  for 3 days, where after they expressed CD83 on their cell surface.

[0212] In a pilot experiment  $5 \cdot 10^5$  immature DCs were seeded in wells of 24-well plates and exposed for 24 hours to 100 and 1000 virus particles per cell of each fiber recombinant virus. Virus tested was adenovirus serotype 5 (Ad5), and the fiber chimeric viruses based on Ad5: Ad5.Fib12, Ad5.Fib16, Ad5.Fib28, Ad5.Fib32, Ad5.Fib40-L (long fiber of serotype 40), Ad5.Fib49, and Ad5.Fib51 (where Fibxx stands for the serotype of which the fiber molecule is derived). these viruses are derived from subgroup C, A, B, D, D, F, D, and B respectively. After 24-hours cells were lysed (1% Triton X-100/ PBS) and luciferase activity was determined using a protocol supplied by the manufacturer (Promega, Madison, WI, USA). The results of this experiment, shown in figure 25, demonstrates that Ad5 poorly infects immature DCs as witnessed by one low level of transgene expression. In contrast, Ad5.Fib16 and Ad5.Fib51 (both a B-group fiber chimeric virus) and also Ad5.Fib40-L (Subgroup F) show efficient infection of immature DCs based on luciferase transgene expression.

[0213] In a second experiment,  $5 \cdot 10^5$  immature and mature DC were infected with 10000 virus particles per cell of Ad5, Ad5.Fib16, Ad5.Fib40-L, and Ad5.Fib51 all carrying the LacZ gene as a marker. LacZ expression was monitored by flow cytometric analysis using a CM-FDG kit system and the instructions supplied by the manufacturer (Molecular probes, Leiden, The Netherlands). The results of this experiment, shown in figure 26, correlates with the previous experiment in that Ad5.Fib16 and Ad5.Fib51 are superior to Ad5 in transducing mature and immature human DCs. Also, this experiment shows that Ad5.Fib40-L is not as good as Ad5.Fib16 and Ad5.Fib51 but better than Ad5.

[0214] Based on these results we tested other chimeric adenoviruses containing fibers of B group viruses e.g. Ad5.Fib11 and Ad5.Fib35 for there capacity to infect DCs. We focussed on immature DCs since these are the cells that process an expressed transgene product into MHC class I and II presentable peptides. Immature DC's were seeded at a cell density of  $5 \cdot 10^5$  cells/well in 24 well plates (Costar) and infected with 1000 and 5000 virus particles per cell after which the cells were cultured for 48 hours under conditions for immature DCs prior to cell lysis and Luciferase activity measurements. The result of this experiment, shown in figure 27, demonstrate that Ad5 based chimeric adenoviruses containing fibers of group-B viruses efficiently infect immature DCs. In a fourth experiment we again infected immature DCs identically as described in the former experiments but this time Ad5, Ad5.Fib16, and Ad5.Fib35 were used carrying green fluorescent protein (GFP) as a markergene. The results on GFP expression measured with a flow cytometer 48 hours after virus exposure is shown in figure 28 and correlates with the data obtained so far.

[0215] Thus, the results so far are consistent in that Ad5 based vectors carrying a fiber from a alternative adenovirus derived from subgroup B predominantly fiber of 35, 51, 16, and 11 are superior to Ad5 for transducing human DCs.

[0216] The adenoviruses disclosed herein are also very suitable in vaccination of animals. To illustrate this, we tested DCs derived from mouse and chimpanzee to identify whether these viruses can be used in these animal models. This latter in particular since the receptor for human adenovirus derived from subgroup B is unknown to date and therefore it is unknown whether this protein is conserved among species. For both species immature DCs were seeded at a density of  $10^5$  cells per well of 24-well plates. Cells were subsequently exposed for 48 hours to 1000 virus particles per cell of Ad5, Ad5.Fib16, and Ad5.Fib51 in case of mouse DC and Ad5, and Ad5.Fib35 in case of chimpanzee DCs (see figure 29). The mouse experiment was performed with viruses carrying luciferase as a marker and demonstrated approximately 10-50 fold increased luciferase activity as compared to Ad5. The chimpanzee DCs were infected with the GFP viruses and were analysed using a flow cytometer. These results, also shown in figure 29, demonstrate that Ad5 (3%) transduces chimpanzee DCs very poorly as compared to Ad5.Fib35 (66.5%).

## Example 11

### *Construction of a plasmid-based vector system to generate Ad11-based recombinant viruses*

[0217] The results of the neutralization experiments described in Example 5 show that Ad11, like Ad35, was also not neutralized in the vast majority of human serum samples.

[0218] Therefore, recombinant adenoviruses based on Ad11 are preferred above the commonly used Ad2 and Ad5-based vectors as vectors for gene therapy treatment and vaccination. Both Ad35 and Ad11 are B-group viruses and are classified as viruses belonging to DNA homology cluster 2 (Wadell, 1984).

[0219] Therefore, the genomes of Ad35 and Ad11 are very similar. To generate a plasmid based system for the production of Ad11-based recombinant viruses the adapter plasmid pAdApt35IP1 generated in Example 7 is modified as follows. Construct pAdApt35IP1 is digested with AvrII and then partially with PacI. The digestion mixture is separated on gel and the 4.4 kb fragment containing the expression cassette and the vector backbone is isolated using the gene-

clean kit (BIO 101, Inc.). Then a PCR amplification is performed on wtAd11 DNA using the primers 35F1 and 35R2 (see Example 7) using Pwo DNA polymerase according to the manufacturers instructions. The obtained PCR fragment of 0.5 kb is purified using the PCR purification kit (LTI) and ligated to the above prepared fragment of pAdApt35IP1. This gives construct pAdApt11-35IP1 in which the 5' adenovirus fragment is exchanged for the corresponding sequence of Ad11. Next, pAdApt11-35IP1 is digested with BglII and partially with PacI. The obtained fragments are separated on gel and the 3.6 kb fragment containing the vector sequences, the 5' adenovirus fragment and the expression cassette is purified from gel as above. Next, a PCR fragment is generated using primers 35F3 and 35R4 (see Example 7) on wtAd11 DNA.

[0220] Amplification is done as above and the obtained 1.3 kb fragment is purified and digested with BglII and PacI. The isolated fragments are then ligated to give construct pAdApt11IP1. This adapter plasmid now contains Ad11 sequences in stead of Ad35 sequences. Correct amplification of PCR amplified Ad11 sequences, is verified by comparison of the sequence in this clone with the corresponding sequence of Ad11 DNA. The latter is obtained by direct sequencing on Ad11 DNA using the indicated PCR primers. The large cosmid clone containing the Ad11 backbone is generated as follows. First, a PCR fragment is amplified on Ad11 DNA using the primers 35F5 and 35R6 with Pwo DNA polymerase as described in Example 7 for Ad35 DNA. The PCR fragment is then purified using the PCR purification kit (LTI) and digested with NotI and NdeI. The resulting 3.1 kb fragment is isolated from gel using the geneclean kit (Bio 101, Inc.). A second PCR fragment is then generated on Ad11 DNA using the primers 35F7 and 35R8 (see Example 7) with Pwo DNA polymerase according to the manufacturers instructions and purified using the PCR purification kit (LTI). This amplified fragment is also digested with NdeI and NotI and the resulting 1.6 kb fragment is purified from gel as above.

[0221] The two digested PCR fragments are then ligated together with cosmid vector pWE15, previously digested with NotI and dephosphorylated using Tsap enzyme (LTI) according to manufacturers instructions. One clone is selected that has one copy of both fragments inserted. Correct clones are selected by analytical NotI digestion that gives a fragment of 4.7 kb. Confirmation is obtained by a PCR reaction using primers 35F5 and 35R8 that gives a fragment of the same size. The correct clone is then linearized with NdeI and isolated from gel. Next, wtAd11 DNA is digested with NdeI and the large 27 kb fragment is isolated from Low melting point agarose gel using agarase enzyme (Roche) according to the manufacturers instructions. Both fragments are then ligated and packaged using  $\lambda$  phage packaging extracts (Stratagene) according to the manufacturers protocol. After infection into STBL-2 cells (LTI) colonies are grown on plates and analysed for the presence of the complete insert.

[0222] The functionality of selected clones is then tested by cotransfection on PER.C6. Hereto, the DNA is digested with NotI and 6  $\mu$ g is cotransfected with 2  $\mu$ g of a PCR fragment generated on Ad11 DNA with primers 35F1 and 35R4 (see example 7). Correct clones give CPE within one week following transfection. The correct clone is named pWE.Ad11.pIX-rlTR.

[0223] Using the above described procedure, a plasmid-based system consisting of an adapter plasmid suitable for insertion of foreign genes and a large helper fragment containing the viral backbone is generated. Recombinant Ad11-based viruses are made using the methods described inhere for Ad35-based recombinant viruses.

## Example 12

### *Neutralization of adenoviruses in samples derived from patients*

[0224] In the neutralization experiments described in Examples 1 and 5, all samples were derived from healthy volunteers.

[0225] Since one of the applications of non-neutralized vectors is in the field of gene therapy, it is interesting to investigate whether Ad35 is also neutralized with a low frequency and with low titers in groups of patients that are candidates for treatment with gene therapy.

#### *- Cardio-vascular disease patients*

[0226] 26 paired serum and pericardial fluid (PF) samples were obtained from patients with heart failure. These were tested against Ad5 and Ad35 using the neutralization assay described in Example 1. The results confirmed the previous data with samples from healthy volunteers. 70% of the serum samples contained NA to Ad5 and 4% to Ad35. In the pericardial fluid samples the titers were lower resulting in a total of 40% with NA to Ad5 and none to Ad35. There was a good correlation between NA in PF and serum i.e. there were no positive PF samples without NA in the paired serum sample. These results show that non-neutralized vectors based on Ad35 are preferred over Ad5 vectors for treatment of cardio-vascular diseases. As is true for all forms of non-neutralized vectors in this application, the vector may be based on the genome of the non-neutralized serotype or may be based on Ad5 (or another serotype) though displaying at least the major capsid proteins (hexon, penton and optionally fiber) of the non-neutralized serotype.

- Rheumatoid Arthritis patients

[0227] The molecular determinant underlying arthritis is not yet known but both T-cell dysfunction and imbalanced growth factor production in joints is known to cause inflammation and hyperplasia of synovial tissue. The synoviocytes start to proliferate and invade the cartilage and bone which leads to destruction of these tissues. Current treatment starts (when in an early stage) with administration of anti-inflammatory drugs (anti-TNF, IL1-RA, IL-10) and/or conventional drugs (e.g. MTX, sulfasalazine). In late stage RA synovectomy is performed which is based on surgery, radiation, or chemical intervention. An alternative or additional option is treatment via gene therapy where an adenoviral vector is delivered directly into the joints of patients and expresses an anti-inflammatory drug or a suicide gene. Previous studies performed in rhesus monkeys suffering from collagen-induced arthritis have shown that Ad5-based vectors carrying a marker gene can transduce synoviocytes. Whether in the human situation adenoviral delivery is hampered by the presence of NA is not known. To investigate the presence of NA in synovial fluid (SF) of RA patients, SF samples were obtained from a panel of 53 random selected patients suffering from rheumatoid arthritis (RA).

[0228] These were tested against several wt adenoviruses using the neutralization assay as described in Example 1. Results of this screen are presented in Table III. Adenovirus type 5 was found to be neutralized in 72% of the SF samples. Most of these samples contain high titers of NA as also the highest dilution of the SF sample that was tested (64x) neutralized Ad5 viruses. This means that adenoviral vector delivery to the synoviocytes in the joints of RA patients will be very inefficient. Moreover, since the titers in the SF are so high it is doubtful whether lavage of the joints prior to vector injection will remove enough of the NA. Of the other serotypes that were tested Ad35 was shown to be neutralized in only 4% of the samples. Therefore, these data confirm the results obtained in serum samples from healthy patients and show that for treatment of rheumatoid arthritis Ad35-based vectors or chimeric vectors displaying at least some of the capsid proteins from Ad35 are preferred vectors.

### Example 13

#### *Modifications in the backbone of Ad35-based viruses*

##### 1) Generation of pBr/Ad35.Pac-rITR and pBr/Ad35.PRn

[0229] Example 4 describes the generation of the Ad35 subclone pBr/Ad35.Eco13.3. This clone contains Ad35 sequences from bp 21943 to the end of the right ITR cloned into the EcoRI and EcoRV sites of pBr322. To extend these sequences to the PacI site located at bp 18137 in Ad35, pBr/Ad35.Eco13.3 (see Example 4) was digested with AatII and SnaBI and the large vector-containing fragment was isolated from gel using the QIAEX II gel extraction kit (Qia-gen). Ad35 wt DNA was digested with PacI and SnaBI and the 4.6 kb fragment was isolated as above. This fragment was then ligated to a double-stranded (ds) linker containing a PacI and an AatII overhang. This linker was obtained after annealing the following oligonucleotides:

A-P1: 5'-CTG GTG GTT AAT-3'

A-P2: 5'-TAA CCA CCA GAC GT-3'

[0230] The ligation mix containing the ds linker and the PacI-SnaBI Ad35 fragment was separated from unligated linker on a LMP gel. The 4.6 kb band was cut out the gel, molten at 65 °C, and then ligated to the purified pBr/Ad35.Eco13.3 vector fragment digested with AatII and SnaBI. Ligations were transformed into electrocompetent DH10B cells (Life Technologies Inc.). The resulting clone, pBr/Ad35.Pac-rITR, contained Ad35 sequences from the PacI site at bp 18137 upto the right ITR.

[0231] Next, a unique restriction site was introduced at the 3' end of the right ITR to be able to free the ITR from vector sequences. Hereto, a PCR fragment was used that covers Ad35 sequences from the NdeI site at bp 33165 to the right ITR having the restriction sites SwaI, NotI and EcoRI attached to the rITR. The PCR fragment was generated using primers 35F7 and 35R8 (described in example 7). After purification, the PCR fragment was cloned into the AT cloning vector (Invitrogen) and sequenced to verify correct amplification.

[0232] The correct amplified clone was then digested with EcoRI, blunted with Klenow enzym and subsequently digested with NdeI and the PCR fragment was isolated. In parallel, the NdeI in the pBr vector in pBr/Ad35.Pac-rITR was removed as follows: A pBr322 vector from which the NdeI site was removed by digestion with NdeI, Klenow treatment and religation, was digested with AatII and NheI. The vector fragment was isolated in LMP gel and ligated to the 16.7 kb Ad35 AatII-NheI fragment from pBr/Ad35.Pac-rITR that was also isolated in an LMP gel. This generated pBr/Ad35.Pac-rITR.ΔNdeI. Next pBr/Ad35.Pac-rITR.ΔNdeI was digested with NheI, the ends were filled in using Klenow enzym and the DNA was then digested with NdeI. The large fragment containing the vector and Ad35 sequences was isolated. Ligation of this vector fragment and the PCR fragment resulted in pBr/Ad35.PRn. In this clone specific



sequences coding for fiber, E2A, E3, E4 or hexon can be manipulated. In addition, promoter sequences that drive for instance the E4 proteins or the E2 can be mutated or deleted and exchanged for heterologous promoters.

## 2) Generation of Ad35-based viruses with fiber proteins from different serotypes.

[0233] Adenoviruses infect human cells with different efficiencies. Infection is accomplished by a two step process involving: 1. the fiber proteins that mediate binding of the virus to specific receptors on the cells, and 2. the penton proteins that mediate internalization by interaction of for example the RGD sequence to integrins present on the cell surface. For subgroup B viruses of which Ad35 is a member, the cellular receptor for the fiber protein is not known. There are striking differences in infection efficiency of human cells of subgroup B viruses compared to subgroup C viruses like Ad5 (see WO 00/03029 and EP 99200624.7). Even within one subgroup infection efficiencies of certain human cells may differ between various serotypes. For example, the fiber of Ad16, when present on an Ad5-based recombinant virus infects primary endothelial cells, smooth muscle cells and synoviocytes of human and rhesus monkey origin better than Ad5 chimeric viruses carrying the fiber of Ad35 or Ad51. Thus, to obtain high infection efficiencies of Ad35-based viruses, it may be necessary to change the fiber protein for a fiber protein of a different serotype. The technology for such fiber chimeras is described for Ad5-based viruses in Example 3, and is below exemplified for Ad35 viruses.

[0234] First, most fiber sequences are deleted from the Ad35 backbone in construct pBr/Ad35.PRN as follows: The left flanking sequences and part of the fiber protein in Ad35 ranging from bp 30225 upstream of a unique MluI site up to bp 30872 (numbers according to wt Ad35 sequence as disclosed in Figure 6) in the tail of fiber are amplified using primers

DF35-1 : 5'-CAC TCA CCA CCT CCA ATT CC-3'

and

DF35-2: 5'-CGG GAT CCC GTA CGG GTA GAC AGG GTT GAA GG-3'

[0235] This PCR amplification introduces an unique BsiWI site in the tail of the fiber gene.

[0236] The right flanking sequences ranging from the end of the fiber protein at bp 31798 to bp 33199 (numbering according to wtAd35 sequence, Figure 6) , 3' from the unique NdeI site is amplified using primers

DF35-3: 5'-CGG GAT CCG CTA GCT GAA ATA AAG TTT AAG TGT TTT TAT TTA AAA TCA C-3'

and

DF35-4: 5'-CCA GTT GCA TTG CTT GGT TGG-3'.

[0237] This PCR introduces a unique NheI site in the place of the fiber sequences. PCR amplification is done with Pwo DNA polymerase (Roche) according to the manufacturers instructions. After amplification the PCR products are purified using a PCR purification kit and the fragments are digested with BamHI and ligated together. The 2 kb ligated fragments are purified from gel and cloned in the PCR Script Amp vector (Stratagene). Correct amplification is checked by sequencing. The PCR fragment is then excised as a MluI/NdeI fragment and cloned in pBr/Ad35.PRN digested with the same enzymes. This generates pBr/Ad35.PRΔfib, a shuttle vector suitable to introduce fiber sequences of alternative serotypes. This strategy is analogous to the fiber modification strategy for Ad5-based viruses as disclosed in WO00/03029. Primers that are listed in Table I of that application were used to amplify fiber sequences of various subgroups of adenovirus. For amplification of fibers that are cloned in the pBr/Ad35.PRΔfib the same (degenerate) primer sequences can be used, however, the NdeI site in the forward primers (tail oligonucleotides A to E) should be changed to a BsiWI site and the NsiI site in the reverse oligo (knob oligonucleotide 1 to 8) should be changed in a NheI site. Thus fiber 16 sequences are amplified using the following degenerate primers:

5'- CCK GTS TAC CCG TAC GAA GAT GAA AGC-3' and 5'-CCG GCT AGC TCA GTC ATC TTC TCT GAT ATA-3'. Amplified sequences are then digested with BsiWI and NheI and cloned into pBr/Ad35.PRΔfib digested with the same enzymes to generate pBr/Ad35.PRfib16. The latter construct is then digested with PacI and SmaI and the insert is isolated from gel. The PacI/SmaI Ad35 fragment with modified fiber is then cloned into the corresponding sites of pWE/Ad35.pIX-rITR to give pWE/Ad35.pIX-rITR:fib16. This cosmid backbone can then be used with an Ad35-based adapter plasmid to generate Ad35 recombinant viruses that display the fiber of Ad16. Other fiber sequences can be amplified with (degenerate) primers as mentioned above. If one of the fibers sequences turns out to have an internal BsiWI or NheI site, the PCR fragment has to be digested partially with that enzyme.

## 3) Generation of Ad35-based viruses with inducible, E1 independent, E4 expression.

[0238] The adenovirus E4 promoter is activated by expression of E1 proteins. It is not known whether the Ad5 E1 proteins are capable of mediating activation of the Ad35 E4 promoter.

[0239] Therefore, to enable production of Ad35 recombinant viruses on PER.C6 cells, it may be advantageous to make E4 expression independent of E1. This can be achieved by replacing the Ad35-E4 promoter by heterologous promoter sequences like, but not limited to, the 7xTetO promoter.

5 [0240] Recombinant E1-deleted Ad5-based vectors are shown to have residual expression of viral genes from the vector backbone in target cells, despite the absence of E1 expression. Viral gene expression increases the toxicity and may trigger a host immune response to the infected cell. For most applications of adenoviral vectors in the field of gene therapy and vaccination it is desired to reduce or diminish the expression of viral genes from the backbone. One way to achieve this is to delete all, or as much as possible, sequences from the viral backbone. By deleting E2A, E2B or E4 genes and/or the late gene functions, one has to complement for these functions during production. This complemen-  
10 tation can either be by means of a helper virus or through stable addition of these functions, with or without inducible transcription regulation, to the producer cell.

[0241] Methods to achieve this have been described for Ad5 and are known in the art. One specific method is replacement of the E4 promoter by promoter sequences that are not active in the target cells. E4 proteins play a role in for example replication of adenoviruses through activation of the E2 promoter and in late gene expression through reg-  
15 ulation of splicing and nuclear export of late gene transcripts. In addition, at least some of the E4 proteins are toxic to cells. Therefore, reduction or elimination of E4 expression in target cells will further improve Ad35-based vectors. One way to achieve this is to replace the E4 promoter by an heterologous promoter that is inactive in the target cells.

[0242] An example of a heterologous promoter/activator system that is inactive in target cells is the tetracyclin inducible TetO system (Gossen and Bujard, 1992). Other prokaryotic or synthetic promoter/activator systems may be  
20 used. In this example, the E4 promoter in the backbone of the viral vector is replaced by a DNA fragment containing 7 repeats of the tetracyclin responsive element from the tet operon (7xTetO).

[0243] A strong transactivator for this promoter is a fusion protein containing the DNA binding domain of the tet repressor and the activation domain of VP16 (Tet transactivator protein, Tta). Strong E4 expression, independent of E1 expression, can be accomplished in PER.C6 cells expressing Tta. Tta expressing PER.C6 cells have been generated  
25 and described (see Example 15). Ad5 derived E1-deleted viruses with E4 under control of 7xTetO can be generated and propagated on these cells. Following infection in cells of human or animal origin (that do not express the Tta trans-activator), E4 expression was found to be greatly diminished compared to E1 deleted viruses with the normal E4 pro-moter.

[0244] Below the construction of pWE/Ad35.pIX-rITR.TetO-E4, a cosmid helper vector to produce viruses with the  
30 E4 promoter replacement, is described.

[0245] First, a fragment was generated by PCR amplification on pBr/Ad35.PRn DNA using the following primers:

35 355ITR: 5'- GAT CCC GAG CTC ACA ACG TCA TTT TCC CAC G-3'  
and  
353ITR: 5'-CGG AAT TCG CGG CCG CAT TTA AAT C-3'

[0246] This fragment contains sequences between bp 34656 (numbering according to wtAd35) and the NotI site 3' or the right ITR in pBr/Ad35.PRn and introduces an SstI site 5' of the right ITR sequence.

[0247] A second PCR fragment was generated on pBr/Ad35.PRn DNA using primers:

40 35DE4: 5'-CCC AAG CTT GCT TGT GTA TAT ATA TTG TGG-3' and  
35F7: See example 7.

[0248] This PCR amplifies Ad35 sequences between bp 33098 and 34500 (numbering according to wtAd35) and  
45 introduces a HindIII site upstream of the E4 Tata-box. With these two PCR reactions the right- and left -flanking sequences of the E4 promoter are amplified. For amplification, Pwo DNA polymerase was used according to manufac-turers instructions. A third fragment containing the 7xTetO promoter was isolated from construct pAAO-E-TATA-7xTetO by digestion with SstI and HindIII. The generation of pAAO-E-TATA-7xTetO is described below. The first PCR fragment (355/353) was then digested with SstI and NotI and ligated to the 7xTetO fragment. The ligation mixture was then  
50 digested with HindIII and NotI and the 0.5 kb fragment is isolated from gel. The second PCR fragment (35DE4/35F7) was digested with NdeI and HindIII and gel purified. These two fragments are then ligated into pBr/Ad35.PRn digested with NdeI and NotI to give pBr/Ad35.PR.TetOE4. The modification of the E4 promoter is then transferred to the Ad35 helper cosmid clone by exchanging the PacI/SwaI fragment of the latter with the one from pBr/Ad35.PR.TetOE4 to give pWE/Ad35.pIX-rITR.TetOE4. pAAO-E-TATA.7xTetO was generated as follows. Two oligonucleotides were synthesized:

55 TATApus: 5'-AGC TTT CTT ATA AAT TTT CAG TGT TAG ACT AGT AAA TTG CTT AAG-3' and  
TATAmin: 5'-AGC TCT TAA GCA ATT TAC TAG TCT AAC ACT GAA AAT TTA TAA GAA-3'  
(The underlined sequences form a modified TATA box).

[0249] The oligonucleotides were annealed to yield a double stranded DNA fragment with 5' overhangs that are compatible with HindIII digested DNA. The product of the annealing reaction was ligated into HindIII digested pGL3-Enhancer Vector (Promega) to yield pAAO-E-TATA. The clone that had the HindIII site at the 5' end of the insert restored was selected for further cloning.

5 [0250] Next, the heptamerized tet-operator sequence was amplified from the plasmid pUHC-13-3 (Gossen and Bujard, 1992) in a PCR reaction using the Expand PCR system (Roche) according to the manufacturers protocol. The following primers were used:

tet3: 5'- CCG GAG CTC CAT GGC CTA ACT CGA GTT TAC CAC TCC C-3'

10 tet5: 5'-CCC AAG CTT AGC TCG ACT TTC ACT TTT CTC-3'

[0251] The amplified fragment was digested with SstI and HindIII (these sites are present in tet3 and tet5 respectively) and cloned into SstI/HindIII digested pAAO-E-TATA giving rise to pAAO-E-TATA-7xtetO

[0252] To test the functionality of the generated pWE/Ad35.pIX-rITR.TetOE4 cosmid clone, the DNA was digested with NotI. The left end of wtAd35 DNA was then amplified using primers 35F1 and 35R4 (see example 7). Following amplification, the PCR mixture was purified and digested with SalI to remove intact viral DNA. Then 4gr of both the digested pWE/Ad35.pIX-rITR.TetOE4 and the PCR fragment was cotransfected into PER.C6-tTA cells that were seeded in T25 flasks the day before. Transfected cells were transferred to T80 flasks after two days and another two days later CPE was obtained, showing that the cosmid backbone is functional.

20

#### Example 14

Generation of cell lines capable of complementing E1-deleted Ad35 viruses

25 *Generation of pIG135 and pIG270*

[0253] Construct pIG.E1A.E1B contains E1 region sequences of Ad5 corresponding to nucleotides 459 to 3510 of the wt Ad5 sequence (Genbank accession number M72360) operatively linked to the human phosphoglycerate kinase promoter (PGK) and the Hepatitis B Virus polyA sequences. The generation of this construct is described in WO97/00326. The E1 sequences of Ad5 were replaced by corresponding sequences of Ad35 as follows. pRSV.Ad35-E1 (described in example 8) was digested with EcoRI and Sse8387I and the 3 kb fragment corresponding to the Ad35 E1 sequences was isolated from gel. Construct pIG.E1A.E1B was digested with Sse8387I completely and partially with EcoRI. The 4.2 kb fragment corresponding to vector sequences without the Ad5 E1 region but retaining the PGK promoter were separated from other fragments on LMP agarose gel and the correct band was excised from gel. Both obtained fragments were ligated resulting in pIG.Ad35-E1.

35 [0254] This vector was further modified to remove the LacZ sequences present in the pUC119 vector backbone. Hereto, the vector was digested with BsaAI and BstXI and the large fragment was isolated from gel. A double stranded oligo was prepared by annealing the following two oligos:

40 BB1: 5'-GTG CCT AGG CCA CGG GG-3' and

BB2: 5'-GTG GCC TAG GCA C-3'

[0255] Ligation of the oligo and the vector fragment resulted in construct pIG135. Correct insertion of the oligo restores the BsaAI and BstXI sites and introduces a unique AvrII site. Next, we introduced a unique site at the 3' end of the Ad35-E1 expression cassette in pIG135. Hereto, the construct was digested with SapI and the 3' protruding ends were made blunt by treatment with T4 DNA polymerase. The thus treated linear plasmid was further digested with BsrGI and the large vector containing fragment was isolated from gel. To restore the 3' end of the HBVpolyA sequence and to introduce a unique site, a PCR fragment was generated using the following primers:

50 270F: 5'- CAC CTC TGC CTA ATC ATC TC -3' and

270R: 5'- GCT CTA GAA ATT CCA CTG CCT TCC ACC -3'

[0256] The PCR was performed on pIG.Ad35.E1 DNA using Pwo polymerase (Roche) according to the manufacturers instructions. The obtained PCR product was digested with BsrGI and dephosphorylated using Tsap enzym (LTI), the latter to prevent insert dimerization on the BsrGI site. The PCR fragment and the vector fragment were ligated to yield construct pIG270.

55

*Ad35 E1 sequences are capable of transforming rat primary cells*

[0257] New born WAG/RIJ rats were sacrificed at 1 week of gestation and kidneys were isolated. After careful removal of the capsule, kidneys were disintegrated into a single cell suspension by multiple rounds of incubation in trypsin/EDTA (LTI) at 37 °C and collection of floating cells in cold PBS containing 1% FBS. When most of the kidney was trypsinized all cells were resuspended in DMEM supplemented with 10% FBS and filtered through a sterile cheese cloth. Baby Rat Kidney (BRK) cells obtained from one kidney were plated in 5 dishes (Greiner, 6 cm). When a confluency of 70-80% was reached, the cells were transfected with 1 or 5 µgr DNA/dish using the CaPO<sub>4</sub> precipitation kit (LTI) according to the manufacturers instructions. The following constructs were used in separate transfections: pIG.E1A.E1B (expressing the Ad5-E1 region), pRSV.Ad35-E1, pIG.Ad35-E1 and pIG270 (the latter expressing the Ad35-E1). Cells were incubated at 37 °C, 5% CO<sub>2</sub> until foci of transformed cells appeared. Table IV shows the number of foci that resulted from several transfection experiments using circular or linear DNA. As expected, the Ad5-E1 region efficiently transformed BRK cells. Foci also appeared in the Ad35-E1 transfected cell layer although with lower efficiency. The Ad35 transformed foci appeared at a later time point: ~2 weeks post transfection compared with 7-10 days for Ad5-E1. These experiments clearly show that the E1 genes of the B group virus Ad35 are capable of transforming primary rodent cells.

[0258] This proves the functionality of the Ad35-E1 expression constructs and confirms earlier findings of the transforming capacity of the B-group viruses Ad3 and Ad7 (Dijkema, 1979).

[0259] To test whether the cells in the foci were really transformed a few foci were picked and expanded. From the 7 picked foci at least 5 turned out to grow as established cell lines.

*Generation of new packaging cells derived from primary human amniocytes*

[0260] Amniotic fluid obtained after amniocentesis was centrifugated and cells were resuspended in AmnioMax medium (LTI) and cultured in tissue culture flasks at 37 °C and 10 % CO<sub>2</sub>. When cells were growing nicely (approximately one cell division/24 hrs.), the medium was replaced with a 1:1 mixture of AmnioMax complete medium and DMEM low glucose medium (LTI) supplemented with Glutamax I (end concentration 4mM, LTI) and glucose (end concentration 4.5 gr/L, LTI) and 10% FBS (LTI). For transfection ~ 5x10<sup>5</sup> cells were plated in 10 cm tissue culture dishes. The day after, cells were transfected with 20 µgr of circular pIG270/dish using the CaPO<sub>4</sub> transfection kit (LTI) according to manufacturers instructions and cells were incubated overnight with the DNA precipitate. The following day, cells were washed 4 times with PBS to remove the precipitate and further incubated for over three weeks until foci of transformed cells appeared.

[0261] Once a week the medium was replaced by fresh medium. Other transfection agents like, but not limited to, LipofectAmine (LTI) or PEI (Polyethylenimine, high molecular weight, water-free, Aldrich) were used. Of these three agents PEI reached the best transfection efficiency on primary human amniocytes: ~1% blue cells 48 hrs. following transfection of pAdApt35.LacZ.

[0262] Foci are isolated as follows. The medium is removed and replaced by PBS after which foci are isolated by gently scraping the cells using a 50-200 µl Gilson pipette with a disposable filter tip. Cells contained in 10 µl PBS were brought in a 96 well plate containing 15 µl trypsin/EDTA (LTI) and a single cell suspension was obtained by pipetting up and down and a short incubation at room temperature.

[0263] After addition of 200 µl of the above described 1:1 mixture of AmnioMax complete medium and DMEM with supplements and 10% FBS, cells were further incubated. Clones that continued to grow were expanded and analysed their ability to complement growth of E1-deleted adenoviral vectors of different sub-groups, specifically ones derived from B-group viruses specifically from Ad35 or Ad11.

*Generation of new packaging cell lines from human embryonic retinoblasts*

[0264] Human retina cells are isolated from the eyes of aborted fetuses and cultured in DMEM medium (LTI) supplemented with 10% FES (LTI). The day before transfection, ~5x10<sup>5</sup> cells are plated in 6 cm dishes and cultured overnight at 37 °C and 10% CO<sub>2</sub>. Transfection is done using the CaPO<sub>4</sub> precipitation kit (LTI) according to the manufacturers instructions. Each dish is transfected with 8-10 µgr pIG270 DNA, either as a circular plasmid or as a purified fragment. To obtain the purified fragment, pIG270 was digested with AvrII and XbaI and the 4 kb fragment corresponding to the Ad35 E1 expression cassette was isolated from gel by agarase treatment (Roche). The following day, the precipitate is washed away carefully by four washes with sterile PBS. Then fresh medium is added and transfected cells are further cultured until foci of transformed cells appear. When large enough (>100 cells) foci are picked and brought into 96-wells as described above. Clones of transformed human embryonic retinoblasts that continue to grow, are expanded and tested for their ability to complement growth of E1-deleted adenoviral vectors of different sub-groups specifically ones derived from B-group viruses specifically from Ad35 or Ad11.



*New packaging cell lines derived from PER.C6*

[0265] As described in example 8, it is possible to generate and grow Ad35 E1-deleted viruses on PER.C6 cells with cotransfection of an Ad35-E1 expression construct, e.g. pRSV.Ad35.E1. However, large scale production of recombinant adenoviruses using this method is cumbersome because for each amplification step a transfection of the Ad35-E1 construct is needed. In addition, this method increases the risk of non-homologous recombination between the plasmid and the virus genome with high chances of generation of recombinant viruses that incorporate E1 sequences resulting in replication competent viruses. To avoid this, the expression of Ad35-E1 proteins in PER.C6 has to be mediated by integrated copies of the expression plasmid in the genome. Since PER.C6 cells are already transformed and express Ad5-E1 proteins, addition of extra Ad35-E1 expression may be toxic for the cells, however, it is not impossible to stably transfect transformed cells with E1 proteins since Ad5-E1 expressing A549 cells have been generated.

[0266] In an attempt to generate recombinant adenoviruses derived from subgroup B virus Ad7, Abrahamsen *et al.* (1997) were not able to generate E1-deleted viruses on 293 cells without contamination of wt Ad7. Viruses that were picked after plaque purification on 293-ORF6 cells (Brough *et al.*, 1996) were shown to have incorporated Ad7 E1B sequences by non-homologous recombination. Thus, efficient propagation of Ad7 recombinant viruses proved possible only in the presence of Ad7-E1B expression and Ad5-E4-ORF6 expression. The E1B proteins are known to interact with cellular as well as viral proteins (Bridge *et al.*, 1993; White, 1995). Possibly, the complex formed between the E1B 55K protein and E4-ORF6 which is necessary to increase mRNA export of viral proteins and to inhibit export of most cellular mRNAs, is critical and in some way serotype specific. The above experiments suggest that the E1A proteins of Ad5 are capable of complementing an Ad7-E1A deletion and that Ad7-E1B expression in adenovirus packaging cells on itself is not enough to generate a stable complementing cell line. To test whether one or both of the Ad35-E1B proteins is/are the limiting factor in efficient Ad35 vector propagation on PER.C6 cells, we have generated an Ad35 adapter plasmid that does contain the E1B promoter and E1B sequences but lacks the promoter and the coding region for E1A. Hereto, the left end of wtAd35 DNA was amplified using the primers 35F1 and 35R4 (both described in Example 7) with Pwo DNA polymerase (Roche) according to the manufacturers instructions. The 4.6 kb PCR product was purified using the PCR purification kit (LTI) and digested with SnaBI and ApaI enzymes. The resulting 4.2 kb fragment was then purified from gel using the QIAExII kit (Qiagen).

[0267] Next, pAdApt35IP1 (Example 7) was digested with SnaBI and ApaI and the 2.6 kb vector containing fragment was isolated from gel using the GeneClean kit (BIO 101, Inc). Both isolated fragments were ligated to give pBr/Ad35.leftITR-pIX. Correct amplification during PCR was verified by a functionality test as follows: The DNA was digested with BstBI to liberate the Ad35 insert from vector sequences and 4 µgr of this DNA was cotransfected with 4 µgr of NotI digested pWE/Ad35.pIX-rITR (Example 7) into PER.C6 cells.

[0268] The transfected cells were passaged to T80 flasks at day 2 and again two days later CPE had formed showing that the new pBr/Ad35.leftITR-pIX construct contains functional E1 sequences. The pBr/Ad35.leftITR-pIX construct was then further modified as follows. The DNA was digested with SnaBI and HindIII and the 5' HindIII overhang was filled in using Klenow enzyme. Religation of the digested DNA and transformation into competent cells (LTI) gave construct pBr/Ad35.leftITR-pIXΔE1A. This latter construct contains the left end 4.6 kb of Ad35 except for E1A sequences between bp 450 and 1341 (numbering according to wtAd35, Figure 6) and thus lacks the E1A promoter and most of the E1A coding sequences. pBr/Ad35.leftITR-pIXΔE1A was then digested with BstBI and 2 µgr of this construct was cotransfected with 6 µgr of NotI digested pWE/Ad35.pIX-rITR (Example 7) into PER.C6 cells. One week following transfection full CPE had formed in the transfected flasks.

[0269] This experiment shows that the Ad35-E1A proteins are functionally complemented by Ad5-e1A expression in PER.C6 cells and that at least one of the Ad35-E1B proteins cannot be complemented by Ad5-E1 expression in PER.C6. It further shows that it is possible to make a complementing cell line for Ad35 E1-deleted viruses by expressing Ad35-E1B proteins in PER.C6. Stable expression of Ad35-E1B sequences from integrated copies in the genome of PER.C6 cells may be driven by the E1B promoter and terminated by a heterologous poly-adenylation signal like, but not limited to, the HBVpA.

[0270] The heterologous pA signal is necessary to avoid overlap between the E1B insert and the recombinant vector, since the natural E1B termination is located in the pIX transcription unit which has to be present on the adenoviral vector.

[0271] Alternatively, the E1B sequences may be driven by a heterologous promoter like, but not limited to the human PGK promoter or by an inducible promoter like, but not limited to the 7xtetO promoter (Gossen and Bujard, 1992). Also in these cases the transcription termination is mediated by a heterologous pA sequence, e.g. the HBV pA. The Ad35-E1B sequences at least comprise one of the coding regions of the E1B 21K and the E1B 55K proteins located between nucleotides 1611 and 3400 of the wt Ad35 sequence. The insert may also include (part of the) Ad35-E1B sequences between nucleotides 1550 and 1611 of the wt Ad35 sequence.



## Example 15

Generation of producer cell lines for the production of recombinant adenoviral vectors deleted in early region 1 and early region 2A

Generation of PER.C6-tTA cells

[0272] Here is described the generation of cell lines for the production of recombinant adenoviral vectors that are deleted in early region 1 (E1) and early region 2A (E2A).

[0273] The producer cell lines complement for the E1 and E2A deletion from recombinant adenoviral vectors *in trans* by constitutive expression of both E1 and E2A genes. The pre-established Ad5-E1 transformed human embryo retinoblast cell line PER.C6 (WO 97/00326) was further equipped with E2A expression cassettes.

[0274] The adenoviral E2A gene encodes a 72 kDa DNA Binding Protein with has a high affinity for single stranded DNA.

[0275] Because of its function, constitutive expression of DBP is toxic for cells. The ts125E2A mutant encodes a DBP which has a Pro→Ser substitution of amino acid 413. Due to this mutation, the ts125E2A encoded DBP is fully active at the permissive temperature of 32°C, but does not bind to ssDNA at the non-permissive temperature of 39°C. This allows the generation of cell lines that constitutively express E2A, which is not functional and is not toxic at the non-permissive temperature of 39°C. Temperature sensitive E2A gradually becomes functional upon temperature decrease and becomes fully functional at a temperature of 32°C, the permissive temperature.

A. Generation of plasmids expressing the wild type E2A- or temperature sensitive ts125E2A gene.

[0276]

pcDNA3wtE2A: The complete wild-type early region 2A (E2A) coding region was amplified from the plasmid pBR/Ad.Bam-rITR (ECACC deposit P97082122) with the primers DBPpcr1 and DBPpcr2 using the Expand™ Long Template PCR system according to the standard protocol of the supplier (Boehringer Mannheim). The PCR was performed on a Biometra Trio Thermoblock, using the following amplification program: 94°C for 2 minutes, 1 cycle; 94°C for 10 seconds + 51°C for 30 seconds + 68°C for 2 minutes, 1 cycle; 94°C for 10 seconds + 58°C for 30 seconds + 68°C for 2 minutes, 10 cycles; 94°C for 10 seconds + 58°C for 30 seconds + 68°C for 2 minutes with 10 seconds extension per cycle, 20 cycles; 68°C for 5 minutes, 1 cycle. The primer DBPpcr1: CGG GAT CCG CCA CCA TGG CCA GTC GGG AAG AGG AG (5' to 3') contains a unique *Bam*HI restriction site (underlined) 5' of the Kozak sequence (*italic*) and start codon of the E2A coding sequence. The primer DBPpcr2: CGG AAT TCT TAA AAA TCA AAG GGG TTC TGC CGC (5' to 3') contains a unique *Eco*RI restriction site (underlined) 3' of the stop codon of the E2A coding sequence. The bold characters refer to sequences derived from the E2A coding region. The PCR fragment was digested with *Bam*HI/*Eco*RI and cloned into *Bam*HI/*Eco*RI digested pcDNA3 (Invitrogen), giving rise to pcDNA3wtE2A.

pcDNA3tsE2A: The complete ts125E2A-coding region was amplified from DNA isolated from the temperature sensitive adenovirus mutant H5ts125. The PCR amplification procedure was identical to that for the amplification of wtE2A. The PCR fragment was digested with *Bam*HI/*Eco*RI and cloned into *Bam*HI/*Eco*RI digested pcDNA3 (Invitrogen), giving rise to pcDNA3tsE2A. The integrity of the coding sequence of wtE2A and tsE2A was confirmed by sequencing.

B. Growth characteristics of producer cells for the production of recombinant adenoviral vectors cultured at 32-, 37- and 39°C.

[0277] PER.C6 cells were cultured in Dulbecco's Modified Eagle Medium (DMEM, Gibco BRL) supplemented with 10% Fetal Bovine Serum (FBS, Gibco BRL) and 10mM MgCl<sub>2</sub> in a 10% CO<sub>2</sub> atmosphere at either 32°C, 37°C or 39°C. At day 0, a total of 1 x 10<sup>6</sup> PER.C6 cells were seeded per 25cm<sup>2</sup> tissue culture flask (Nunc) and the cells were cultured at either 32°C, 37°C or 39°C. At day 1-8, cells were counted. Figure 30 shows that the growth rate and the final cell density of the PER.C6 culture at 39°C are comparable to that at 37°C. The growth rate and final density of the PER.C6 culture at 32°C were slightly reduced as compared to that at 37°C or 39°C. No significant cell death was observed at any of the incubation temperatures. Thus PER.C6 performs very well both at 32°C and 39°C, the permissive and non-permissive temperature for ts125E2A, respectively.

## C. Transfection of PER.C6 with E2A expression vectors; colony formation and generation of cell lines

[0278] One day prior to transfection,  $2 \times 10^6$  PER.C6 cells were seeded per 6 cm tissue culture dish (Greiner) in DMEM, supplemented with 10% FBS and 10mM  $MgCl_2$  and incubated at 37°C in a 10%  $CO_2$  atmosphere. The next day, the cells were transfected with 3, 5 or 8µg of either pcDNA3, pcDNA3wtE2A or pcDNA3tsE2A plasmid DNA per dish, using the LipofectAMINE PLUS™ Reagent Kit according to the standard protocol of the supplier (Gibco BRL), except that the cells were transfected at 39°C in a 10%  $CO_2$  atmosphere. After the transfection, the cells were constantly kept at 39°C, the non-permissive temperature for ts125E2A. Three days later, the cells were put in DMEM supplemented with 10% FBS, 10mM  $MgCl_2$  and 0.25mg/ml G418 (Gibco BRL), and the first G418 resistant colonies appeared at 10 days post transfection. As shown in table 1, there was a dramatic difference between the total number of colonies obtained after transfection of pcDNA3 (~200 colonies) or pcDNA3tsE2A (~100 colonies) and pcDNA3wtE2A (only 4 colonies). These results indicate that the toxicity of constitutively expressed E2A can be overcome by using a temperature sensitive mutant of E2A (ts125E2A) and culturing of the cells at the non-permissive temperature of 39°C.

[0279] From each transfection, a number of colonies was picked by scraping the cells from the dish with a pipette. The detached cells were subsequently put into 24 wells tissue culture dishes (Greiner) and cultured further at 39°C in a 10%  $CO_2$  atmosphere in DMEM, supplemented with 10% FBS, 10mM  $MgCl_2$  and 0.25mg/ml G418. As shown in table 1, 100% of the pcDNA3 transfected colonies (4/4) and 82% of the pcDNA3tsE2A transfected colonies (37/45) were established to stable cell lines (the remaining 8 pcDNA3tsE2A transfected colonies grew slowly and were discarded). In contrast, only 1 pcDNA3wtE2A-transfected colony could be established. The other 3 died directly after picking.

[0280] Next, the E2A expression levels in the different cell lines were determined by Western blotting. The cell lines were seeded on 6 well tissue culture dishes and sub-confluent cultures were washed twice with PBS (NPBI) and lysed and scraped in RIPA (1% NP-40, 0.5% sodium deoxycholate and 0.1% SDS in PBS, supplemented with 1mM phenylmethylsulfonylfluoride and 0.1 mg/ml trypsin inhibitor). After 15 minutes incubation on ice, the lysates were cleared by centrifugation. Protein concentrations were determined by the Bio-Rad protein assay, according to standard procedures of the supplier (BioRad). Equal amounts of whole-cell extract were fractionated by SDS-PAGE on 10% gels. Proteins were transferred onto Immobilon-P membranes (Millipore) and incubated with the αDBP monoclonal antibody B6. The secondary antibody was a horseradish-peroxidase-conjugated goat anti mouse antibody (BioRad). The Western blotting procedure and incubations were performed according to the protocol provided by Millipore. The complexes were visualized with the ECL detection system according to the manufacturer's protocol (Amersham). Figure 31 shows that all of the cell lines derived from the pcDNA3tsE2A transfection expressed the 72-kDa E2A protein (left panel, lanes 4-14; middle panel, lanes 1-13; right panel, lanes 1-12). In contrast, the only cell line derived from the pcDNAwtE2A transfection did not express the E2A protein (left panel, lane 2). No E2A protein was detected in extract from a cell line derived from the pcDNA3 transfection (left panel, lane 1), which served as a negative control. Extract from PER.C6 cells transiently transfected with pcDNA3ts125 (left panel, lane 3) served as a positive control for the Western blot procedure. These data confirmed that constitutive expression of wtE2A is toxic for cells and that using the ts125 mutant of E2A could circumvent this toxicity.

## D. Complementation of E2A deletion in adenoviral vectors on PER.C6 cells constitutively expressing full-length ts125E2A.

[0281] The adenovirus Ad5.dl802 is an Ad 5 derived vector deleted for the major part of the E2A coding region and does not produce functional DBP. Ad5.dl802 was used to test the E2A trans-complementing activity of PER.C6 cells constitutively expressing ts125E2A. Parental PER.C6 cells or PER.C6tsE2A clone 3-9 were cultured in DMEM, supplemented with 10% FBS and 10mM  $MgCl_2$  at 39°C and 10%  $CO_2$  in 25 cm<sup>2</sup> flasks and either mock infected or infected with Ad5.dl802 at an m.o.i. of 5. Subsequently the infected cells were cultured at 32°C and cells were screened for the appearance of a cytopathic effect (CPE) as determined by changes in cell morphology and detachment of the cells from the flask. Full CPE appeared in the Ad5.dl802 infected PER.C6tsE2A clone 3-9 within 2 days.

[0282] No CPE appeared in the Ad5.dl802 infected PER.C6 cells or the mock infected cells. These data showed that PER.C6 cells constitutively expressing ts125E2A complemented in trans for the E2A deletion in the Ad5.dl802 vector at the permissive temperature of 32°C.

## E. Serum-free suspension culture of PER.C6tsE2A cell lines.

[0283] Large-scale production of recombinant adenoviral vectors for human gene therapy requires an easy and scaleable culturing method for the producer cell line, preferably a suspension culture in medium devoid of any human or animal constituents. To that end, the cell line PER.C6tsE2A c5-9 (designated c5-9) was cultured at 39°C and 10%  $CO_2$  in a 175 cm<sup>2</sup> tissue culture flask (Nunc) in DMEM, supplemented with 10% FBS and 10mM  $MgCl_2$ . At sub-confluency (70-80% confluent), the cells were washed with PBS (NPBI) and the medium was replaced by 25 ml serum free

suspension medium Ex-cell™ 525 (JRH) supplemented with 1 x L-Glutamine (Gibco BRL), hereafter designated SFM. Two days later, cells were detached from the flask by flicking and the cells were centrifuged at 1,000 rpm for 5 minutes. The cell pellet was resuspended in 5 ml SFM and 0.5 ml cell suspension was transferred to a 80 cm<sup>2</sup> tissue culture flask (Nunc), together with 12 ml fresh SFM. After 2 days, cells were harvested (all cells are in suspension) and counted in a Burkert cell counter. Next, cells were seeded in a 125 ml tissue culture erlenmeyer (Corning) at a seeding density of 3 x 10<sup>5</sup> cells per ml in a total volume of 20 ml SFM. Cells were further cultured at 125 RPM on an orbital shaker (GFL) at 39°C in a 10% CO<sub>2</sub> atmosphere. Cells were counted at day 1-6 in a Burkert cell counter. In Figure 4, the mean growth curve from 8 cultures is shown. PER.C6tsE2A c5-9 performed well in serum free suspension culture. The maximum cell density of approximately 2 x 10<sup>6</sup> cells per ml is reached within 5 days of culture.

#### F. Growth characteristics of PER.C6 and PER.C6/E2A at 37°C and 39°C.

[0284] PER.C6 cells or PER.C6ts125E2A (c8-4) cells were cultured in Dulbecco's Modified Eagle Medium (DMEM, Gibco BRL) supplemented with 10% Fetal Bovine Serum (FBS, Gibco BRL) and 10mM MgCl<sub>2</sub> in a 10% CO<sub>2</sub> atmosphere at either 37°C (PER.C6) or 39°C (PER.C6ts125E2A c8-4). At day 0, a total of 1 x 10<sup>6</sup> cells were seeded per 25cm<sup>2</sup> tissue culture flask (Nunc) and the cells were cultured at the respective temperatures. At the indicated time points, cells were counted. The growth of PER.C6 cells at 37°C was comparable to the growth of PER.C6ts125E2A c8-4 at 39°C (Figure 33).

[0285] This shows that constitutive expression of ts125E2A encoded DBP had no adverse effect on the growth of cells at the non-permissive temperature of 39°C.

#### G. Stability of PER.C6ts125E2A

[0286] For several passages, the PER.C6ts125E2A cell line clone 8-4 was cultured at 39°C and 10% CO<sub>2</sub> in a 25 cm<sup>2</sup> tissue culture flask (Nunc) in DMEM, supplemented with 10% FBS and 10 mM MgCl<sub>2</sub> in the absence of selection pressure (G418). At sub-confluency (70-80% confluent), the cells were washed with PBS (NPBI) and lysed and scraped in RIPA (1% NP-40, 0.5% sodium deoxycholate and 0.1% SDS in PBS, supplemented with 1mM phenylmethylsulfonylfluoride and 0.1 mg/ml trypsin inhibitor). After 15 minutes incubation on ice, the lysates were cleared by centrifugation. Protein concentrations were determined by the BioRad protein assay, according to standard procedures of the supplier (BioRad). Equal amounts of whole-cell extract were fractionated by SDS-PAGE in 10% gels. Proteins were transferred onto Immobilon-P membranes (Millipore) and incubated with the αDBP monoclonal antibody B6. The secondary antibody was a horseradish-peroxidase-conjugated goat anti mouse antibody (BioRad). The Western blotting procedure and incubations were performed according to the protocol provided by Millipore. The complexes were visualized with the ECL detection system according to the manufacturer's protocol (Amersham). The expression of ts125E2A encoded DBP was stable for at least 16 passages, which is equivalent to approximately 40 cell doublings (Figure 34). No decrease in DBP levels was observed during this culture period, indicating that the expression of ts125E2A was stable, even in the absence of G418 selection pressure.

#### Example 16

#### Generation of tTA expressing packaging cell lines

##### A. Generation of a plasmid from which the tTA gene is expressed.

[0287]

pcDNA3.1-tTA: The tTA gene, a fusion of the tetR and VP16 genes, was removed from the plasmid pUHD 15-1 (Gossen and Bujard, 1992) by digestion using the restriction enzymes *Bam*HI and *Eco*RI. First, pUHD15-1 was digested with *Eco*RI.

The linearized plasmid was treated with Klenow enzyme in the presence of dNTPs to fill in the *Eco*RI sticky ends. Then, the plasmid was digested with *Bam*HI. The resulting fragment, 1025 bp in length, was purified from agarose. Subsequently, the fragment was used in a ligation reaction with *Bam*HI/*Eco*RV digested pcDNA 3.1 HYGRO (-) (Invitrogen) giving rise to pcDNA3.1-tTA. After transformation into competent *E. Coli* DH5α (Life Techn.) and analysis of ampicillin resistant colonies, one clone was selected that showed a digestion pattern as expected for pcDNA3.1-tTA.

*B. Transfection of PER.C6 and PER.C6/E2A with the tTA expression vector; colony formation and generation of cell lines*

[0288] One day prior to transfection,  $2 \times 10^6$  PER.C6 or PER.C6/E2A cells were seeded per 60 mm tissue culture dish (Greiner) in Dulbecco's modified essential medium (DMEM, Gibco BRL) supplemented with 10% FBS (JRH) and 10 mM  $MgCl_2$  and incubated at 37°C in a 10%  $CO_2$  atmosphere. The next day, cells were transfected with 4-8  $\mu g$  of pcDNA3.1-tTA plasmid DNA using the LipofectAMINE PLUS™ Reagent Kit according to the standard protocol of the supplier (Gibco BRL). The cells were incubated with the LipofectAMINE PLUS™-DNA mixture for four hours at 37°C and 10%  $CO_2$ . Then, 2 ml of DMEM supplemented with 20% FBS and 10 mM  $MgCl_2$  was added and cells were further incubated at 37°C and 10%  $CO_2$ . The next day, cells were washed with PBS and incubated in fresh DMEM supplemented with 10% FBS, 10 mM  $MgCl_2$  at either 37°C (PER.C6) or 39°C (PER.C6/E2A) in a 10%  $CO_2$  atmosphere for three days. Then, the media were exchanged for selection media; PER.C6 cells were incubated with DMEM supplemented with 10% PBS, 10 mM  $MgCl_2$  and 50  $\mu g/ml$  hygromycin B (GIBCO) while PER.C6/E2A cells were maintained in DMEM supplemented with 10% PBS, 10 mM  $MgCl_2$  and 100  $\mu g/ml$  hygromycin B.

[0289] Colonies of cells that resisted the selection appeared within three weeks while nonresistant cells died during this period.

[0290] From each transfection, a number of independent, hygromycin resistant cell colonies were picked by scraping the cells from the dish with a pipette and put into 2.5  $cm^2$  dishes (Greiner) for further growth in DMEM containing 10% FBS, 10 mM  $MgCl_2$  and supplemented with 50  $\mu g/ml$  (PER.C6 cells) or 100  $\mu g/ml$  (PER.C6/E2A cells) hygromycin in a 10%  $CO_2$  atmosphere and at 37°C or 39°C, respectively.

[0291] Next, it was determined whether these hygromycin-resistant cell colonies expressed functional tTA protein.

[0292] Therefore, cultures of PER.C6/tTA or PER/E2A/tTA cells were transfected with the plasmid pUHC 13-3 that contains the reporter gene luciferase under the control of the 7xtetO promoter (Gossens and Bujard, 1992). To demonstrate that the expression of luciferase was mediated by tTA, one half of the cultures was maintained in medium without doxycycline.

[0293] The other half was maintained in medium with 8  $\mu g/ml$  doxycycline (Sigma). The latter drug is an analogue of tetracycline and binds to tTA and inhibits its activity. All PER.C6/tTA and PER/E2A/tTA cell lines yielded high levels of luciferase, indicating that all cell lines expressed the tTA protein (Figure 35). In addition, the expression of luciferase was greatly suppressed when the cells were treated with doxycycline. Collectively, the data showed that the isolated and established hygromycin-resistant PER.C6 and PER/E2A cell clones all expressed functional tTA.

Legend to the figures:

[0294]

Figure 1:

Bar graph showing the percentage of serum samples positive for neutralisation for each human wt adenovirus tested (see example 1 for description of the neutralisation assay).

Figure 2:

Graph showing absence of correlation between the VP/CCID50 ratio and the percentage of neutralisation.

Figure 3:

Schematic representation of a partial restriction map of Ad35 (taken from Kang *et al.*, 1989) and the clones generated to make recombinant Ad35-based viruses.

Figure 4: Bar graph presenting the percentage sera samples that show neutralizing activity to a selection of adenovirus serotypes. Sera were derived from healthy volunteers from Belgium and the UK.

Figure 5: Bar graph presenting the percentage sera samples that show neutralizing activity to adenovirus serotypes 5, 11, 26, 34, 35, 48 and 49. Sera were derived from five different locations in Europe and the United States.

Figure 6: Sequence of human adenovirus type 35. As explained in the text the nucleotide sequence of the terminal ends of the virus are not definite resolved.

Figure 7: Map of pAdApt

Figure 8: Map of pIPspAdapt



Figure 9: Map of pIPspAdapt1

Figure 10: Map of pIPspAdapt3

5 Figure 11: Map of pAdApt35IP3

Figure 12: Map of pAdApt35IP1

10 Figure 13: Schematic representation of the steps undertaken to construct pWE.Ad35.pIX-rITR

Figure 14: Map of pWE.Ad35.pIX-rITR

Figure 15: Map of pRSV.Ad35-E1

15 Figure 16: Map of PGKneopA

20 Figure 17: Map of pRSVpNeo Figure 18: Map of pRSVhvbNeo

Figure 19: Flow cytometric analyses on Green fluorescent protein (GFP) expression in human TF-1 cells. Non-transduced TF-1 cells were used to set a background level of 1%. GFP expression in cells transduced with Ad5, Ad5.Fib16, Ad5.Fib17, Ad5.Fib40-L, Ad5.Fib35, and Ad5.Fib51 is shown.

25 Figure 20: Transduction of primary human fibroblast-like stroma. Cells were analysed 48 hours after a two hour exposure to the different chimaeric fiber viruses. Shown is percentage of cells found positive for the transgene: green fluorescent protein (GFP) using a flow cytometer. Non-transduced stroma cells were used to set a background at 1%. Results of different experiments (n=3) are shown  $\pm$  standard deviation.

30 Figure 21: Transduction of primary human fibroblast-like stroma, CD34<sup>+</sup> cells and CD34<sup>+</sup>Lin<sup>-</sup> cells. Cells were analyzed 5 days after a two hour exposure to the different chimaeric fiber viruses. Shown is percentage of cells found positive for the transgene: green fluorescent protein (GFP) using a flow cytometer. Non-transduced cells were used to set a background at 1%. Also shown is the number of GFP positive events divided by the total number of events analysed (between brackets).

35 Figure 22 A) Flow cytometric analysis of GFP positive cells after transduction of CD34<sup>+</sup> cells with Ad5.Fib51. All cells gated in R2-R7 are positive for CD34 but differ in their expression of early differentiation markers CD33, CD38, and CD71 (Lin). Cells in R2 are negative for CD333, CD38, CD71 whereas cells in R7 are positive for these markers. To demonstrate specificity of Ad5.Fib51 the percentage of GFP positive cells was determined in R2-R7 which  
40 proofed to decline from 91% (R2) to 15% (R7). B) Identical experiment as shown under A (X-axis is R2-R7) but for the other Ad fiber chimaeric viruses showing that Ad5.Fib35, and Ad5.Fib16 behave similar as Ad5.Fib51.

Figure 23: Alignment of the chimeric fiber proteins of Ad5fib16, Ad5fib35 and Ad5fib51 with the Ad5 fiber sequence.

45 Figure 24: Toxicity of Adenovirus exposure to primitive human Bone marrow cells and Stem cells. Cell cultures were counted just before and 5 days after adenovirus transduction. Shown is the percentage of primitive human bone marrow cells (CD34<sup>+</sup>) and HSCs (CD34<sup>+</sup>Lin<sup>-</sup>) recovered as compared to day 0.

50 Figure 25: Transduction of immature DCs at a virus dose of 100 or 1000 virus particles per cell. Virus tested is Ad5 and Ad5 based vectors carrying the fiber of serotype 12 (Ad5.Fib12), 16 (Ad5.Fib16), 28 (Ad5.Fib28), 32 (Ad5.Fib32), the long fiber of 40 (Ad5.Fib40-L, 49 (Ad5.Fib49), 51 (Ad5.Fib51). Luciferase transgene expression is expressed as relative light units per microgram of protein.

55 Figure 26: Flow cytometric analyses of LacZ expression on immature and mature DCs transduced with 10000 virus particles per cell of Ad5 or the fiber chimaeric vectors Ad5.Fib16, Ad5.Fib40-L, or Ad5.Fib51. Percentages of cells scored positive are shown in upper left corner of each histogram.

Figure 27: Luciferase transgene expression in human immature DCs measured 48 hours after transduction with

1000 or 5000 virus particles per cell. Virus tested were fiber chimaeric viruses carrying the fiber of subgroup B members (serotypes 11, 16, 35, and 51).

Figure 28: Green fluorescent protein (GFP) expression in immature human DCs 48 hours after transduction with 1000 virus particles per cell of Ad5, Ad5.Fib16, and Ad5.Fib35.

Non-transduced cells were used to set a background level of approximately 1% (-).

Figure 29: Transduction of mouse and chimpanzee DCs.

Luciferase transgene expression measured in mouse DCs 48 hours after transduction is expressed as relative light units per microgram of protein. Chimpanzee DCs were measured 48 hours after transduction using a flow cytometer. GFP expression demonstrates the poor transduction of Ad (35) in contrast to Ad5.Fib35 (66%).

Figure 30: Temperature dependent growth of PER.C6.

PER.C6 cells were cultured in Dulbecco's Modified Eagle Medium supplemented with 10% Fetal Bovine Serum (FBS, Gibco BRL) and 10mM MgCl<sub>2</sub> in a 10% CO<sub>2</sub> atmosphere at either 32°C, 37°C or 39°C. At day 0, a total of 1 x 10<sup>6</sup> PER.C6 cells were seeded per 25cm<sup>2</sup> tissue culture flask (Nunc) and the cells were cultured at either 32°C, 37°C or 39°C. At day 1-8, cells were counted. The growth rate and the final cell density of the PER.C6 culture at 39°C are comparable to that at 37°C.

The growth rate and final density of the PER.C6 culture at 32°C were slightly reduced as compared to that at 37°C or 39°C.

PER.C6 cells were seeded at a density of 1 x 10<sup>6</sup> cells per 25 cm<sup>2</sup> tissue culture flask and cultured at either 32-, 37- or 39°C. At the indicated time points, cells were counted in a Burkert cell counter. PER.C6 grows well at both 32-, 37- and 39°C.

Figure 31: DBP levels in PER.C6 cells transfected with pcDNA3, pcDNA3wtE2A or pcDNA3ts125E2A.

Equal amounts of whole-cell extract were fractionated by SDS-PAGE on 10% gels. Proteins were transferred onto Immobilon-P membranes and DBP protein was visualized using the αDBP monoclonal B6 in an ECL detection system. All of the cell lines derived from the pcDNA3ts125E2A transfection express the 72-kDa E2A-encoded DBP protein (left panel, lanes 4-14; middle panel, lanes 1-13; right panel, lanes 1-12). In contrast, the only cell line derived from the pcDNAwtE2A transfection did not express the DBP protein (left panel, lane 2). No DBP protein was detected in extract from a cell line derived from the pcDNA3 transfection (left panel, lane 1), which serves as a negative control. Extract from PER.C6 cells transiently transfected with pcDNA3ts125 (left panel, lane 3) served as a positive control for the Western blot procedure. These data confirm that constitutive expression of wtE2A is toxic for cells and that using the ts125 mutant of E2A can circumvent this toxicity.

Figure 32: Suspension growth of PER.C6ts125E2A C5-9.

The tsE2A expressing cell line PER.C6tsE2A.c5-9 was cultured in suspension in serum free Ex-cell™. At the indicated time points, cells were counted in a Burkert cell counter. The results of 8 independent cultures are indicated. PER.C6tsE2A grows well in suspension in serum free Ex-cell™ medium.

Figure 33: Growth curve PER.C6 and PER.C6tsE2A.

PER.C6 cells or PER.C6ts125E2A (c8-4) cells were cultured at 37°C or 39°C, respectively. At day 0, a total of 1 x 10<sup>6</sup> cells was seeded per 25cm<sup>2</sup> tissue culture flask. At the indicated time points, cells were counted. The growth of PER.C6 cells at 37°C is comparable to the growth of PER.C6ts125E2A c8-4 at 39°C. This shows that constitutive overexpression of ts125E2A has no adverse effect on the growth of cells at the non-permissive temperature of 39°C.

Figure 34: Stability of PER.C6ts125E2A.

For several passages, the PER.C6ts125E2A cell line clone 8-4 was cultured at 39°C in medium without G418. Equal amounts of whole-cell extract from different passage numbers were fractionated by SDS-PAGE on 10% gels. Proteins were transferred onto Immobilon-P membranes and DBP protein was visualized using the αDBP monoclonal B6 in an ECL detection system. The expression of ts125E2A encoded DBP is stable for at least 16 passages, which is equivalent to approximately 40 cell doublings. No decrease in DBP levels were observed during this culture period, indicating that the expression of ts125E2A is stable, even in the absence of G418 selection pressure.

Figure 35: tTA activity in hygromycin resistant PER.C6/tTA (A) and PER/E2A/tTA (B) cells.

Sixteen independent hygromycin resistant PER.C6/tTA cell colonies and 23 independent hygromycin resistant

PER/E2A/tTA cell colonies were grown in 10 cm<sup>2</sup> wells to sub-confluency and transfected with 2 µg of pUHC 13-3 (a plasmid that contains the reporter gene luciferase under the control of the 7xtetO promoter). One half of the cultures was maintained in medium containing doxycycline to inhibit the activity of tTA. Cells were harvested at 48 hours after transfection and luciferase activity was measured. The luciferase activity is indicated in relative light units (RLU) per µg protein.

Table I:

Serotype	Elution [NaCl] mM	VP/ml	CCID50	log <sub>10</sub> VP/CCID50 ratio
1	597	8.66x10 <sup>10</sup>	5.00x10 <sup>7</sup>	3.2
2	574	1.04x10 <sup>12</sup>	3.66x10 <sup>11</sup>	0.4
3	131	1.19x10 <sup>11</sup>	1.28x10 <sup>7</sup>	4.0
4	260	4.84x10 <sup>11</sup>	2.50x10 <sup>8</sup>	3.3
5	533	5.40x10 <sup>11</sup>	1.12x10 <sup>10</sup>	1.7
6	477	1.05x10 <sup>12</sup>	2.14x10 <sup>10</sup>	1.7
7	328	1.68x10 <sup>12</sup>	2.73x10 <sup>9</sup>	2.4
9	379	4.99x10 <sup>11</sup>	3.75x10 <sup>7</sup>	4.1
10	387	8.32x10 <sup>12</sup>	1.12x10 <sup>9</sup>	3.9
12	305	3.64x10 <sup>11</sup>	1.46x10 <sup>7</sup>	4.4
13	231	4.37x10 <sup>12</sup>	7.31x10 <sup>8</sup>	3.8
15	443	5.33x10 <sup>12</sup>	1.25x10 <sup>9</sup>	3.6
16	312	1.75x10 <sup>12</sup>	5.59x10 <sup>8</sup>	3.5
17	478	1.39x10 <sup>12</sup>	1.45x10 <sup>9</sup>	3.0
19	430	8.44x10 <sup>11</sup>	8.55x10 <sup>7</sup>	4.0
20	156	1.41x10 <sup>11</sup>	1.68x10 <sup>7</sup>	3.9
21	437	3.21x10 <sup>11</sup>	1.12x10 <sup>8</sup>	3.5
22	385	1.43x10 <sup>10</sup>	5.59x10 <sup>10</sup>	3.2
23	132	2.33x10 <sup>11</sup>	1.57x10 <sup>7</sup>	4.2
24	405	5.12x10 <sup>12</sup>	4.27x10 <sup>8</sup>	4.1
25	405	7.24x10 <sup>11</sup>	5.59x10 <sup>7</sup>	4.1
26	356	1.13x10 <sup>12</sup>	1.12x10 <sup>8</sup>	4.0
27	342	2.00x10 <sup>12</sup>	1.28x10 <sup>8</sup>	4.2
28	347	2.77x10 <sup>12</sup>	5.00x10 <sup>7</sup>	4.7
29	386	2.78x10 <sup>11</sup>	2.00x10 <sup>7</sup>	4.1
30	409	1.33x10 <sup>12</sup>	5.59x10 <sup>9</sup>	3.4
31	303	8.48x10 <sup>10</sup>	2.19x10 <sup>7</sup>	3.6
33	302	1.02x10 <sup>12</sup>	1.12x10 <sup>7</sup>	5.0
34	425	1.08x10 <sup>12</sup>	1.63x10 <sup>11</sup>	0.8
35	446	3.26x10 <sup>12</sup>	1.25x10 <sup>11</sup>	1.4
36	325	9.26x10 <sup>12</sup>	3.62x10 <sup>9</sup>	3.4
37	257	5.86x10 <sup>12</sup>	2.8x10 <sup>9</sup>	3.3
38	337	3.61x10 <sup>12</sup>	5.59x10 <sup>7</sup>	4.8
39	241	3.34x10 <sup>11</sup>	1.17x10 <sup>7</sup>	4.5
42	370	1.95x10 <sup>12</sup>	1.12x10 <sup>8</sup>	4.2

Continued on next page.

Serotype #	Elution [NaCl] mM	VP/ml	CCID50	log <sub>10</sub> VP/CCID50 ratio
43	284	2.42x10 <sup>12</sup>	1.81x10 <sup>8</sup>	4.1
44	295	8.45x10 <sup>11</sup>	2.00x10 <sup>7</sup>	4.6
45	283	5.20x10 <sup>11</sup>	2.99x10 <sup>7</sup>	4.2
46	282	9.73x10 <sup>12</sup>	2.50x10 <sup>8</sup>	4.6
47	271	5.69x10 <sup>11</sup>	3.42x10 <sup>7</sup>	4.2
48	264	1.68x10 <sup>12</sup>	9.56x10 <sup>8</sup>	3.3
49	332	2.20x10 <sup>12</sup>	8.55x10 <sup>7</sup>	4.4
50	459	7.38x10 <sup>12</sup>	2.80x10 <sup>9</sup>	3.4
51	450	8.41x10 <sup>11</sup>	1.88x10 <sup>8</sup>	3.7

Legend to table I:

[0295] All human adenoviruses used in the neutralisation experiments were produced on PER.C6 cells (ECACC deposit number 96022940) (Fallaux *et al.*, 1998) and purified on CsCl as described in example 1. The NaCl concentration at which the different serotypes eluted from the HPLC column is shown. Virus particles/ml (VP/ml) were calculated from an Ad5 standard. The titer in the experiment (CCID50) was determined on PER.C6 cells (ECACC deposit number 96022940) as described in example 1 by titrations performed in parallel with the neutralisation experiment. The CCID50 is shown for the 44 viruses used in this study and reflects the dilution of the virus needed to obtain CPE in 50% of the wells after 5 days. The ratio of VP/CCID50 is depicted in log<sub>10</sub> and is a measurement of the infectivity of the different batches on PER.C6 cells (ECACC deposit number 96022940).

Table II

AdApt35.LacZ viruses escape neutralization by human serum.						
Virus	Human serum dilution					
	no serum	10x	50x	250x	1250x	6250x
AdApt5.LacZ moi: 5 VP/cell	100 %	0 %	0 %	1 %	40 %	80 %
AdApt35.LacZ 250 µl crude lysate	100 %	100 %	100 %	100 %	100 %	100 %

Table III

Percentage of synovial fluid samples containing neutralizing activity (NA) to wt adenoviruses of different serotypes.		
	% of SF samples with NA (all positives)	% of SF samples with NA (positives at ≥64x dilution)
Ad5	72	59
Ad26	66	34
Ad34	45	19
Ad35	4	0
Ad48	42	4



Table IV

The numbers of foci obtained with the different E1 expression constructs in BRK transformation experiments.			
Average # of foci/dish:			
	Construct	1 µgr	5 µgr
Experiment 1	pIG.E1A.E1 B	nd	60
	pIG.E1A.E1 B	nd	35
	pRSVAd35E1	0	3
	pIG.Ad35.E 1	3	7
Experiment 2	pIG.E1A.E1 B	37	nd
	pIG.Ad35.E 1	nd	2
Experiment 3	pIG.E1A.E1 B	nd	140
	pIG.Ad35.E 1	nd	20
	pIG270	nd	30

## REFERENCES

[0296]

Abrahamsen, K., Kong, H-L., Mastrangeli, A., Brough, D., Lizonova, A., Crystal, R. and Falck-Pedersen, E. (1997) Construction of an adenovirus type 7a E1A vector. *J. Virol.* 71, no. 11, p8946-8951.

Athappilly, F.K., Murali, R., Rux, J.J., Cai, Z. and Burnett, R.M. (1994). The refined crystal structure of hexon, the major coat protein of adenovirus type 2, at 2.9 Å resolution. *J. Mol. Biol.* 242, 430-455.

Basler, C.F., Droguett, G., Horwitz, M.S. (1996). Sequence of the immunoregulatory early region 3 and flanking sequences of adenovirus type 35. *Gene* 170:249-54

Bridge, E., Medghalchi, S., Ubol, S., Leesong, M. and Ketner, G. (1993) Adenovirus early region 4 and viral DNA synthesis. *Virology* 193, 794-801.

Brody, S.L. and Crystal, R.G. (1994) Adenovirus mediated in vivo gene transfer. *Ann. N. Y. Acad. Sci.* 716:90-101.

Boshart, M., Weber, F., Jahn, G., Dorsch-Hasler, K., Fleckenstein B. and Schaffner W. (1985). A very strong enhancer is located upstream of an immediate early gene of human cytomegalovirus. *Cell* 41, 521-530,

Dijkema, R., Dekker, B.M.M., van der Feltz, M.J.M. and van der Eb, A.J. (1979). Transformation of primary rat kidney cells by DNA fragments of weakly oncogenic adenoviruses. *J. Virol.* 32, No 3, 943-950.

Fallaux, F.J., Bout, A., van der Velde, I., van den Wollenberg, D.J., Hehir, K.M., Keegan, J., Auger, C., Cramer, S.J., van Ormondt, H., van der Eb, A.J., Valerio, D. and Hoebe, R.C. (1998). New helper cells and matched early region 1-deleted adenovirus vectors prevent generation of replication competent adenoviruses. *Hum. Gene Ther.* 9, 1909-1917.

Gossen, M., and H. Bujard (1992) Tight control of gene expression in mammalian cells by tetracyclin-responsive promoters. *Proc. Natl. Acad. Sci. USA* 89; 5547-5551.

Flomenberg, P.R., Chen, M., Munk, G., Horwitz, M.S. (1987). Molecular epidemiology of adenovirus type 35 infections in immunocompromised hosts. *J. Infect Dis.* 155(6): 1127-34.

5 Francki, R.I.B., Fauquet, C.M., Knudson, D.L. and Brown, F. (1991). Classification and nomenclature of viruses. Fifth report of the international Committee on taxonomy of viruses. *Arch. Virol. Suppl.* 2: 140-144.

Gahery-Segard, H., Farace, F., Godfrin, D., Gaston, J., Lengagne, R., Tursz, P., Boulanger, P. and Guillet, J.-G. (1998). Immune response to recombinant capsid proteins of adenovirus in humans: Antifiber and anti-penton base antibodies have a synergistic effect on neutralising activity. *J. Virol.* 72, 2388-2397.

10 He, T.-C., Zhou, S., Da Costa, L.T., Yu, J., Kinzler, K.W., Vogelstein, B. (1998). A simplified system for generating recombinant adenoviruses. *Proc. Natl. Acad. Sci. USA* 95, 2509-2514.

15 Hierholzer, J.C., Wigand, R., Anderson, L.J., Adrian, T., and Gold, J.W.M. (1988) Adenoviruses from patients with AIDS: a plethora of serotypes and a description of five new serotypes of subgenus D (types 43-47). *J. Infect. Dis.* 158, 804-813.

20 De Jong, P.J., Valderrama, G., Spigland, I. and Horwitz, M.S. (1983). Adenovirus isolates from urine of patients with acquired immunodeficiency syndrome. *Lancet* 1(8337): 1293-1296.

Kay, R., Takei, F., and Humphries, R.K. (1990). Expression cloning of a cDNA encoding M1/69. *J. Immunol.* 145, 1952-1959.

25 Kang, W.G., Berencsi, G., Takacs, M., Ascher, Z., Fejer, G., Nasz, I. (1989a). Molecular cloning and physical mapping of the DNA of human adenovirus type 35. *Acta Microbiol Hung* 36(1): 67-75.

Kang, W.G., Berencsi, G., Banrevi, A., Ascher, Z., Fejer, G., Takacs, M., Kiss, A., Nasz, I. (1989b). Relationship of E1 and E3 regions of human adenovirus 35 to those of human adenovirus subgroups A, C and D. *Acta Microbiol Hung* 36(4): 445-57.

30 Levrero, M., Barban, V., Manteca, S., Ballay, A., Balsamo, C., Avantaggiati, M.L., Natoli, G., Skellekens, H., Tiollais, P., and Perricaudet, M. (1991). Defective and nondefective adenovirus vectors for expressing foreign genes in vitro and in vivo. *Gene* 101, 195-202.

35 Li, Q.G., Hambræus, J. and Wadell, G. (1991). Genetic relationship between thirteen genomes of types of adenovirus 11, 34, and 35 with different tropisms. *Interviol.* 32, 338-350.

Prince, H.M. (1998). Gene transfer: a review of methods and applications. *Pathology* 30(4), 335-347.

40 Robbins, P.D. and Ghivizzani, S.C. (1998). Viral vectors for gene therapy. *Pharmacol Ther.* 80, 35-47.

Schnurr, D and Dondero, M.E. (1993). Two new candidate adenovirus serotypes. *Interviol.* 36, 79-83.

45 Schulick, A.H., Vassalli, G., Dunn, P.F., Dong, G., Rade, J.J., Zamarron, C. and Dichek, D.A. (1997). Established immunity precludes adenovirus-mediated gene transfer in rat carotid arteries. Potential for immunosuppression and vector engineering to overcome barriers of immunity. *J. Clin. Invest.* 99(2), 209-19.

50 Shabram, P.W., Giroux, D.D., Goudreau, A.M., Gregory, R.J., Horn, M.T., Huyghe, B.G., Liu, X., Nunnally, M.H., Sugarman, B.J. and Sutjipto, S. (1997) Analytical anion-exchange HPLC of recombinant type-5 adenoviral particles. *Hum. Gene Ther.* 8(4): 453-465.

Toogood et al., 1989; *J. Gen Virol.* 70, 3203-3214

55 Toogood, C.I., Murali, R., Burnett, R.M., Hay, R.T. (1989). The adenovirus type 40 hexon: sequence, predicted structure and relationship to other adenovirus hexons. *J. Gen. Virol.* 70, 3203-14.

Valderrama-Leon, G., Flomenberg, P., Horwitz, M.S. (1985). Restriction endonuclease mapping of adenovirus 35, a type isolated from immunocompromised hosts. *J Virol.* 56(2):647-50.

EP 1 054 064 A1

Wadell, G. (1984). Molecular epidemiology of adenoviruses. Curr. Top. Microbiol. Immunol. 110, 191-220.

White, E. (1995) Regulation of p53-dependent apoptosis by E1a and E1b. Curr. Top. Microbiol. Immunol. 199, 34-58.

5

SEQUENCE LISTING

10

<110> Introgene B.V.

<120> Serotype of Adenovirus and uses Thereof

<130> P49636EP10

15

<140> 00201738.2

<141> 2000-05-16

<150> EP 99201545.3

<151> 1999-05-17

20

<160> 88

<170> PatentIn Ver. 2.1

25

<210> 1

<211> 23

<212> DNA

<213> Artificial Sequence

30

<220>

<223> Description of Artificial Sequence: linker  
containing a PacI site

35

<220>

<221> misc\_feature

<222> (1)..(23)

<400> 1

aattgtctta attaaccgct taa

23

40

<210> 2

<211> 19

<212> DNA

<213> Artificial Sequence

45

<220>

<223> Description of Artificial Sequence: oligonucleotide

50

<220>

<221> misc\_feature

<222> (1)..(19)

<400> 2

aattgtctta attaaccgc

19

55

<210> 3

5  
 <211> 19  
 <212> DNA  
 <213> Artificial Sequence

10  
 <220>  
 <223> Description of Artificial Sequence: oligonucleotide

15  
 <220>  
 <221> misc\_feature  
 <222> (1)..(19)

<400> 3  
 aattgcggtt aattaagac 19

20  
 <210> 4  
 <211> 39  
 <212> DNA  
 <213> Artificial Sequence

25  
 <220>  
 <223> Description of Artificial Sequence: primer ITR-EPH

30  
 <220>  
 <221> misc\_feature  
 <222> (1)..(39)

<400> 4  
 cggaattctt aattaagtta acatcatcaa taatatacc 39

35  
 <210> 5  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

40  
 <220>  
 <223> Description of Artificial Sequence: primer Ad101

45  
 <220>  
 <221> misc\_feature  
 <222> (1)..(20)

50  
 <400> 5  
 tgattcacat cggtcagtgc 20

55  
 <210> 6  
 <211> 23  
 <212> DNA  
 <213> Artificial Sequence



<220>  
<223> Description of Artificial Sequence: primer delta E2A.SnaBI

5  
<220>  
<221> misc\_feature  
<222> (1)..(23)

<400> 6  
10 ggcgtacgta gccctgtcga aag 23

<210> 7  
<211> 35  
15 <212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer delta E2A.DBP-start

20  
<220>  
<221> misc\_feature  
<222> (1)..(35)

<400> 7  
25 ccaatgcatt cgaagtactt ccttctecta taggc 35

<210> 8  
30 <211> 22  
<212> DNA  
<213> Artificial Sequence

<220>  
35 <223> Description of Artificial Sequence: primer delta E2A.DBP-stop

<220>  
<221> misc\_feature  
<222> (1)..(22)

40  
<400> 8  
ccaatgcata cggcgcagac gg 22

<210> 9  
45 <211> 21  
<212> DNA  
<213> Artificial Sequence

<220>  
50 <223> Description of Artificial Sequence: primer delta E2A.BamHI

<220>

55

<221> misc\_feature

<222> (1)..(21)

5

<400> 9

gaggtggatc ccatggacga g

21

10

<210> 10

<211> 47

<212> DNA

<213> Artificial Sequence

15

<220>

<223> Description of Artificial Sequence: primer LTR-1

<220>

<221> misc\_feature

20

<222> (1)..(47)

<400> 10

ctgtacgtac cagtgcactg gcctaggcat ggaaaaatac ataactg

47

25

<210> 11

<211> 64

<212> DNA

<213> Artificial Sequence

30

<220>

<223> Description of Artificial Sequence: primer LTR-2

<220>

35

<221> misc\_feature

<222> (1)..(64)

<400> 11

gcggatcctt cgaaccatgg taagcttggt accgctagcg ttaaccgggc gactcagtc 60  
atcg 64

40

<210> 12

<211> 28

<212> DNA

45

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer HSA1

50

<220>

<221> misc\_feature

<222> (1)..(28)

55

5  
 <400> 12  
 gcgccacat gggcagagcg atggtggc 28

10  
 <210> 13  
 <211> 50  
 <212> DNA  
 <213> Artificial Sequence

15  
 <220>  
 <223> Description of Artificial Sequence: primer HSA2

20  
 <220>  
 <221> misc\_feature  
 <222> (1)..(50)

25  
 <400> 13  
 gttagatcta agcttgctga catcgatcta ctaacagtag agatgtagaa 50

30  
 <210> 14  
 <211> 10  
 <212> DNA  
 <213> Artificial Sequence

35  
 <220>  
 <223> Description of Artificial Sequence: oligonucleotide

40  
 <220>  
 <221> misc\_feature  
 <222> (1)..(10)

45  
 <400> 14  
 ttaagtcgac 10

50  
 <210> 15  
 <211> 23  
 <212> DNA  
 <213> Artificial Sequence

55  
 <220>  
 <223> Description of Artificial Sequence: linker

55  
 <220>  
 <221> misc\_feature  
 <222> (1)..(23)

55  
 <400> 15  
 aattgtctta attaaccgca att 23

5  
 <210> 16  
 <211> 67  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: oligonucleotide PLL-1

10  
 <220>  
 <221> misc\_feature  
 <222> (1)..(67)

15  
 <400> 16  
 gccatcccta ggaagcttgg taccggtgaa ttcgctagcg ttaacggatc ctctagacga 60  
 gatctgg 67

20  
 <210> 17  
 <211> 67  
 <212> DNA  
 <213> Artificial Sequence

25  
 <220>  
 <223> Description of Artificial Sequence: oligonucleotide PLL-2

30  
 <220>  
 <221> misc\_feature  
 <222> (1)..(67)

35  
 <400> 17  
 ccagatctcg tctagaggat ccgttaacgc tagcgaattc accggtacca agcttcctag 60  
 ggatggc 67

40  
 <210> 18  
 <211> 39  
 <212> DNA  
 <213> Artificial Sequence

45  
 <220>  
 <223> Description of Artificial Sequence: primer CMVplus

50  
 <220>  
 <221> misc\_feature  
 <222> (1)..(39)

55  
 <400> 18  
 gatcggtacc actgcagtgg tcaatattgg ccattagcc 39



5  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: primer CMVminA  
 <220>  
 <221> misc\_feature  
 <222> (1)..(29)  
 <400> 19  
 gatcaagctt ccaatgcacc gttcccggc 29  
 15  
 <210> 20  
 <211> 32  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: primer  
 <220>  
 <221> misc\_feature  
 <222> (1)..(32)  
 <400> 20  
 ggggtggcca gggtagctct aggcttttgc aa 32  
 30  
 <210> 21  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: primer  
 <220>  
 <221> misc\_feature  
 <222> (1)..(29)  
 <400> 21  
 ggggggatcc ataaacaagt tcagaatcc 29  
 45  
 <210> 22  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 55

<223> Description of Artificial Sequence: primer delta hex1

<220>

<221> misc\_feature

<222> (1)..(18)

<400> 22

cctggtgctg ccaacagc

18

<210> 23

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer delta hex2

<220>

<221> misc\_feature

<222> (1)..(30)

<400> 23

ccggatccac tagtggaag cgggcgcgcg

30

<210> 24

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer delta hex3

<220>

<221> misc\_feature

<222> (1)..(35)

<400> 24

ccggatccaa ttgagaagca agcaacatca acaac

35

<210> 25

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer delta hex4

<220>

<221> misc\_feature

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

<222> (1)..(19)

<400> 25  
gagaagggca tggaggctg 19

<210> 26  
<211> 32  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer Hex-up2

<220>  
<221> misc\_feature  
<222> (1)..(32)

<400> 26  
gactagtcaa gatggcyacc chtcgatga tg 32

<210> 27  
<211> 33  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer Hex-do2

<220>  
<221> misc\_feature  
<222> (1)..(33)

<400> 27  
gctggccaat tggtatgtkg tkgcgttrcc ggc 33

<210> 28  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: primer DP5-F

<220>  
<221> misc\_feature  
<222> (1)..(20)

<400> 28  
ctgttgctgc tgctaatagc 20

5  
 <210> 29  
 <211> 32  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: primer DP5-R  
 10  
 <220>  
 <221> misc\_feature  
 <222> (1)..(32)  
 15  
 <400> 29  
 cgcggtatcct gtacaactaa ggggaataca ag 32  
 20  
 <210> 30  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: primer DP3-F  
 25  
 <220>  
 <221> misc\_feature  
 <222> (1)..(33)  
 30  
 <400> 30  
 cgcggtatccc ttaaggcaag catgtccatc ctt 33  
 35  
 <210> 31  
 <211> 27  
 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: primer DP3-3R  
 40  
 <220>  
 <221> misc\_feature  
 <222> (1)..(27)  
 45  
 <400> 31  
 aaaacacgtt ttacgcgtcg acctttc 27  
 50  
 <210> 32  
 <211> 33  
 55



5 <212> DNA  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: primer P3-for  
 <220>  
 10 <221> misc\_feature  
 <222> (1)..(33)  
 <400> 32  
 gctcgatgta caatgaggag acgagccgtg cta 33  
 15  
 <210> 33  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence  
 20 <220>  
 <223> Description of Artificial Sequence: primer P3-rev  
 <220>  
 25 <221> misc\_feature  
 <222> (1)..(33)  
 <400> 33  
 gctcgactta agttagaaag tgcggcttga aag 33  
 30  
 <210> 34  
 <211> 35  
 <212> DNA  
 <213> Artificial Sequence  
 35 <220>  
 <223> Description of Artificial Sequence: primer P17F  
 <220>  
 40 <221> misc\_feature  
 <222> (1)..(35)  
 <400> 34  
 45 gctcgatgta caatgaggcg tgcgggtgtg tcttc 35  
 <210> 35  
 <211> 34  
 <212> DNA  
 <213> Artificial Sequence  
 50 <220>  
 55

<223> Description of Artificial Sequence: primer P17R

<220>

<221> misc\_feature

<222> (1)..(34)

<400> 35

gctcgactta agttagaagg tgcgactgga aagc

34

<210> 36

<211> 42

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer NY-up

<220>

<221> misc\_feature

<222> (1)..(42)

<400> 36

cgacatatgt agatgcatta gtttgtgta tgtttcaacg tg

42

<210> 37

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer NY-down

<220>

<221> misc\_feature

<222> (1)..(18)

<400> 37

ggagaccact gccatgtt

18

<210> 38

<211> 10

<212> DNA

<213> Adenoviridae

<220>

<221> misc\_feature

<222> (1)..(10)

<223> /note="Partial sequence of an adenovirus ITR"

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

<400> 38  
catcatcaat 10

<210> 39  
<211> 14  
<212> DNA  
<213> Adenoviridae

<220>  
<221> misc\_feature  
<222> (1)..(14)  
<223> /note="Partial sequence of an adenovirus ITR"

<400> 39  
ccaataatat acct 14

<210> 40  
<211> 21  
<212> DNA  
<213> Adenoviridae

<220>  
<221> misc\_feature  
<222> (1)..(21)  
<223> /note="Partial sequence of an adenovirus ITR"

<400> 40  
aggtatatta ttgatgatgg g 21

<210> 41  
<211> 18  
<212> DNA  
<213> Adenoviridae

<220>  
<221> misc\_feature  
<222> (1)..(18)  
<223> /note="Partial sequence of an adenovirus ITR"

<400> 41  
catcatcaat aatatacc 18

<210> 42  
<211> 13  
<212> DNA  
<213> Adenoviridae

<220>

5  
 <221> misc\_feature  
 <222> (1)..(13)  
 <223> /note="Partial sequence of an adenovirus ITR"

<400> 42  
 catcatcaat aat 13

10  
 <210> 43  
 <211> 47  
 <212> DNA  
 <213> Artificial Sequence

15  
 <220>  
 <223> Description of Artificial Sequence: oligo ExSalPacF

20  
 <220>  
 <221> misc\_feature  
 <222> (1)..(47)

<400> 43  
 tcgatggcaa acagctatta tgggtattat gggttcgaat taattaa 47

25  
 <210> 44  
 <211> 47  
 <212> DNA  
 <213> Artificial Sequence

30  
 <220>  
 <223> Description of Artificial Sequence: oligo ExSalPacR

35  
 <220>  
 <221> misc\_feature  
 <222> (1)..(47)

40  
 <400> 44  
 tcgattaatt aattcgaacc cataataccc ataatagctg ttgcca 47

45  
 <210> 45  
 <211> 42  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: primer PCLIPMSF

50  
 <220>  
 <221> misc\_feature  
 <222> (1)..(42)

55



&lt;400&gt; 45

ccccaattgg tcgaccatca tcaataatat acettatttt gg

42

5

&lt;210&gt; 46

&lt;211&gt; 22

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

10

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: primer pCLIPBSRGI

15

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)..(22)

&lt;400&gt; 46

gcgaaaattg tcacttcctg tg

22

20

&lt;210&gt; 47

&lt;211&gt; 37

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

25

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Ecolinker+

30

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)..(37)

&lt;400&gt; 47

aattcggcgc gccgtcgacg atatcgatag cggccgc

37

35

&lt;210&gt; 48

&lt;211&gt; 37

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

40

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: Ecolinker-

45

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)..(37)

50

&lt;400&gt; 48

aattgcggcc gctatcgata tcgtcgacgg cgcgccg

37

55

5  
 <210> 49  
 <211> 49  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: linker HindXba+

10  
 <220>  
 <221> misc\_feature  
 <222> (1)..(49)

15  
 <400> 49  
 agctctagag gatccgtaa cgctagcgaa ttcaccggta ccaagctta 49

20  
 <210> 50  
 <211> 49  
 <212> DNA  
 <213> Artificial Sequence

25  
 <220>  
 <223> Description of Artificial Sequence: linker HindXba-

30  
 <220>  
 <221> misc\_feature  
 <222> (1)..(49)

35  
 <400> 50  
 ctagtaagct tggtagcggg gaattcgcta gcgttaacgg atcctctag 49

40  
 <210> 51  
 <211> 44  
 <212> DNA  
 <213> Artificial Sequence

45  
 <220>  
 <223> Description of Artificial Sequence: primer 35F1

50  
 <220>  
 <221> misc\_feature  
 <222> (1)..(44)

55  
 <400> 51  
 cggaattctt aattaatcga catcatcaat aatatacctt atag 44

<210> 52  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence

5  
 <220>  
 <223> Description of Artificial Sequence: primer 35R2

10  
 <220>  
 <221> misc\_feature  
 <222> (1)..(33)

15  
 <400> 52  
 ggtggtccta ggctgacacc tacgtaaaaa cag 33

20  
 <210> 53  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

25  
 <220>  
 <223> Description of Artificial Sequence: primer 35F3

30  
 <220>  
 <221> misc\_feature  
 <222> (1)..(30)

35  
 <400> 53  
 tggtggagat ctggtgagta ttgggaaaac 30

40  
 <210> 54  
 <211> 37  
 <212> DNA  
 <213> Artificial Sequence

45  
 <220>  
 <223> Description of Artificial Sequence: primer 35R4

50  
 <220>  
 <221> misc\_feature  
 <222> (1)..(37)

55  
 <400> 54  
 cggaattctt aattaaggga aatgcaaatc tgtgagg 37

55  
 <210> 55  
 <211> 34  
 <212> DNA  
 <213> Artificial Sequence

55  
 <220>  
 <223> Description of Artificial Sequence: primer 35F5

5  
 <220>  
 <221> misc\_feature  
 <222> (1)..(34)  
 <400> 55  
 cggaattcgc ggccgcggtg agtattggga aaac 34

10  
 <210> 56  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence

15  
 <220>  
 <223> Description of Artificial Sequence: primer 35R6

20  
 <220>  
 <221> misc\_feature  
 <222> (1)..(22)  
 <400> 56  
 cgccagatcg tctacagaac ag 22

25  
 <210> 57  
 <211> 23  
 <212> DNA  
 <213> Artificial Sequence

30  
 <220>  
 <223> Description of Artificial Sequence: primer 35F7

35  
 <220>  
 <221> misc\_feature  
 <222> (1)..(23)  
 <400> 57  
 gaatgctggc ttcagttgta atc 23

40  
 <210> 58  
 <211> 42  
 <212> DNA  
 <213> Artificial Sequence

45  
 <220>  
 <223> Description of Artificial Sequence: primer 35R8

50  
 <220>  
 <221> misc\_feature  
 <222> (1)..(42)



<400> 58

cggaattcgc ggccgcattt aaatcatcat caataatata cc

42

5

<210> 59

<211> 33

<212> DNA

<213> Artificial Sequence

10

<220>

<223> Description of Artificial Sequence: primer 35F11

15

<220>

<221> misc\_feature

<222> (1)..(33)

<400> 59

ggggtaccga attctcgcta gggtatttat acc

33

20

<210> 60

<211> 38

<212> DNA

<213> Artificial Sequence

25

<220>

<223> Description of Artificial Sequence: primer 35F10

30

<220>

<221> misc\_feature

<222> (1)..(38)

<400> 60

gctctagacc tgcaggttag tcagtttctt ctccactg

38

35

<210> 61

<211> 27

<212> DNA

<213> Artificial Sequence

40

<220>

<223> Description of Artificial Sequence: primer HBV-F

45

<220>

<221> misc\_feature

<222> (1)..(27)

<400> 61

ggctctagag atccttcgcg ggacgtc

27

50

55

5  
 <210> 62  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: primer HBV-R  
  
 10  
 <220>  
 <221> misc\_feature  
 <222> (1)..(26)  
  
 <400> 62  
 15 ggcgaattca ctgcctcca ccaagc 26  
  
 <210> 63  
 <211> 12  
 20 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence: oligonucleotide A-P1  
 25  
 <220>  
 <221> misc\_feature  
 <222> (1)..(12)  
  
 <400> 63  
 30 ctggtggtta at 12  
  
 <210> 64  
 <211> 14  
 35 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence:  
 40 oligonucleotide A-P2  
  
 <220>  
 <221> misc\_feature  
 <222> (1)..(14)  
 45  
 <400> 64  
 taaccaccag acgt 14  
  
 50  
 <210> 65  
 <211> 20  
 <212> DNA  
  
 55

5  
 <213> Artificial Sequence  
 <220>  
 <223> Description of Artificial Sequence: primer DF35-1  
 <220>  
 <221> misc\_feature  
 <222> (1)..(20)  
 10  
 <400> 65  
 cactcaccac ctccaattcc 20  
 15  
 <210> 66  
 <211> 32  
 <212> DNA  
 <213> Artificial Sequence  
 20  
 <220>  
 <223> Description of Artificial Sequence: primer DF35-2  
 <220>  
 <221> misc\_feature  
 <222> (1)..(32)  
 25  
 <400> 66  
 cgggatcccg tacgggtaga cagggttgaa gg 32  
 30  
 <210> 67  
 <211> 49  
 <212> DNA  
 <213> Artificial Sequence  
 35  
 <220>  
 <223> Description of Artificial Sequence: primer DF35-3  
 <220>  
 <221> misc\_feature  
 <222> (1)..(49)  
 40  
 <400> 67  
 cgggatccgc tagctgaaat aaagttaag tggttttatt taaaatcac 49  
 45  
 <210> 68  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence  
 50  
 <220>  
 <223> Description of Artificial Sequence: primer DF35-4  
 55

5  
 <220>  
 <221> misc\_feature  
 <222> (1)..(21)  
 <400> 68  
 ccagttgcat tgcttggtg g 21

10  
 <210> 69  
 <211> 27  
 <212> DNA  
 <213> Artificial Sequence

15  
 <220>  
 <223> Description of Artificial Sequence: degenerated primer

20  
 <220>  
 <221> misc\_feature  
 <222> (1)..(27)  
 <400> 69  
 cckgtstacc cgtacgaaga tgaaagc 27

25  
 <210> 70  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

30  
 <220>  
 <223> Description of Artificial Sequence: degenerated primer

35  
 <220>  
 <221> misc\_feature  
 <222> (1)..(30)  
 <400> 70  
 ccggctagct cagtcattt ctctgatata 30

40  
 <210> 71  
 <211> 31  
 <212> DNA  
 <213> Artificial Sequence

45  
 <220>  
 <223> Description of Artificial Sequence: primer 355ITR

50  
 <220>  
 <221> misc\_feature  
 <222> (1)..(31)

55



<400> 71

gatccggagc tcacaacgtc attttccac g

31

5

<210> 72

<211> 25

<212> DNA

<213> Artificial Sequence

10

<220>

<223> Description of Artificial Sequence: primer 353ITR

<220>

15

<221> misc\_feature

<222> (1)..(25)

<400> 72

cggaattcgc ggccgcattt aaatc

25

20

<210> 73

<211> 30

<212> DNA

<213> Artificial Sequence

25

<220>

<223> Description of Artificial Sequence: primer 35DE4

<220>

30

<221> misc\_feature

<222> (1)..(30)

<400> 73

ccaagcttg cttgtgtata tatattgtg

30

35

<210> 74

<211> 45

<212> DNA

<213> Artificial Sequence

40

<220>

<223> Description of Artificial Sequence: oligonucleotide TATA-plus

<220>

<221> misc\_feature

<222> (1)..(45)

45

<400> 74

agctttctta taaatttca gtgtagact agtaaattgc ttaag

45

50

55

5  
 <210> 75  
 <211> 45  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: oligonucleotide TATA-min

10  
 <220>  
 <221> misc\_feature  
 <222> (1)..(45)

15  
 <400> 75  
 agctcttaag caatttacta gtctaact gaaaatttat aagaa 45

20  
 <210> 76  
 <211> 37  
 <212> DNA  
 <213> Artificial Sequence

25  
 <220>  
 <223> Description of Artificial Sequence: primer tet3

30  
 <220>  
 <221> misc\_feature  
 <222> (1)..(37)

35  
 <400> 76  
 ccggagctcc atggcctaac tcgagtttac cactccc 37

40  
 <210> 77  
 <211> 30  
 <212> DNA  
 <213> Artificial Sequence

45  
 <220>  
 <223> Description of Artificial Sequence: primer tet5

50  
 <220>  
 <221> misc\_feature  
 <222> (1)..(30)

55  
 <400> 77  
 cccaagctta gctcgacttt cacttttctc 30

50  
 <210> 78  
 <211> 17  
 <212> DNA

5 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: oligo BB1

<220>

<221> misc\_feature

10 <222> (1)..(17)

<400> 78

gtgcctaggc cacgggg 17

15 <210> 79

<211> 13

<212> DNA

<213> Artificial Sequence

20 <220>

<223> Description of Artificial Sequence: oligo BB2

<220>

<221> misc\_feature

25 <222> (1)..(13)

<400> 79

gtggcctagg cac 13

30 <210> 80

<211> 20

<212> DNA

<213> Artificial Sequence

35 <220>

<223> Description of Artificial Sequence: primer 270F

<220>

<221> misc\_feature

40 <222> (1)..(20)

<400> 80

cacctctgcc taatcatctc 20

45 <210> 81

<211> 27

<212> DNA

<213> Artificial Sequence

50 <220>

<223> Description of Artificial Sequence: primer 270R

55

5  
 <220>  
 <221> misc\_feature  
 <222> (1)..(27)  
 <400> 81  
 gctctagaaa ttccactgcc ttccacc 27

10  
 <210> 82  
 <211> 35  
 <212> DNA  
 <213> Artificial Sequence

15  
 <220>  
 <223> Description of Artificial Sequence: primer DBPpcr1

20  
 <220>  
 <221> misc\_feature  
 <222> (1)..(35)  
 <400> 82  
 cgggatccgc caccatggcc agtcgggaag aggag 35

25  
 <210> 83  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence

30  
 <220>  
 <223> Description of Artificial Sequence: primer DBPpcr2

35  
 <220>  
 <221> misc\_feature  
 <222> (1)..(33)  
 <400> 83  
 cggaattctt aaaaatcaaa ggggttctgc cgc 33

40  
 <210> 84  
 <211> 34794  
 <212> DNA  
 <213> Adenoviridae

45  
 <220>  
 <221> misc\_feature  
 <222> (1)..(34794)  
 <223> /note="Nucleic acid sequence of Ad35"

50  
 <400> 84

55

catcatcaat aatatacctt atagatggaa tggcgccaat atgtaaatga ggtgatttta 60  
 aaaagtgtgg gccgtgtggt gattggctgt ggggttaacg gttaaaaggg gcggcgcggc 120  
 cgtgggaaaa tgacgtttta tgggggtgga gttttttgc aagtgtcgc gggaaatgtt 180  
 5 acgcataaaa aggcctcttt tctcacgga ctacttagtt tccccaggt atttaacagg 240  
 aaatgaggta gttttgaccg gatgcaagt aaaatgtctg attttcgcgc gaaaactgaa 300  
 tgaggaagtg tttttctgaa taatgtgta tttatggcag ggtggagtat ttgttcagg 360  
 ccaggtagac ttgacccat tacgtggagg ttctgattac cgtgttttt acctgaattt 420  
 ccgcgtaccg tgcataagtc ttctgtttt acgtagggtg cagctgatcg ctaggtatt 480  
 10 tatacctcag ggtttgtgc aagaggccac tcttgagtgc cagcgagaag agttttctcc 540  
 tctgcgcggc cagttaata ataaaaaat gagagatttg cgatttctgc ctcaggaaat 600  
 aatctctgct gagactggaa atgaaatatt ggagctgtg gtgcacgcc tgatgggaga 660  
 cgatccggag ccacctgtgc agcttttga gctcctacg cttcaggaac tgtatgatt 720  
 agaggtagag gcatcggagg attctaata ggaagctgtg aatggctttt ttaccgattc 780  
 15 tatgcttta gctgtaatg aaggattaga attagatccg ctttggaca cttcaatac 840  
 tccaggggtg attgtgaaa gcggtacagg tgtaagaaaa ttacctgatt tgagttccgt 900  
 ggactgtat ttgactgct atgaagacgg gtttctccg agtgatgagg aggaccatga 960  
 aaaggagcag tccatgcaga ctgcagcggg tgaggagtg aaggctcca atgttggtt 1020  
 tcagttgat tgcgggagc ttctggaca tggctgaag tctgtgaat ttacaggaa 1080  
 20 aaatactgga gtaaggaac tggtatgtc gtttgttat atgagaacgc actgccactt 1140  
 tattacagt aagtgtgtt aagtaaaat ttaaaggaat atgctgttt tcatatgat 1200  
 attgagtgtg agttttgtc ttctattat aggtctgtg tctgatgtg atgaatcac 1260  
 atctctgat tctactact cactctga tattcaagca cctgtctg tggacgtgc 1320  
 caagccatt cctgtgaagc ttaagcctgg gaaacgtcca gcagtggaga aacttgagg 1380  
 25 cttgttacag ggtggggagc gacctttga cttgagtaca cggaaacgtc caagacaata 1440  
 agtgttcat atcgtgtt acttaagggt acgtcaatat ttgtgtgaga gtgcaatga 1500  
 ataaaaat gtaactgtt cactggttt tattgcttt tggcgggga ctcaggata 1560  
 taagtagaag cagacctgt tggtagctc ataggagctg gcttcatcc atggaggtt 1620  
 gggccattt ggaagacct aggaagacta ggcaactgt agagagcgt tggacggag 1680  
 30 tctccggtt ttggagattc tggctgcta gtaattagc taggtagtt ttaggataa 1740  
 aacaggacta taaacaagaa ttgaaaagt tgttggtaga ttgccagga cttttgaag 1800  
 ctcttaattt gggccatcag gttcactta aaaaaaagt tttatcagt ttagacttt 1860  
 caacccagg tagaactgct gctgtgtg cttttctac tttatatta gataaatga 1920  
 tccgcagac tcatttcagc aggggatac ttttgattt catagccaca gcattgtga 1980  
 35 gaacatgga ggttcgcaag atgaggacaa tcttaggtta ctggccagt cagccttgg 2040  
 gtgtagcggg aatcctgagg catccaccg tcatgccagc gttctggag gaggaacagc 2100  
 aagaggacaa ccgagagcc gccctggacc ctccagtga ggaggcggag tagctgact 2160  
 gtctctgaa ctgcaacggg tgcctactg atctacgtc actggacggg atagggcgt 2220  
 taagaggag agggcatcca gtggtactg tctagatct gatttgctt taagttaat 2280  
 40 gactgcaga cgtctgaaa ccatttggtg gcatgaggt cagaaagagg gaaggatga 2340  
 agtttctga ttgcaggaga aatattcact ggaacagggt aaaacatgtt gttggagcc 2400  
 agaggatgat tggcggtg ccattaaaaa ttatgccaa atagctttga ggctgataa 2460  
 acagtataag atcagtagac ggattaatat ccggaatgct tttacatat ctggaatgg 2520  
 ggctgaggtg gtaatagata ctcaagacaa gacagttatt agatgtgca tgatggatat 2580  
 45 gtggcctgga gtagtcgga tggagcagt cacttttga aatgttaagt ttaggggaga 2640  
 tggttataat ggaatagtgt ttatggcaa taccaactt atattgcat gtgtagctt 2700  
 ttttggttc acaatacct gtgtagatc ctggggacag gtagtgtac ggggtgtg 2760  
 tttctatgc tgttgattg ccacagctg cagaaccaag agtcaattgt ctctgaagaa 2820  
 atcatattc caaagatga acctggcat tctgaatgaa ggcaagcaa ggtccgtca 2880  
 50 ctgccttct acagatactg gatgtttat ttaattaag ggaaatgcca gcgtaagca 2940  
 taacatgatt tgtgtgctt ccgatgagag gccttatcaa atgctcact gtgctggtg 3000  
 gcattgtaat atgctggcta ctgtcatat ttttccat caacgcaaaa aatggcctgt 3060  
 tttgatcac aatgtgtga ccaagtgcac catgcatga ggtggcgta gaggaatgtt 3120



tatgccttac cagtgtaca tgaatcatgt gaaagtgttg ttggaaccag atgccttttc 3180  
 cagaatgagc ctaacaggaa tctttgacat gaacacgcaa atctggaaga tcttgaggta 3240  
 tgatgatacg agatcgaggg tgcgcgcatg cgaatgcgga ggcaagcatg ccagggtcca 3300  
 gccggtgtgt gtagatgtga ccgaagatct cagaccggat catttggtta ttgcccgcac 3360  
 tggagcagag ttccgatcca gtggagaaga aactgactaa ggtgagtatt gggaaaactt 3420  
 tggggtggga tttcagatg gacagattga gtaaaaattt gtttttctg tcttcagct 3480  
 gacatgagtg gaaatgcttc ttttaagggg ggagtcttca gcccttatct gacagggcgt 3540  
 ctccatcctt gggcaggagt tcttcagaat gttatgggat ctactgtgga tggagaccc 3600  
 gttcaaccg ccaattcttc aacgtgacc tatgtactt taagtcttc accttggac 3660  
 gcagctgcag ccgtgcgc gcctctgtc gccgtaaca ctgtgcttg aatgggttac 3720  
 tatggaagca tctgggctaa ttccacttc tctaataacc ctctacact gactcaggac 3780  
 aagtacttg tcttttggc ccagctggag gcttgaccc aacgtctgg tgaacttct 3840  
 cagcaggtgg ccgagtggc agtacaact gagtctgtc tggcagcgc aaagtctaaa 3900  
 taaaaaaaaat tccagaatca atgaataaat aaacgagctt gttgttgatt taaatcaag 3960  
 tgttttatt tcaatttctg cgcacggtat gccctggacc accgatctc atcattgaga 4020  
 actcgggtgga tttttccag aatctatag aggtgggatt gaatgttag atacatgggc 4080  
 attaggccgt ctttgggtg gagatagctc cattgaaggg attcatgctc cgggtagtg 4140  
 ttgtaataca ccagtcata acaaggctgc agtgcaggt gttgcacaat atcttttaga 4200  
 agtaggtga ttgccacaga taagcccttg gttaggtgt ttacaacccg gttgagctgg 4260  
 gaggggtgca ttccaggtga aattatgtc attttggtt ggattttta gttggcaata 4320  
 ttgcccaa gatccgtct tgggtcatg ttatgaagga ctaccaagac ggtgtatccg 4380  
 gtacatttag gaaattatc gtgcagctg gatggaaaag cgtggaaaaa ttggagaca 4440  
 ccctgtgtc ctccgagatt ttccatgac tcatcatga taatagcaat ggggcgtg 4500  
 gcagcggcgc gggcaaacac gttccgtgg tctgacacat catagttatg ttctgagtt 4560  
 aatcatcat aagccattt aatgaattg gggcggagcg taccagattg gggtatgaat 4620  
 gttccttcgg gcccgagc atagttccc tcacagattt gcatttcca agcttccagt 4680  
 tctgaggtg gaatcatgc cacctgggg gctatgaaga acaccgttc ggggcgggg 4740  
 gtgattagtt gggatgatag caagtctg agcaattgag atttccaca tccggtggg 4800  
 ccataaataa ttccgattac aggttgcagg tggtagtta gggaacggca actgccgtc 4860  
 tctgaagca agggggccac ctggttcatc atttccctta catgatatt ttccgcacc 4920  
 aatccatta ggagcgctc tctcctagt gatagaagt cttgtagtga ggaaaagtt 4980  
 ttccagcgtt ttgaccgtc agcatggg atttggaaa gatttctg caaaagttc 5040  
 agtctgtcc acagttcagt gatgtgtct atggcatctc gatccagcag acctcctct 5100  
 ttccgggtt tggacggctc ctggagtagg gtatgagacg atggcgctc agcgtgcca 5160  
 ggggtcgtc ctccaggtt ctgagtctc gactcagggt tgttccgtc acagtgaagg 5220  
 ggtgtgcgc tcttggggc cttgccagg tgccttcag actcattctg ctggtggaga 5280  
 acttctgtc cttggcgccc tgtatgtcg ccaagtagca gttaccatg agttcgtagt 5340  
 tgagcgtc ggctgcgtg ctttggcgc ggagcttacc ttggaagt ttcttcata 5400  
 ccgggcagta taggcattc agcgcatata gcttggcgc aaggaaaatg gattctggg 5460  
 agtatgcat cgcgcgcag gaggcgcaa cagtttaca ttccaccagc caggttaaat 5520  
 ccggttcatt ggggtcaaaa acaagtttc cccatattt ttgatgcgt ttcttacct 5580  
 tggctccat aagttcgtt cctcgtttag tgacaaacag gctgtccgta tctccgtaga 5640  
 ctgattttac aggcctctc tccagtggag tgcctcgtc ttcttcgtac aggaactct 5700  
 accactctga tacaaggcg cgcgtccagg ccagcaciaa ggaggtatg tggaggggt 5760  
 agcgtcgtt gtcaaccagg ggtccacct ttccaaagt atgcaaacac atgtaccct 5820  
 cttaacatc caggaatgtg attggttgt aggtgtatt cacgtgacct ggggtcccc 5880  
 ctgggggggt ataaaaggg gcggttctt gctcttctc actgtcttc ggtcgtgt 5940  
 ccaggaaagt cagctgttg gtaggtatt cctctcgaa ggcggcatg acctctgcac 6000  
 tcaggtgtc agtttctaag aacgaggagg atttgaatt gacagtccg gttgagatgc 6060  
 cttcatgag gtttctec atttgcag aaaacacaat tttttattg tcaagtttg 6120  
 tggcaaatga tccatacag gcgttgata aaagtttgc aatggatgc atggttgg 6180  
 tctttcctt gtccgcgc tcttggcg cgatgtgag ttggacatac tgcgtgcca 6240

5 ggcacttcca ttcggggaag atagttgtta attcatctgg cacgattctc acttgccacc 6300  
 ctcgattatg caaggtaatt aaatccacac tgggtggccac ctgcctcga aggggttcat 6360  
 tggccaaca gagcctacct ctttcttag aacagaaagg gggaaagtgg tctagcataa 6420  
 gttcatcggg agggctcga tccatggtta agattcccgg aagtaaatec ttatcaaat 6480  
 agctgatggg agtgggtgca tctaaggcca ttgccattc tcgagctgcc agtgcgcgt 6540  
 catatgggtt aaggggactg cccagggca tgggatgggt gagagcagag gcatacatgc 6600  
 cacagatgac atagacgtag atgggatcct caaagatgcc tatgtagggt ggatagcatc 6660  
 gccccctct gatacttct cgcacatagt catatagttc atgtgatggc gctagcagcc 6720  
 10 cgggaccaa gttgggtgca ttgggtttt ctgtctgta gacgatctgg cgaaagatgg 6780  
 cgtgagaatt ggaagagatg gtgggtcttt gaaaaatgtt gaaatgggca tgaggtagac 6840  
 ctacagagtc tctgacaaag tgggcataag attcttgaag ctgtgttacc agttcggcgg 6900  
 tgacaagtac gtctagggcg cagtagtcaa gtgtttcttg aatgatgtca taacctggtt 6960  
 ggttttctt tcccacagt tcgcggtga gaaggtattc ttgcgatcc ttccagtact 7020  
 15 cttctagcgg aaaccgtct ttgtctgac ggtaagatcc tagcatgtag aactgattaa 7080  
 ctgccttga agggcagcag ccttctcta cgggtagaga gtatgctga gcagctttc 7140  
 gtagcgaagc gtgagtaagg gcaaagggtt ctctgacat gactttgaga aattggtatt 7200  
 tgaagtccat gtcgtcacag gctccctgtt cccagagttg gaagtctacc cgtttctgt 7260  
 aggcggggtt gggcaagcg aaagtaacat cattgaagag aatcttaccg gctctgggca 7320  
 20 taaaattgcg agtgatggcg aaaggctgtg gtactccgc tcgattgttg atcacctggg 7380  
 cagctaggac gatttcgtcg aaaccgtga tgtgtgtcc tacgatgtat aattctatga 7440  
 aacgcggcgt gcctctgacg tgaggtagct tactgagctc atcaaagggt aggtctgtgg 7500  
 ggtcagataa ggctgagtgt tcgagagccc attcgtgcag gtgaggattt gcatgtagga 7560  
 atgatgacca aagatctacc gccagtctg tttgtaactg gtcccgatac tgacgaaat 7620  
 25 gccggcaat tgccatttt tctggagtga cacagtagaa ggttctgggg tcttgttgc 7680  
 atcgatcca cttgagtta atggctagat cgtgggcat gttgacgaga cgctctctc 7740  
 ctgagagttt catgaccagc atgaaaggaa ctagtgttt gccaaaggat cccatccagg 7800  
 tgtaagttc cacatcgtag gtcaggaaga gtcttctgt gcgaggatga gagccgatcg 7860  
 ggaagaactg gatttcctgc caccagttgg aggatggct gttgatgtga tggaagtaga 7920  
 30 agttctgcg gcgcgcgag cattcgtgtt tgtcttga cagacggcg cagtagtgc 7980  
 agcgttgac ggggtgtatc tctgaatga gctgtacct gcttccctg acgagaaatt 8040  
 tcagtgggaa gccagggcct ggcgattgta tctcgtctc ttctatattc gctgtatcg 8100  
 cctgtctac tctgtttcg atgggtgca tctgacgag ccccgcggg aggcaagtcc 8160  
 agacctggc gcgggagggg cggagctgaa ggacgagagc gcgcaggctg gagctgtcca 8220  
 35 gagtctgag acgtcggga ctgaggttag taggtaggga cagaagatta acttgcatga 8280  
 tctttccag ggcgtcggg aggttcagat ggtactgtat ttccacaggt tcgtttgtag 8340  
 agacgtcaat ggcttcagg gttccgtgtc cttggggc cactaccgta ctttgtttt 8400  
 ttctttgat cgtgtgtgc tctctgtct ctgcatgct cagaagcgt gacggggacg 8460  
 cgcgcgggc ggcagcgtt gttccggacc cgggggcatg gctggtagt gcacgtcggc 8520  
 40 gccgcgacg ggcaggttct ggtattgcg tctgagaaga cttgcgtcg ccaccacgcg 8580  
 tcgattgacg tctgtatct gacgtctct ggtgaaagct accggcccc tgagcttgaa 8640  
 cctgaaagag agttcaacag aatcaatttc ggtatcgtta acggcagctt gtctcagtat 8700  
 ttctgtacg tcaccagagt tctctggta ggcgatctcc gccatgaact gctcgatttc 8760  
 ttctctga agatctcgc gaccgctct ttcgacgtg gccgcaggt cattggagat 8820  
 45 accggccatg agttgggaga atgattcat gccgcctcg ttccagacgc ggctgtaaac 8880  
 cacggcccc tcggagtctc ttgcgcgat caccacctga gcgaggttaa gctccacgtg 8940  
 tctggtgaag accgcatagt tgcattggcg ctgaaaaagg tagttgagt tggtggcaat 9000  
 gtgttcggcg acgaagaaat acatgatcca tctctcagc ggcatttcgc taacatcgcc 9060  
 cagagcttc aagectcca tggcctcgt gaagtccac gcaaaattaa aaaactggga 9120  
 50 gtttcgcgcg gacacgtca attctctc gagaagacgg atgagttcgg ctatggtggc 9180  
 ccgtacttc gttcgaagg ctcccggat ctctcttc tctctatct cttctccac 9240  
 taacatctct tctctctt caggcgggg cggaggggg acgcggcgac gtcgacggcg 9300  
 cacgggcaaa cggctgatga atcgttcaat gacctctcg cggcgcggc gcatggttc 9360

55

agtgacggcg cggccgttct cgcgcggtcg cagagtaaaa acaccgccgc gcattctctt 9420  
 aaagtgggta ctgggagggt ctccgtttgg gagggagagg gcgtgatta tacattttat 9480  
 taattggccc gtagggactg cgcgcagaga tctgacgtg tcaagatcca cgggatctga 9540  
 aaacctttcg acgaaagcgt ctaaccagtc acagtcacaa ggtaggctga gtacggcttc 9600  
 ttgtggcgcg ggggtggtat gtgttcggtc tgggtcttct gtttcttctt catctcggga 9660  
 aggtgagacg atgctgctgg tgatgaaatt aaagtaggca gttctaagac ggcggtatgt 9720  
 ggcgaggagc accagggtctt tgggtccggc ttgctggata cgcaggcgat tggccattcc 9780  
 ccaagcatta tctgacatc tagcaagatc tttgtagtag tcttgcata gccgttctac 9840  
 gggcacttct tctcaccg tcttccatg catacgtgtg agtccaaatc cgcgattgg 9900  
 ttgtaccagt gccaaagtcag ctacgactct ttcggcgagg atggcttgc gtacttgggt 9960  
 aagggtgggt tgaaagtcac caaaatccac aaagcgggtg taagcccctg tattaatgt 10020  
 gtaagcacag ttggccatga ctgaccagtt aactgtctgg tgaccagggc gcacagctc 10080  
 ggtgtattta aggcgcgaat aggcgcgggt gtcaaagatg taatcgttc aggtgcgcac 10140  
 cagatactgg tacctataa gaaaatgcgg cgggtggtgg cggtagagag gccatcgttc 10200  
 ttagctgga gcgccagggg cgaggtcttc caacataagg cggtgatagc cgtagatgta 10260  
 cctggacatc cagggtgatt ctgcggcggt agtagaagcc cgaggaaact cgcgtacgcg 10320  
 gttccaaatg ttgcgtagcg gcataagta gttcattgta ggcacggtt gaccagtgcg 10380  
 gcgcgcgcag tcattgatgc tctatagaca cggagaaaat gaaagcgttc agcgactga 10440  
 ctccgtagcc tggaggaacg tgaacgggtt gggtcgcggt gtaccccggt tcgagactg 10500  
 tactcgagcc ggccggagcc gcggctaacg tggattggc actcccgctc cgaccagcc 10560  
 tacaanaatc caggatagcg aatcgagtcg tttgctggt ttcggaatgg cagggaagt 10620  
 agtctattt tttttttt tttgcgctc agatgatcc cgtctgcga cagatgcgc 10680  
 cccaacaaca gccccctc cagcagcagc agcagcaacc aaaaaggct gtcctgcaa 10740  
 ctactgcaac tgcgcgctg agcgggtgcg gacagccgc ctatgatctg gacttgaag 10800  
 agggcgaagg actggcacgt ctagggtgcg cttcgccga gcggcatccg cgagtcaac 10860  
 tgaanaaga ttctcgcgag gcgtatgtc ccaacagaa cctatttaga gacagaagcg 10920  
 gcgaggagcc ggaggagatg cgagcttcc gcttaacgc gggctgtgag ctgcgtcacg 10980  
 gtttgaccg aagacgagtg ttgcgagacg aggtttcga agttgatgaa gtgacagga 11040  
 tcagtctgc cagggcacac gtggctgcg ccaacctgt atcggttac gagcagacag 11100  
 taaaggaaga gcgtaactc caaaagtctt ttaataatca tgtgcgaacc ctgattgcc 11160  
 gcgaagaagt taccttgggt ttgatgcatt tgtgggatt gatggaagct atcattcaga 11220  
 acctactag caaacctctg accgccagc tgttctggt ggtgcaacac agcagagaca 11280  
 atgaggctt cagagaggcg ctgctgaaca tcaccgaacc cgaggggaga tggttgatg 11340  
 atcttatcaa cattctacag agtatcatg tgcaggagcg gacgtgggc ctggccgaga 11400  
 aggtagctgc catcaattac tcggtttga gcttgggaaa atattacgt cgcaaatct 11460  
 acaagactcc atacgttccc atagacaagg aggtgaagat agatgggttc tacatgcga 11520  
 tgacgtcaa ggtcttgacc ctgagcgatg atcttgggt gtatcgcaat gacagaatgc 11580  
 atcgcgcggt tagcgcagc aggaggcgcg agttaagcga cagggaactg atgcacagt 11640  
 tgcaagagc tctgactgga gctggaaccg agggtagaa ttacttcgac atgggagctg 11700  
 acttgagtg gcagcctagt cgcagggtc tgagcgccgc gacggcagga tgtgagcttc 11760  
 cttacataga agaggcggt gaaggcgagg aggaagagg cgagtactg gaagactgat 11820  
 ggcacaacc gtgtttttg ctatggaag cagcaagcac cggatccgc aatgcggcg 11880  
 gcgtgcaga gccagcgtc cggcattaac tctcgagc attggacca ggccatgaa 11940  
 cgtatcatg cgttgacgac tcgcaacccc gaagcctta gacagcaacc ccaggccaac 12000  
 cgtctatcg ccatcatgga agctgtatg cttcccgat ctaatccac tcatgagaag 12060  
 gtcttgcca tctgaacgc gttggtggag acaaaagta ttcgtccaga tgaggccgga 12120  
 ctggtatata acgtctctt agaacgcgtg gctcgtaca acagtagcaa tgtgcaaac 12180  
 aatttgacc gtatgataac agatgtacgc gaagcgtgt ctgagcgca aaggttcag 12240  
 cgtgatgcca acctgggttc gctggtggcg ttaaatgctt tcttgagtac tcagcctgt 12300  
 aatgtgccg gtgtcaaca ggattatact aacttttaa gtgcttgag actgatgta 12360  
 tcagaagtac ctgagcgga agtgtatcag tccgtctg attactctt tcagactagc 12420  
 agacagggt tgcagacgt aaatctgagc caagcttta aaacctaa aggtttgtg 12480



ggagtgcag ccccggtagg agaaagagca accgtgtcta gcttgtaac tccgaactcc 12540  
 cgctgttat tactgttgg agtcctttc accgacagcg gtagcatga ccgtaattcc 12600  
 tatttgggtt acctactaaa cctgtatcgc gaagccatag ggcaaagtca ggtggacgag 12660  
 5 cagacctatc aagaaattac ccaagtcagt cgcgcttgg gacaggaaga cactggcagt 12720  
 ttggaagcca ctctgaactt ctgtctacc aatcggctc aaaagatccc tctcaatat 12780  
 gctcttactg cggaggagga gaggatcctt agatatgtc agcagagcgt gggattgtt 12840  
 ctgatgcaag agggggcaac tccgactgca gcactggaca tgacagcgcg aaatatggag 12900  
 cccagcatgt atgccagtaa ccgaccttc attacaaac tgctggacta ctgcacaga 12960  
 10 gctccgcta tgaactctga ttatttcacc aatgccatc taaaccgca ctggctgcc 13020  
 ccacctggtt tctacacggg cgaatatgac atgccgacc ctaatgacgg atttctgtg 13080  
 gacgacgtg acagegatgt ttttcacct cttctgac atcgacgtg gaaaaaggaa 13140  
 ggcggtgata gaatgcattc ttctgcatc ctgtccggg tcatgggtg taccgcggt 13200  
 gagcccgagt ctgcaagtcc tttcctagt ctacctttt ctctacacag tgtacgtagc 13260  
 15 agcgaagtgg gtagaataag tgcocgagt ttaatggcg aagaggagta cctaaacgat 13320  
 tcttgctca gaccggcaag agaaaaaat tcccaaca atggaataga aagtttggg 13380  
 gataaaatga gtagatggaa gacttatgt caggatcaca gagacgagcc tgggatcatg 13440  
 gggactacaa gtagagcgag ccgtagacgc cagcgccatg acagacagag ggtcttgtg 13500  
 tgggacgatg aggattcggc cgatgatagc agcgtgttg acttgggtg gagaggaagg 13560  
 20 ggcaaccgtt ttgtcattt gcgcctcgc ttgggtgta tgtgtgaaa aaaaataaaa 13620  
 aagaaaaact caccaaggcc atggcgacga gcgtacgtc gttctctt attatctgtg 13680  
 tctagtataa tgaggcgagt cgtgctaggc ggagcgggtg tgtatccgga gggctcctt 13740  
 ccttcgtacg agagcgtgat gcagcagcag caggcgacgg cggatgatga atccccactg 13800  
 gaggtccct ttgtccctc gcgatactg gcacctacg agggcagaaa cagcattcgt 13860  
 25 tactcggaac tggcaccica gtacgatacc accaggtgt atctggtgga caacaagtcg 13920  
 gggacattg cttctctgaa ctatcagaat gaccacagca acttctgac caggtgtg 13980  
 cagaacaatg actttacccc tacggaagcc agcaccaga ccattaactt tgatgaacga 14040  
 tgcggtggg gcggtcagct aaagaccatc atgcatacta acatgccaaa cgtgaacgag 14100  
 tatatgtta gtaacaagt caaagcgcgt gtgatggtt ccagaaaacc tccgacggt 14160  
 30 gctgcagtg gggatactta tgatcacaag caggatattt tggaatatga gtggttcgag 14220  
 ttactttgc cagaaggcaa ctttcagtt actatgacta ttgattgat gaacaatgcc 14280  
 atcatagata attacttga agtgggtaga cagaatggag tgcttgaag tgacattgt 14340  
 gtaagttc acaccaggaa cttcaagcgt ggtgggac ccgaaacca gttgatcatg 14400  
 cctggagtgt atacgtatga agccttccat cctgacattg tcttactgcc tggctcgga 14460  
 35 gtggatttta ccgagagtcg tttagcaac cttcttgta tcagaaaaa acagccattt 14520  
 caagagggtt ttaagattt gtatgaagat ttagaagggt gtaatatcc gccctctt 14580  
 gatgtagatg cctatgagaa cagtaagaaa gaacaaaag ccaaaataga agctgctaca 14640  
 gctgctgcag aagetaagc aaacatagtt gccagcgact ctacaagggt tgctaacgt 14700  
 ggagaggta gagagacaa tttgcgcca acacctgtc cgactgcaga atcattattg 14760  
 40 gccgatgtt ctgaaggaa ggacgtgaaa ctactattc aacctgtaga aaaagatagt 14820  
 aagaatagaa gctataatgt gttggaagac aaatcaaca cagcctatc cagttggtat 14880  
 cttcgtaca attatggcga tccgaaaaa ggagtgcgt cctggacatt gctaccacc 14940  
 tcagatgta cctgcggagc agagcaggtt tactggctc tccagacat gatgaaggat 15000  
 cctgtcactt tccgtccac tagacaagtc agtaactacc ctgtggtggg tgcagagctt 15060  
 45 atgcccgtt tctcaaagag cttctacaac gaacaagctg tgtactcca gcagctccg 15120  
 cagtcacct cgttacgca cgtctcaac cgtttctg agaaccagat ttaatccgt 15180  
 ccgcccgc ccaccattac caccgtcagt gaaaacgtc ctgctctac agatcacggg 15240  
 accctgccgt tgcgacgag tatccggga gtccaacgtg tgaccgttac tgacccaga 15300  
 cgcgcacct gtcctacgt gtacaaggca ctgggcatag tgcaccgcg cgtccttca 15360  
 50 agccgcacti tctaaaaaaa aaaaatgtc attcttatc cggcagtaa taacaccgt 15420  
 tgggtctgc gcgtccaag caagatgtac ggaggcgac gcaaacgtc taccaacat 15480  
 cccgtgcgt ttcgcgaca tttcgcgt ccatgggtg cctcaagg cgcactgc 15540  
 gttcgaacca ccgtgatga tgtaatcgt caggtggtg ccgacgccg taattatact 15600

cctactgccc ctacatctac tgtggatgca gttattgaca gtgtagtggc tgacgctgc 15660  
 aactatgctc gacgtaagag ccggcgaagg cgcattgccca gacgccaccg agctaccact 15720  
 gccatgcgag ccgcaagagc tctgctacga agagctagac gcgtggggcg aagagccatg 15780  
 5 cttagggcgg ccagacgtgc agcttcgggc gccagcgccg gcaggtcccg caggcaagca 15840  
 gccgctgtcg cagcggcgac tattgccgac atggcccaat cgcgaagagg caatgtatac 15900  
 tgggtgcgtg acgctgccac cggtaacgt gtaccggtgc gcacccgtcc cctcgcact 15960  
 tagaagatac tgagcagtct ccgatgttgt gtcccagcgg cgaggatgtc caagcgcaaa 16020  
 tacaaggaag aaatgctgca ggtatcgc cctgaagtct acggccaacc gttgaaggat 16080  
 10 gaaaaaaaac cccgcaaaat caagcgggtt aaaaaggaca aaaaagaaga ggaagatggc 16140  
 gatgatgggc tggcggagtt tgtgcgcgag ttgccccac ggcgacgcgt gcaatggcgt 16200  
 gggcgcaag ttcgacatgt gttgagacct ggaacttcgg tggctttac acccggcgag 16260  
 cgttcaagcg ctactttta gcgttcctat gatgaggtgt acggggatga tgatattctt 16320  
 gagcaggcgg ctgaccgatt aggcgagttt gcttatggca agcgtagtag aataacttcc 16380  
 15 aaggatgaga cagtgtcaat acccttggat catggaaatc ccaccctag tcttaaacg 16440  
 gtcactttgc agcaagtgtt acccgtaact ccgcgaacag gtgttaaacg cgaaggatga 16500  
 gatttgatc ccactatgca actgatgta cccaaacgcc agaagtggga ggacgtttg 16560  
 gagaaagtaa aagtggatcc agatattcaa cctgaggtta aagtgagacc cattaagcag 16620  
 gtagcgctg gtctgggggt acaaactgta gacattaaga tccactga aagtatggaa 16680  
 20 gtgcaactg aaccgcaaa gcctactgcc acctccactg aagtcaaac ggatccatgg 16740  
 atgccatgc ctattacaac tgacgcgcc ggtccactc gaagatccc acgaaagtac 16800  
 ggtccagcaa gtctgtgat gcccaattat gttgtacacc catctattat tctactcct 16860  
 ggttaccgag gcactcgtc ctatcgcagc cgaacagta cctcccgcc tgcgcgaag 16920  
 acacctgcaa atgcagtcg tgcgcgtaga cgcacaagca aaccgactcc cggcgccctg 16980  
 25 gtgcggcaag tgtaccgcaa tggtagtgc gaaccttga cactgcgcg tgcgcgttac 17040  
 catccgagta tcatcactta atcaatgtt cgcgtccct cttgcagata tggccctac 17100  
 ttgtgcctt cgcgttccca tcactggta ccgaggaaga aactgcgcc gtagaagagg 17160  
 gatgttggga cgcggaatgc gacgtacag gcgacggcgt gctatccga agcaattgcg 17220  
 ggggtgttt ttaccagct taattccat tategtgt gcaattggcg cgataccagg 17280  
 30 catagcttc gtggcggtc aggcctcgca acgacattga cattggaaaa aaacgtata 17340  
 aataaaaaaa aatacaatgg actctgacac tctgttct gtgactatgt tttcttagag 17400  
 atggaagaca tcaattttc atccttggct ccgcgacag gcacgaagcc gtacatgggc 17460  
 acctggagcg acatcggcac gagccaactg aacggggggc cttcaattg gagcagtatc 17520  
 tggagcgggc ttaaaaattt tggctcaacc ataaaaacat acgggaacaa agcttggaa 17580  
 35 agcagtacag gacagcgct tagaaataaa cttaaagacc agaactcca acaaaaagta 17640  
 gtcgatggga tagcttcgg catcaatgga tggtagatt tggtaacca ggtctgcag 17700  
 aaaaagataa acagtcgtt ggaccgccc ccagcaaccc caggtgaaat gcaagtggag 17760  
 gaagaaattc ctccgcaga aaaacgagc gacaagcgc cgcgtcccga ttggaagag 17820  
 acgttgtga cgcgcgtaga tgaaccgct tcttatgagg aagcaacgaa gcttggaaatg 17880  
 40 cccaccata gaccgatag ccaatggcc accgggtga tgaaccttc tcagttgcat 17940  
 cgacccgtca ccttggattt gccctccc cctgtgcta ctgtgtacc cgcttctaag 18000  
 cctgtcgtg cccgaaacc agtcgcgta gccaggtcac gtccggggg cgctcctcgt 18060  
 ccaaatgcgc actggcaaaa tactctgaac agcatcgtg gtctaggcgt gcaaagtga 18120  
 aaacgcgct gctgcttta attaatatg gagtagcgt taacttgcct atctgtgtat 18180  
 45 atgtgtcatt acagcgcgt acagcagcag aggaaaaaag gaagaggtcg tgcgtcgac 18240  
 ctgagttact ttcaagatgg ccaccctac gatgtgccc caatgggcat acatgcacat 18300  
 cgccggacag gatgcttcgg agtacctgag tccgggtctg gtgcagttcg cccgcgccac 18360  
 agacacctac ttcaatctgg gaaataagt tagaaatccc accgtagcgc cgaccacga 18420  
 tgtgaccacc gaccgtagc agcggctcat gttgcgttc gtgcccgtt accgggagga 18480  
 50 caatacatc tcttacaag tgcgtacac cctggccgtg ggcgacaaca gagtgtgga 18540  
 tatggcagc agttctttg acattagggg cgtgttgac agaggtcca gttcaaac 18600  
 ctattctgtt acgcttaca actctctgc tctaaaggc gtcctaatg catctaatg 18660  
 gattgcaaaa ggcgtaccaa ctgcagcagc cgcaggcaat ggtgaagaag aacatgaac 18720



agaggagaaa actgctactt acacttttgc caatgctcct gtaaaagccg aggctcaaat 18780  
 tacaaaagag ggcttaccac taggtttgga gatttcagct gaaaacgaat ctaaaccat 18840  
 ctatgcagat aaactttatc agccagaacc tcaagtggga gatgaaactt ggactgacct 18900  
 5 agacggaaaa accgaagagt atggaggcag ggctctaaag cctactacta acatgaaacc 18960  
 ctgttacggg tctatgcga agcctactaa tttaaaaggt ggtcaggcaa aaccgaaaaa 19020  
 ctcggaaccg tcgagtgaac aaattgaata tgatattgac atggaatitt ttgataactc 19080  
 atcgcaaga acaaaacttca gtctaaaaat tgtcatgtat gcagaaaatg taggtttgga 19140  
 aacgccagac actcatgtag tgtacaaacc tggacagaa gacacaagtt ccgaagctaa 19200  
 10 ttgggacaa cagtctatgc ccaacagacc caactacatt ggcttcagag ataactttat 19260  
 tggactcatg tactataaca gtactggtaa catgggggtg ctggctggtc aagcgtctca 19320  
 gttaatgca gtggtgact tgcaggacag aaacacagaa cttcttacc aactcttct 19380  
 tgactctctg ggagacagaa ccagatactt tagcatgttg aatcaggctg tggacagtta 19440  
 tgatctgat gtacgtgta ttgaaaatca tgggtggaa gatgaacttc ccaactattg 19500  
 15 tttccactg gacggcatag gtgtccaac aaccagttac aaatcaatag ttccaatgg 19560  
 agaagataat aataattgga aagaacctga agtaaatgga acaagtgaac tcggacaggg 19620  
 taatttgtt gccatggaaa ttaacctca agccaatca tggcgaagtt tctttatc 19680  
 caatgtggt ctgtatctc cagactcgt ccaatacacc ccgtccaatg tcaactctcc 19740  
 agaaaacaaa aacacctacg actacatgaa cgggagggtg gtgcgccat ctctagtaga 19800  
 20 cacctatgtg aacattgggt ccagggtggt tctggatgcc atggacaatg tcaaccatt 19860  
 caaccaccac cgtaacgtg gcttgctta ccgatctatg cttctgggtg acggacgtta 19920  
 tgtgcttcc cactataag tgcctcaaaa attctctgct gttaaaaacc tctgcttct 19980  
 ccaggctcc taccattatg agtggaaact taggaaggat gtgaacatgg ttctacagag 20040  
 ttccctcgt aacgacctgc gggtagatgg ccagacatc agtttcacga gcatcaacct 20100  
 25 ctatgctact ttttccca tggctcaaa caccgttcc accctgaag ccatgctgcg 20160  
 gaatgacacc aatgatcagt cattcaaga ctacctatc gcagtaaca tgcctaccc 20220  
 cattctgcc aatgcaacca atattccat ttccattct tctcgcaact gggcggtt 20280  
 cagaggctgg tcattacca gactgaaac caaagaaact cctctttgg ggtctggatt 20340  
 tgacctac ttgtctatt ctggtctat tccctacgt gatggtacct tctacctga 20400  
 30 ccacacttt aagaagggtt ccatcatgtt tgactctca gtgagctggc ctggaaatga 20460  
 caggttacta tctctaacg aattgaaat aaagcgcact gtggatggcg aaggctaca 20520  
 cgtagccaa tgcaacatga ccaagactg gttcttggt cagatgctc ccaactaca 20580  
 catcgctat cagggttct acattccaga aggatacaa gatcgcatgt attcatttt 20640  
 cagaaactc cagccatga gcaggcaggt ggtgatgag gtcaattaca aagactcaa 20700  
 35 ggccgtgcc atacctacc aacacaaca ccttggtt gtgggttaca tggctccgac 20760  
 catgcgcaa gtcaacct atccgctaa ctatccat cactcattg gaacaactgc 20820  
 cgtaaatagt gttacgaga aaaagttct gtgtgacaga accatgtgc gcataccgt 20880  
 ctgagcaac tcatgtcta tgggggacct tacagactg ggacagaata tgcctatgc 20940  
 caactcagct catgctctg acatgacct tgaggtgat ccatggatg agccacct 21000  
 40 gctttatct ctctcgaag tttcgact ggtcagagt catcagccac acccggtat 21060  
 catcaggea gtctacctgc gtacacct ctggccgt aacgtacca cgtagaagc 21120  
 ttctgttc ttgaaatag cagctgcaac catggctgc ggtccaaa accgctccag 21180  
 cgagcaagag ctgagacca ttgtcaaga cctgggtg ggacctatt tttgggaac 21240  
 ctacgataag cgcttccgg gttcatgg cccgataag ctgcctgtg ccattgtaa 21300  
 45 tacggccga cgtgagacgg ggggagagca ctggttgg ttgggtgga acccagttc 21360  
 taacacctgc taccctttg atcttttg attctcggat gatcgtctca aacagattta 21420  
 ccagttgaa tatgagggtc tctgcgag cagcgtctt gctaccaagg accgtgtat 21480  
 tacgtggaa aaatctacc agaccgtga gggcccggt tctgcgct gcggacttt 21540  
 ctgctcatg ttcttcacg cctttgtga ctggctgac cgtccatgg acggaaacc 21600  
 50 caccatgaa ttgtaactg gattgcaaa caacatgct cattctcta agtccagcc 21660  
 caccctgtg gacaatcaa aagactcta ccttttct aataccatt cgcctattt 21720  
 tgcctcat cgtacacaca tcgaaaggc cactgcgtc gaccgatgg atgtcaata 21780  
 atgactcat taaacaacgt gttcaataa catcattta ttttttaca tgtatcaag 21840

ctctggatta cttatttatt tacaagtcga atgggttctg acgagaatca gaatgaccg 21900  
 caggcagtga tacgttgcgg aactgatact tgggttgcca cttgaattcg ggaatcacca 21960  
 acttgggaac cggatatcg ggcaggatgt cactccacag cttcttggtc agctgcaaag 22020  
 5 ctccaagcag gtcaggagcc gaaatcttga aatcacaatt aggaccagtg ctctgagcgc 22080  
 gagagttgcg gtacaccgga ttgcagcact gaaacacat cagcgacgga tgtctcacgc 22140  
 ttgccagcac ggtgggatct gcaatcatgc ccacatccag atcttcagca ttggcaatgc 22200  
 tgaacggggt catcttcag gtctgcctac ccatggcggg cacccaatta ggcttgggt 22260  
 tgcaatcgca gtgcagggg atcagtatca tcttgccctg atctgtctg attctggat 22320  
 10 acacggctct catgaaagca tcatattgct tgaaagcctg ctgggcttta ctacctcg 22380  
 tataaaacat cccgaggac ctgctgaaa actggttagc tgcacagccg gcattatca 22440  
 cacagcagcg ggcgtcattg ttggtattt gcaccacact tctgcccag cggtttggg 22500  
 tgatttgggt tgcctcgga ttctcttta aggtcgttg tccgttctg ctggccacat 22560  
 ccatctcgat aatctgtcc ttctgaatca taatattgcc atgcaggcac ttcagcttg 22620  
 15 cctcataatc attgcagcca tgaggccaca acgcacagcc tgtacattcc caattatgg 22680  
 gggcagatcg agaaaaagaa tgtatcattc cctgcagaaa tcttccatc atcgtgtca 22740  
 gtgtcttggt actagtgaat gtaactgga tgcctcgtg ctctcgtt acgtactgg 22800  
 gacagatgcg cttgtattgt tctgttctg caggcattag tttaaaacag gttctaagt 22860  
 cgttatccag cctgtactt tccatcagca gacacatcac ttcattgct ttctcccaag 22920  
 20 cagacaccag gggcaagcta atcggttct taacagtga ggcagcagct ccttagcca 22980  
 gagggtcctc tttagcgatc ttctaatgc ttcttttgc atcttctca acgatgcga 23040  
 cgggcgggta gctgaaaccc actgctacaa gttagcctc ttcttttct tcttctgt 23100  
 cttgactgat gtcttgcatg gggatatgt tggcttctt ttgggggta 23160  
 tccgaggagg aggactgtc ctccgtccg gagacagga ggattgtgac gttcgtca 23220  
 25 ccattacca ctgactgtc gtagaagaac ctgaccac acggcgacag gtgttttct 23280  
 tggggggcag aggtggagc gattgcgaag gctgcggtc cgacctgga ggcgatgac 23340  
 tggcagaacc cttccgct tgggggtgt gctcctgtg gcggtcgtt aactgattc 23400  
 cttcgcggt ggccattgt ttctctagg cagagaaaca acagacatgg aaactcagc 23460  
 attgtgtca acatgccac gattgccatc acatctctc ctacgcagc aggaagaag 23520  
 30 gcagagctta agcattccac cggcagtc tgcaccac tctacctag aagataagga 23580  
 ggtcgacga tctcatgaca tgcagaataa aaaagcgaaa gattctgaga cagacatga 23640  
 gcaagacccg ggtatgtga caccggtgga acacagggaa gattgaaac gctttctaga 23700  
 gagagaggat gaaaactgcc caaacagcg agcagataac tataccaag atgtggaaa 23760  
 tagggatcag aacaccgact acctatagg gcttgacgg gaagacgcgc tcttaaca 23820  
 35 tctagcaaga cagtgcctca tagtcaagga tgcattattg gacagaactg aagtgcctc 23880  
 cagtgtgga gagtcagct gcgctacga gcttaacct tttcacctc gtactcccc 23940  
 caaacgtcag ccaaacggca cctgcagcc aaatcctgc ttaacttt atccagctt 24000  
 tgctgtgcca gaagtactg ctacctatca catctttt aaaaatcaaa aaattccagt 24060  
 ctctgcccgc gtaatcgca cccgcgcca tgcctactc aatctgggac ctggttcag 24120  
 40 cttacctgat atagcttct tgaagaggt tccaaagatc ttcgagggtc tgggcaata 24180  
 tgagactcg ggcgaatg ctctgcaaaa gggagaaaat ggcatggatg agcatcacag 24240  
 cgttctggtg gaattggaag gcgataatgc cagactcgca gtactcaagc gaagcgtca 24300  
 ggtcacacac ttgcatac ccgtgtcaa cctgccccct aaagtcatga cggcggtcat 24360  
 ggaccagta ctcattaagc gcgcaagtc ctttcagaa gacatgcatg acccagatg 24420  
 45 ctgtgatgag ggtaaaccag tggtcagtga tgagcagcta acccgatggc tgggcaccga 24480  
 ctctccccgg gatttgaag agcgtcgca gcttatgat gccgtggtc tggttaccgt 24540  
 agaactagag tgtctccgac gtttcttac cgattcagaa acctgcgca aactcgaaga 24600  
 gaactgcac tacacttta gacacggct tgtgcggcag gcatgcaaga tatctaact 24660  
 ggaactcacc aacctggtt cctacatggg tattctgcat gagaatgcc taggacaaag 24720  
 50 cgtgctgac agcaccctta agggggaag ccgcgtgat tacatccgc attgtgtca 24780  
 tctctacctg tgcacacgt ggcaaacgg catgggtgta tggcagcaat gtttagaaga 24840  
 acagaactg aaagagctt acaagctct acagaaatct ctttaagggtc tgtggacagg 24900  
 gttcgacgag cgcaccgtc cttccgact ggcagacct atctccagc agcgtctcag 24960

ggtaactttg cgaaacggat tgctgactt tatgagccag agcatgctta acaattttcg 25020  
 ctctttcacc ctggaacgt cgggtatcct gccgcccacc tgcgcccac tgcctccga 25080  
 ctttgtgct ctcacctacc gcgagtcccc ccgcccgtta tggagtcact gctacctgtt 25140  
 5 ccgtctggcc aactatctct cctaccactc gcatgtgac gaggatgtga gcggagacgg 25200  
 cttgctggag tgccactgcc gctgcaatct gtgcacgcc caccggtccc tagcttgcaa 25260  
 cccccagttg atgagcgaaa ccagataat aggcaccttt gaattgcaag gcccagcag 25320  
 ccaaggcgat gggctctctc ctgggcaaag tttaaaactg accccgggac tgtggacctc 25380  
 cgctacttg cgcaagtttg ctccggaaga ttaccacccc tatgaaatca agttctatga 25440  
 10 ggaccaatca cagcctccaa aggcggaact ttccggttgc gtcacaccc agggggcaat 25500  
 tctggccaa ttgcaagcca tccaaaaatc ccgccaagaa ttctactga aaaagggtaa 25560  
 gggggtctac cttgaccccc agaccggcga ggaactcaac acaagggtcc ctcaggatgt 25620  
 cccaacgacg agaaaacaag aagtgaagg tgcagccgc gcccccagaa gatatggagg 25680  
 aagattggga cagtcaggca gaggaggcgg aggaggacag tctggaggac agtctggagg 25740  
 15 aagacagttt ggaggaggaa aacgaggagg cagaggaggt ggaagaagta accgccgaca 25800  
 aacagttatc ctggctgcg gagacaagca acagcgctac catctccgt ccgagtcgag 25860  
 gaaccggcg gcgtcccagc agtagatggg acgagaccgg acgttcccg aaccaacca 25920  
 gcgttccaa gaccgtaag aaggatcggc agggatacaa gtctggcgg gggcataaga 25980  
 atgccatcat ctctgcttg catgagtgcg ggggcaacat atcttcacg cggcgtact 26040  
 20 tgctattcca ccatgggtg aactttcgc gcaatgttt gcattactac cgtcacctcc 26100  
 acagcccta ctatagccag caaatccga cagtctgac agataaagac agcggcgcg 26160  
 acctcaaca gaaaaccagc agcggcagtt agaaaataca caacaagtgc agcaacagga 26220  
 ggattaaaga ttacagccaa cgagccagcg caaacccgag agttaagaaa tcggatcttt 26280  
 ccaacctgt atgccatctt ccagcagagt cggggtcaag agcaggaact gaaaataaaa 26340  
 25 aaccgatctc tgcgtcgt caccagaagt tgtttgtat acaagagcga agatcaactt 26400  
 cagcgactc tcgaggacgc cgaggtctc ttaacaagt actgcgcgt gactcttaa 26460  
 gaggtagcag cgaccgctt tattcaaaaa agcggggaat tacatcatcc tcgacatgag 26520  
 taaagaaatt ccacgcctt acatgtggag ttatcaacc caaatgggat tggcagcagg 26580  
 cgctcccag gactactcca ccgcatgaa ttggctcag gccgggctt ctatgattc 26640  
 30 tcgagttaat gatatacgc cctaccgaaa ccaataact ttggaacagt cagctcttac 26700  
 caccacgcc cgccaacacc ttaatccag aaattggccc gccgcctag tgtaccagga 26760  
 aagtcgcgt ccaccactg tattacttc tcgagacgc caggccgaag tccaatgac 26820  
 taatgcaggt gcgcagtag ctggcggtc caccatgt cgtcacagg ctcggcataa 26880  
 tataaacgc ctgatgatca gagccgagg tatccagtc aacgacgagt cgtgagctc 26940  
 35 tccgttgg ctacgaccag acggaatct tcagattgc ggtgcggga gatcttctt 27000  
 caccctcgt caggtgttc tgactttga aagtctgt tcgcaaccgc gtcgggcgg 27060  
 aatcgggacc gttaatttg tagaggagtt tactctctt gtctactca acccttctc 27120  
 cggatctct gggcactacc cggacgagt cataccgaac ttcagcga ttagcagtc 27180  
 agtggacgc tacgattgat gtctggtag gcgctgagc tatctcggt gcgacatca 27240  
 40 gaccactgcc gccgtttc ctgctttgc cgggaacta ttgagttcat ctactcgaa 27300  
 ctcccagg atcacctca aggtccggc caggagtg gcgattact cgaaggcaaa 27360  
 atagactct gcctgcaac aattttctc cagcgcccg tctgatcga gcgagaccag 27420  
 ggaaacacca cggtttcat ctactgcat tgaatcacc ccgattgca tgaaagcctt 27480  
 tctgtctta tgtgtactga gttaataaa aactgaatia agactctct acggactgcc 27540  
 45 gcttctcaa ccggatttt acaaccagaa gaacaaaact ttctgtcg tccaggactc 27600  
 tgttaactc accttcta ctacaaact agaagtcga cgactacac gctttccag 27660  
 aagcatttc cctactaata ctacttcaa aaccggaggt gagctccac gctccctac 27720  
 agaaaacct tgggtggaag cggccttgt agtactagga attcttgcg gtggcctgt 27780  
 gattattct tctacctat acacacctt ctacttct ctagtggtgt tctggtatt 27840  
 50 gtttaaaaa tggggccat actagtctt ctgttttac ttctgtttt ggaaccgggt 27900  
 tctgcaatt acgatcatg tctagactt gaccagaaa actgcacact tacttttga 27960  
 cccgacaaa gccgatctg tggagtctt attaatgag gatgggaat caggtccgtt 28020  
 gaaattacac acaatacaa aacctggaac aatacctat ccaccatg ggagccagga 28080



gttcccgagt ggtacactgt ctctgtccga ggctctgacg gttccatccg cattagtaac 28140  
 aacactttca ttttttctga aatgtgcgat ctggccatgt tcatgagcaa acagtattct 28200  
 ctatggcctc ctagcaagga caacatcgta acgtttccca ttgcttattg ctgtgcgct 28260  
 5 tgccttctta ctgctttact gtgcgtatgc atacacctgc ttgtaaccac tcgcatcaaa 28320  
 aacgccaata acaaaagaaa aatgccttaa cctctttctg tttacagaca tggcttctct 28380  
 tacatctctc atattgtca gcattgtcac tgcgctcac ggacaaacag tcgtctctat 28440  
 cccactagga cataattaca ctctcatagg accccaatc acttcagagg tcatctggac 28500  
 caaactggga agcgttgatt actttgatat aatctgtaac aaaacaaaac caataatagt 28560  
 10 aacttgcaac atacaaaatc ttacattgat taatgttagc aaagtttaca gcggttacta 28620  
 ttatggttat gacagataca gtagtcaata tagaaattac ttggttcgtg ttaccagtt 28680  
 gaaaaccacg aaaatgccaa atatggcaaa gattcgatcc gatgacaatt ctctagaaac 28740  
 tttacatct cccaccacac ccgacgaaaa aaacatccca gattcaatga ttgcaattgt 28800  
 tgcagcgggtg gcagtgggtga tggcactaat aataatatgc atgcttttat atgcttgcg 28860  
 15 ctacaaaaag tttcatccta aaaaacaaga tctctacta aggettaaca tttattttct 28920  
 tttatacag ccatggtttc cactaccaca ttcttatgc ttactagtct cgcaactctg 28980  
 actctgctc gctcacact cactgtaact ataggctcaa actgcacact aaaaggacct 29040  
 caagggtggtc atgtcttttg gtggagaata tatgacaatg gatggtttac aaaaccatgt 29100  
 gaccaacctg gtagattttt ctgcaacggc agagacctaa ccattatcaa cgtgacagca 29160  
 20 aatgacaaag gcttctatta tggaaacgac tataaaagta gtttagatta taacattatt 29220  
 gtactgcat ctaccactcc agcacccgc acaactactt tctctagcag cagtgtcgt 29280  
 aacaatacaa ttccaatcc aaccttgcc gcgttttaa aacgcactgt gaataattct 29340  
 acaactcac atacaacaat ttccactca acaatcagca tcatgctgc agtgacaatt 29400  
 ggaatatcta ttctgtttt taccataacc tactacgct gctgctatag aaaagacaaa 29460  
 25 cataaagggtg atccattact tagattgat atttaattg ttctttttt ttatttacag 29520  
 tatggtgaac accaatcatg gtacctagaa atttcttct caccatactc atctgtgctt 29580  
 ttaatgttg cgtactttc acagcagtag ccacagcaac ccagactgt ataggagcat 29640  
 ttgcttcta tgcactttt gctttgtta ctgcatctg cgtatgtac atagtctgc 29700  
 tggttattaa tttttccaa ctctagact ggatcctgt gcgaattgcc tacctgcgc 29760  
 30 accatccga ataccgcaac caaatatcg cggcactct tagactcctc taaaaccatg 29820  
 caggctatac taccaatatt ttgttcta ttgttccct acgtgtctc aacccagct 29880  
 gcctatagta ctccaccaga acacctaga aaatgcaaat tccaacaacc gtggtcattt 29940  
 ctgtgtgtc atcgagaaaa atcagaaatc ccccaaat taataatgat tgetggaata 30000  
 attaatataa tctgttgac cataattca ttttgatat accccctatt tgattttgc 30060  
 35 tggaaatgct ccaatgcaca tgatcatcca caagaccag aggaacacat tccccacaa 30120  
 aacatgcaac atccaatagc gctaatagat tacgaaagt aaccacaacc cccactactc 30180  
 cctgtatta gttactcaa cctaaccgc ggagatgact gaaacactca ccactccaa 30240  
 ttccgcgag gatctgctc atatggacgg ccgctctca gaacaacgac ttgccaact 30300  
 acgcatccgc cagcagcagg aacgctggc caaagagctc agagatgca tccaaattca 30360  
 40 ccaatgcaaa aaaggcatat tctgtttgtt aaaacaagcc aagatattct acgagatcac 30420  
 cgtactgac catcgctct ctacgaact tggcccccac cgacaaaaat ttacctgcat 30480  
 ggtgggaatc aacccatag ttatcacca acaaagtga gatactaagg gttgcattca 30540  
 ctgctctgc gattccatcg agtgaccta cacctgctg aagacctat geggctaag 30600  
 agacctgcta ccaatgaatt aaaaaaaat gattaataaa aatcactta ctgaaatca 30660  
 45 gcaataagg ctctgttga attttctcc agcagcact cactccctc tteccaactc 30720  
 tggattcta aacccgttc agcgcatatc ttctccata cttaaagg gatgtcaat 30780  
 tttagctct ctctgtacc cacaatttc atgttttct tccagatga ccaagagagt 30840  
 ccgctcagt gactcttca acctgtcta cccctatga gatgaaagca cctcccaaca 30900  
 ccccttata aaccagggt ttatttccc aaatggctc acacaaagcc cagacggagt 30960  
 50 tcttacttta aaatgttaa cccactaac aaccacagc ggatctctac agctaaaagt 31020  
 gggaggggga ctacagtgg atgacactga tggtaacct caagaaaaca tacgtctac 31080  
 agcaacctt actaaaaata atcactctgt agaactatc attggaatg gattagaac 31140  
 tcaaaacaat aaactatgt ccaattggg aaatgggtta aaatttaaca acggtgacat 31200

ttgtataaag gatagtatta acaccttatg gactggaata aacctccac ctaactgtca 31260  
 aattgtggaa aacactaata caaatgatgg caaacttact ttagtattag taaaaaatgg 31320  
 agggcttgtt aatggctacg tgtctctagt tgggtatca gacactgtga accaaatgtt 31380  
 5 cacacaaaag acagcaaaaca tccaattaag attatatttt gactcttctg gaaatctatt 31440  
 aactgaggaa tcagacttaa aaattccact taaaaataaa tcttctacag cgaccagtga 31500  
 aactgtagcc agcagcaaag cctttatgcc aagtactaca gtttatccct tcaacaccac 31560  
 tactagggat agtgaanaact acattcatgg aatatgttac tacatgacta gttatgatag 31620  
 aagtctattt ccttgaaca tttctataat gctaaacagc cgtatgattt ctccaatgt 31680  
 10 tgcctatgcc atacaatttg aatggaatct aaatgcaagt gaatctccag aaagcaacat 31740  
 agctacgctg accacatccc ccttttctt tctttacatt acagaagacg acaactaaaa 31800  
 taaagtttaa gtgtttttat ttaaaatcac aaaatcgag tagttatttt gcctccacct 31860  
 tccatttga cagaatacac caatctctcc ccacgcacag cttaaacat ttggatacca 31920  
 ttagagatag acattgtttt agattccaca ttccaaacag tttcagagcg agccaatctg 31980  
 15 gggtcagtga tagataaaaa tccatcgca tagtctttta aagcgcttc acagtccaac 32040  
 tgcctcggat gcgactccgg agtttggatc acggtcatct ggaagaagaa cgtatgggaat 32100  
 cataatccga aaacggtatc ggacgattgt gtcctatcaa acccacaagc agccgctgtc 32160  
 tgcgtcgtc cgtgcgactg ctgtttatgg gatcagggtc cacagtttcc tgaagcatga 32220  
 ttttaatagc ccttaacatc aactttctgg tgcgatgccc gcagcaacgc attctgattt 32280  
 20 cactcaaatc tttgcagttag gtacaacaca ttattacaat attgtttaat aaaccataat 32340  
 taaaagcgct ccagccaaaa ctcatatctg atataatgc ccttgcatac ccatcatacc 32400  
 aaagttaat ataaatlaaa tgacgttccc taaaaaacac actaccacac tacatgatct 32460  
 ctttggcat gtcatatta acaatctgtc tgtaccatgg acaacgttgg ttaatcatgc 32520  
 aacccaatat aaccttccgg aaccacactg ccaacacgc tccccagcc atgcattgaa 32580  
 25 gtgaacctg ctgattacaa tgacaatgaa gaaccaatt ctctcgaccg tgaatcactt 32640  
 gagaatgaaa aatatctata gtggcacaac atagacataa atgcattcat ctctcataa 32700  
 ttttaactc ctgaggattt agaaacatat cccaggggaat aggaagctct tgcagaacag 32760  
 taaagctggc agaacaagga agaccagaa cacaacttac actatgcata gtcattat 32820  
 cacaatctgg caacagcggg tggcttctag tcatagaagc tgggtttca tttctctac 32880  
 30 aacgtggtaa ctgggctctg gtgtaagggt gatgtctggc gcatgatgtc gagcgtgccc 32940  
 gcaacctgt cataatggag ttgcttctg acattctgt attttgata gcaaaacgcg 33000  
 gcctggcag aacacactct tcttgcctt ctatcctgcc gcttagcgtg ttcggtgta 33060  
 tagttcaagt acagccacac tcttaagtgt gcaaaaagaa tgcgtgctc agttgtaatc 33120  
 aaaactccat cgcattctat tgtctgagg aaatcatcca cggtagcata tgcaaatccc 33180  
 35 aaccaagcaa tgcaactgga ttgcgttca agcaggagag gagagggaag agacggaaga 33240  
 accatgttaa tttttattc aaacgatctc gcagtactc aaattgtaga tgcgcagat 33300  
 ggcatctctc gccccactg tgttgggtgaa aaagcacagc taaatcaaaa gaaatgcgat 33360  
 ttcaagggtg ctcaacggtg gttccaaca aagcctccac gcgcacatcc aagaacaaaa 33420  
 gaatacaaaa agaaggagca ttttctaact ctcaatcat catattacat tctgcacca 33480  
 40 tcccagata attttcagct tccagcctt gaattattc tgcagttct tgtggtaaat 33540  
 ccaatccaca cattacaaac aggtcccggg gggcgccctc caccaccatt cttaaacaca 33600  
 cctcataat gacaaaatat cttgtcctg tgcacctgt agcgaattga gaatggcaac 33660  
 atcaattgac atgcccctgg ctctaagttc tctttaagt tctagttgta aaaactctct 33720  
 catattatca ccaactgct tagccagaag cccccggg acaagagcag gggacgctac 33780  
 45 agtgcagtac aagcgcagac ctcccaatt ggctccagca aaaacaagat tggaataagc 33840  
 atattgggaa ccaccagtaa tatcatcgaa gttcgtgaa atataatcag gcagagttc 33900  
 tttagaaat tgaataaaag aaaaatttgc caaaaaaca ttcaaacct ctgggatgca 33960  
 aatgcaatag gttaccgccc tgcgtccaa cattgttagt tttgaattag tctgcaaaa 34020  
 taaaaaaaaa acaagcgtca tatcatagta gcctgacgaa caggtggata aatcagtctt 34080  
 50 tccatcaca gacaagccac aggtcttcca gtcgacct cgtaaaacct gtcacgtga 34140  
 ttaacaaca gcaccgaaag ttctcgcggg tgaccagcat gaataagtct tgatgaagca 34200  
 tacaatccag acatgttagc atcagttaag gagaaaaaac agccaacata gcctttgggt 34260  
 ataattatgc ttaatcgtaa gtatagcaaa gccacccctc gggatataa agtaaaaggc 34320



acaggagaat aaaaaatata attatttctc tgctgctgtt taggcaacgt cgcccccggt 34380  
 ccctctaaat acacatacaa agcctcatca gccatggett accagagaaa gtacagcggg 34440  
 cacacaaacc acaagctcta aagtcactct ccaacctstc cacaatatat atacacaagc 34500  
 cctaaactga cgtaatggga ctaaagtgt aaaaatcccg ccaaaccxaa cacacacccc 34560  
 gaaactgagt caccagggaa aagtacagt tcacttcgc aatcccaaca agcgtcactt 34620  
 cctctttctc acggtacgtc acatccatt aacttacaac gtcattttcc cacggccgcg 34680  
 ccgccccitt taaccgttaa cccacagcc aatcaccaca cgccccacac tttttaaatt 34740  
 cacctcattt acatattggc accattccat ctataaggta tattattgat gatg 34794

<210> 85

<211> 585

<212> PRT

<213> Adenoviridae

<220>

<221> CHAIN

<222> (1)..(585)

<223> /note="Fiber protein of Adenovirus serotype 5"

<400> 85

Met Leu Leu Gln Met Lys Arg Ala Arg Pro Ser Glu Asp Thr Phe Asn.  
 1 5 10 15

Pro Val Tyr Pro Tyr Asp Thr Glu Thr Gly Pro Pro Thr Val Pro Phe  
 20 25 30

Leu Thr Pro Pro Phe Val Ser Pro Asn Gly Phe Gln Glu Ser Pro Pro  
 35 40 45

Gly Val Leu Ser Leu Arg Leu Ser Glu Pro Leu Val Thr Ser Asn Gly  
 50 55 60

Met Leu Ala Leu Lys Met Gly Asn Gly Leu Ser Leu Asp Glu Ala Gly  
 65 70 75 80

Asn Leu Thr Ser Gln Asn Val Thr Thr Val Ser Pro Pro Leu Lys Lys  
 85 90 95

Thr Lys Ser Asn Ile Asn Leu Glu Ile Ser Ala Pro Leu Thr Val Thr  
 100 105 110

Ser Glu Ala Leu Thr Val Ala Ala Ala Pro Leu Met Val Ala Gly  
 115 120 125

Asn Thr Leu Thr Met Gln Ser Gln Ala Pro Leu Thr Val His Asp Ser  
 130 135 140

Lys Leu Ser Ile Ala Thr Gln Gly Pro Leu Thr Val Ser Glu Gly Lys  
 145 150 155 160

Leu Ala Leu Gln Thr Ser Gly Pro Leu Thr Thr Thr Asp Ser Ser Thr

	165	170	175
5	Leu Thr Ile Thr Ala Ser Pro Pro Leu Thr Thr Ala Thr Gly Ser Leu		
	180	185	190
	Gly Ile Asp Leu Lys Glu Pro Ile Tyr Thr Gln Asn Gly Lys Leu Gly		
	195	200	205
10	Leu Lys Tyr Gly Ala Pro Leu His Val Thr Asp Asp Leu Asn Thr Leu		
	210	215	220
	Thr Val Ala Thr Gly Pro Gly Val Thr Ile Asn Asn Thr Ser Leu Gln		
15	225	230	235 240
	Thr Lys Val Thr Gly Ala Leu Gly Phe Asp Ser Gln Gly Asn Met Gln		
	245	250	255
20	Leu Asn Val Ala Gly Gly Leu Arg Ile Asp Ser Gln Asn Arg Arg Leu		
	260	265	270
	Ile Leu Asp Val Ser Tyr Pro Phe Asp Ala Gln Asn Gln Leu Asn Leu		
	275	280	285
25	Arg Leu Gly Gln Gly Pro Leu Phe Ile Asn Ser Ala His Asn Leu Asp		
	290	295	300
	Ile Asn Tyr Asn Lys Gly Leu Tyr Leu Phe Thr Ala Ser Asn Asn Ser		
30	305	310	315 320
	Lys Lys Leu Glu Val Asn Leu Ser Thr Ala Lys Gly Leu Met Phe Asp		
	325	330	335
35	Ala Thr Ala Ile Ala Ile Asn Ala Gly Asp Gly Leu Glu Phe Gly Ser		
	340	345	350
	Pro Asn Ala Pro Asn Thr Asn Pro Leu Lys Thr Lys Ile Gly His Gly		
	355	360	365
40	Leu Glu Phe Asp Ser Asn Lys Ala Met Val Pro Lys Leu Gly Thr Gly		
	370	375	380
	Leu Ser Phe Asp Ser Thr Gly Ala Ile Thr Val Gly Asn Lys Asn Asn		
	385	390	395 400
45	Asp Lys Leu Thr Leu Trp Thr Thr Pro Ala Pro Ser Pro Asn Cys Arg		
	405	410	415
	Leu Asn Ala Glu Lys Asp Ala Lys Leu Thr Leu Val Leu Thr Lys Cys		
50	420	425	430
	Gly Ser Gln Ile Leu Ala Thr Val Ser Val Leu Ala Val Lys Gly Ser		
	435	440	445

55

Leu Ala Pro Ile Ser Gly Thr Val Gln Ser Ala His Leu Ile Ile Arg  
450 455 460

Phe Asp Glu Asn Gly Val Leu Leu Asn Asn Ser Phe Leu Asp Pro Glu  
465 470 475 480

Tyr Trp Asn Phe Arg Asn Gly Asp Leu Thr Glu Gly Thr Ala Tyr Thr  
485 490 495

Asn Ala Val Gly Phe Met Pro Asn Leu Ser Ala Tyr Pro Lys Ser His  
500 505 510

Gly Lys Thr Ala Lys Ser Asn Ile Val Ser Gln Val Tyr Leu Asn Gly  
515 520 525

Asp Lys Thr Lys Pro Val Thr Leu Thr Ile Thr Leu Asn Gly Thr Gln  
530 535 540

Glu Thr Gly Asp Thr Thr Pro Ser Ala Tyr Ser Met Ser Phe Ser Trp  
545 550 555 560

Asp Trp Ser Gly His Asn Tyr Ile Asn Glu Ile Phe Ala Thr Ser Ser  
565 570 575

Tyr Thr Phe Ser Tyr Ile Ala Gln Glu  
580 585

<210> 86  
<211> 356  
<212> PRT  
<213> Adenoviridae

<220>  
<221> CHAIN  
<222> (1)..(356)  
<223> /note="Fiber protein of Adenovirus serotype 16"

<400> 86  
Met Leu Leu Gln Met Lys Arg Ala Arg Pro Ser Glu Asp Thr Phe Asn  
1 5 10 15

Pro Val Tyr Pro Tyr Glu Asp Glu Ser Ser Ser Gln His Pro Phe Ile  
20 25 30

Asn Pro Gly Phe Ile Ser Ser Asn Gly Phe Ala Gln Ser Pro Asp Gly  
35 40 45

Val Leu Thr Leu Lys Cys Val Asn Pro Leu Thr Thr Ala Ser Gly Pro  
50 55 60

5 Leu Gln Leu Lys Val Gly Ser Ser Leu Thr Val Asp Thr Ile Asp Gly  
 65 70 75 80  
 Ser Leu Glu Glu Asn Ile Thr Ala Ala Ala Pro Leu Thr Lys Thr Asn  
 85 90 95  
 10 His Ser Ile Gly Leu Leu Ile Gly Ser Gly Leu Gln Thr Lys Asp Asp  
 100 105 110  
 Lys Leu Cys Leu Ser Leu Glu Asp Gly Leu Val Thr Lys Asp Asp Lys  
 115 120 125  
 15 Leu Cys Leu Ser Leu Gly Asp Gly Leu Ile Thr Lys Asn Asp Val Leu  
 130 135 140  
 Cys Ala Lys Leu Gly His Gly Leu Val Phe Asp Ser Ser Asn Xaa Ile  
 145 150 155 160  
 20 Thr Ile Glu Asn Asn Thr Leu Trp Thr Gly Ala Lys Pro Ser Ala Asn  
 165 170 175  
 Cys Val Ile Lys Glu Gly Glu Asp Ser Pro Asp Cys Lys Leu Thr Leu  
 180 185 190  
 25 Val Leu Val Lys Asn Gly Gly Leu Ile Asn Gly Tyr Ile Thr Leu Met  
 195 200 205  
 30 Gly Ala Ser Glu Tyr Thr Asn Thr Leu Phe Lys Asn Asn Gln Val Thr  
 210 215 220  
 Ile Asp Val Asn Leu Ala Phe Asp Asn Thr Gly Gln Ile Ile Thr Tyr  
 225 230 235 240  
 35 Leu Ser Ser Leu Lys Ser Asn Leu Asn Phe Lys Asp Asn Gln Asn Met  
 245 250 255  
 40 Ala Thr Gly Thr Ile Thr Ser Ala Lys Gly Phe Met Pro Ser Thr Thr  
 260 265 270  
 Ala Tyr Pro Phe Ile Thr Tyr Ala Thr Glu Thr Leu Asn Glu Asp Tyr  
 275 280 285  
 45 Ile Tyr Gly Glu Cys Tyr Tyr Lys Ser Thr Asn Gly Thr Leu Phe Pro  
 290 295 300  
 Leu Lys Val Thr Val Thr Leu Asn Arg Arg Met Leu Ala Ser Gly Met  
 305 310 315 320  
 50 Ala Tyr Ala Met Asn Phe Ser Trp Ser Leu Asn Ala Glu Glu Ala Pro  
 325 330 335

55

Glu Thr Thr Glu Val Thr Leu Ile Thr Ser Pro Phe Phe Phe Ser Tyr  
 340 345 350

Ile Arg Glu Asp  
 355

<210> 87  
 <211> 325  
 <212> PRT  
 <213> Adenoviridae

<220>  
 <221> CHAIN  
 <222> (1)..(325)  
 <223> /note="Fiber protein of Adenovirus serotype 35"

<400> 87  
 Met Leu Leu Gln Met Lys Arg Ala Arg Pro Ser Glu Asp Thr Phe Asn  
 1 5 10 15

Pro Val Tyr Pro Tyr Glu Asp Glu Ser Thr Ser Gln His Pro Phe Ile  
 20 25 30

Asn Pro Gly Phe Ile Ser Pro Asn Gly Phe Thr Gln Ser Pro Asp Gly  
 35 40 45

Val Leu Thr Leu Lys Cys Leu Thr Pro Leu Thr Thr Thr Gly Gly Ser  
 50 55 60

Leu Gln Leu Lys Val Gly Gly Gly Leu Thr Val Asp Asp Thr Asp Gly  
 65 70 75 80

Thr Leu Gln Glu Asn Ile Arg Ala Thr Ala Pro Ile Thr Lys Asn Asn  
 85 90 95

His Ser Val Glu Leu Ser Ile Gly Asn Gly Leu Glu Thr Gln Asn Asn  
 100 105 110

Lys Leu Cys Ala Lys Leu Gly Asn Gly Leu Lys Phe Asn Asn Gly Asp  
 115 120 125

Ile Cys Ile Lys Asp Ser Ile Asn Thr Leu Trp Thr Gly Ile Asn Pro  
 130 135 140

Pro Pro Asn Cys Gln Ile Val Glu Asn Thr Asn Thr Asn Asp Gly Lys  
 145 150 155 160

Leu Thr Leu Val Leu Val Lys Asn Gly Gly Leu Val Asn Gly Tyr Val  
 165 170 175



5

Ser Leu Val Gly Val Ser Asp Thr Val Asn Gln Met Phe Thr Gln Lys  
180 185 190

Thr Ala Asn Ile Gln Leu Arg Leu Tyr Phe Asp Ser Ser Gly Asn Leu  
195 200 205

10

Leu Thr Glu Glu Ser Asp Leu Lys Ile Pro Leu Lys Asn Lys Ser Ser  
210 215 220

Thr Ala Thr Ser Glu Thr Val Ala Ser Ser Lys Ala Phe Met Pro Ser  
225 230 235 240

15

Thr Thr Ala Tyr Pro Phe Asn Thr Thr Thr Arg Asp Ser Glu Asn Tyr  
245 250 255

Ile His Gly Ile Cys Tyr Tyr Met Thr Ser Tyr Asp Arg Ser Leu Phe  
260 265 270

20

Pro Leu Asn Ile Ser Ile Met Leu Asn Ser Arg Met Ile Ser Ser Asn  
275 280 285

25

Val Ala Tyr Ala Ile Gln Phe Glu Trp Asn Leu Asn Ala Ser Glu Ser  
290 295 300

Pro Glu Ser Asn Ile Met Thr Leu Thr Thr Ser Pro Phe Phe Phe Ser  
305 310 315 320

30

Tyr Ile Thr Glu Asp  
325

35

<210> 88  
<211> 325  
<212> PRT  
<213> Adenoviridae

40

<220>  
<221> CHAIN  
<222> (1)..(325)  
<223> /note="Fiber protein of Adenovirus serotype 51"

45

<400> 88  
Met Leu Leu Gln Met Lys Arg Ala Arg Pro Ser Glu Asp Thr Phe Asn  
1 5 10 15

50

Pro Val Tyr Pro Tyr Glu Asp Glu Ser Thr Ser Gln His Pro Phe Ile  
20 25 30

Asn Pro Gly Phe Ile Ser Pro Asn Gly Phe Thr Gln Ser Pro Asp Gly  
35 40 45

55

EP 1 054 064 A1

Val Leu Thr Leu Asn Cys Leu Thr Pro Leu Thr Thr Thr Gly Gly Pro  
50 55 60

5 Leu Gln Leu Lys Val Gly Gly Gly Leu Ile Val Asp Asp Thr Asp Gly  
65 70 75 80

Thr Leu Gln Glu Asn Ile Arg Val Thr Ala Pro Ile Thr Lys Asn Asn  
85 90 95

10 His Ser Val Glu Leu Ser Ile Gly Asn Gly Leu Glu Thr Gln Asn Asn  
100 105 110

Lys Leu Cys Ala Lys Leu Gly Asn Gly Leu Lys Phe Asn Asn Gly Asp  
15 115 120 125

Ile Cys Ile Lys Asp Ser Ile Asn Thr Leu Trp Thr Gly Ile Lys Pro  
130 135 140

20 Pro Pro Asn Cys Gln Ile Val Glu Asn Thr Asp Thr Asn Asp Gly Lys  
145 150 155 160

Leu Thr Leu Val Leu Val Lys Asn Gly Gly Leu Val Asn Gly Tyr Val  
165 170 175

25 Ser Leu Val Gly Val Ser Asp Thr Val Asn Gln Met Phe Thr Gln Lys  
180 185 190

Ser Ala Thr Ile Gln Leu Arg Leu Tyr Phe Asp Ser Ser Gly Asn Leu  
30 195 200 205

Leu Thr Asp Glu Ser Asn Leu Lys Ile Pro Leu Lys Asn Lys Ser Ser  
210 215 220

35 Thr Ala Thr Ser Glu Ala Ala Thr Ser Ser Lys Ala Phe Met Pro Ser  
225 230 235 240

Thr Thr Ala Tyr Pro Phe Asn Thr Thr Thr Arg Asp Ser Glu Asn Tyr  
245 250 255

40 Ile His Gly Ile Cys Tyr Tyr Met Thr Ser Tyr Asp Arg Ser Leu Val  
260 265 270

Pro Leu Asn Ile Ser Ile Met Leu Asn Ser Arg Thr Ile Ser Ser Asn  
45 275 280 285

Val Ala Tyr Ala Ile Gln Phe Glu Trp Asn Leu Asn Ala Lys Glu Ser  
290 295 300

50 Pro Glu Ser Asn Ile Ala Thr Leu Thr Thr Ser Pro Phe Phe Phe Ser  
305 310 315 320

Tyr Ile Ile Glu Asp

55

Claims

1. A gene delivery vehicle comprising at least one of the adenovirus serotype 35 elements or a functional equivalent thereof, responsible for avoiding or deminishing neutralising activity against adenoviral elements by the host to which the gene is to be delivered and a gene of interest.
2. Adenovirus serotype 35 or a functional homologue thereof or a chimaeric virus derived therefrom, or a gene delivery vehicle based on said virus its homologue or its chimaera for use as a pharmaceutical.
3. A gene delivery vehicle according to claim 1, whereby said elements comprise adenovirus 35 E3 expression products or the genes encoding them.
4. A gene delivery vehicle according to claim 1 or 3, whereby said elements comprise adenovirus 35 fiber, penton and/or hexon proteins or a gene encoding either.
5. A gene delivery according to any one of claims 1,3, or 4 which is a chimaera of adenovirus 35 with at least one other adenovirus.
6. A gene delivery vehicle according to any one of claims 1,3,4 or 5 which has a different tropism than adenovirus 35.
7. A nucleic acid encoding at least a functional part of a gene delivery vehicle according to any one of claims 1 or 3-6, or a virus, homologue or chimaera thereof according to claim 2.
8. A nucleic acid encoding at least one of the adenovirus serotype 35 elements or a functional equivalent thereof, responsible for avoiding or deminishing neutralising activity against adenoviral elements by the host to which the gene is to be delivered and having a site for introducing a gene of interest therein.
9. A nucleic acid according to claim 7 or 8, further comprising at least one ITR.
10. A nucleic acid according to claim 7, 8 or 9 further comprising a region of nucleotides designed or useable for homologous recombination.
11. At least one set of two nucleic acids comprising a nucleic acid according to any one of claims 7-10, whereby said set of nucleic acids is capable of a single homologous recombination event with each other, which leads to a nucleic acid encoding a functional gene delivery vehicle.
12. A cell comprising a nucleic acid according to any one of claims 7-10 or a set of nucleic acids according to claim 11.
13. A cell according to claim 12 which complements the necessary elements for adenoviral replication which are absent from the nucleic acid according to any one of claims 7-10 or a set of nucleic acids according to claim 11.
14. A cell according to claim 12 or 13, which originates from a PER.C6 cell (ECACC deposit number 96022940).
15. A method for producing a gene delivery vehicle according to claim 1, or 3-6, comprising expressing a nucleic acid according to any one of claims 7-10 in a cell according to claim 12 or 13 and harvesting the resulting gene delivery vehicle.
16. A method for producing a gene delivery vehicle according to claim 1, or 3-6, comprising culturing a cell according to claim 12 or 13 in a suitable culture medium and harvesting the resulting gene delivery vehicle.
17. A gene delivery vehicle obtainable by a method according to claims 15 or 16.
18. A gene delivery vehicle according to any one of claims 1,3-6 or 17, which is derived from a chimaera of an adenovirus and an integrating virus.
19. A gene delivery vehicle according to claim 18, wherein said integrating virus is adeno associated virus.
20. A gene delivery vehicle according to any one of claims 1, 3-6 or 17-19, which has the tropism determining parts of

adenovirus 16 or functional equivalents thereof.

21. A gene delivery vehicle according to anyone of claims 1, 3-6 or 17-20 for use as a pharmaceutical.

5 22. A pharmaceutical formulation comprising a gene delivery vehicle according to any one of claims 1, 3-6 or 17-20 and a suitable excipient.

23. A pharmaceutical formulation comprising an adenovirus, a chimaera thereof, or a functional homologue thereof according to claim 2 and a suitable excipient.

10

15

20

25

30

35

40

45

50

55

**Figure 1:**  
**% of human sera with neutralising capacity for human adenovirus (n=100)**





Figure 2






Figure 3: Ad35 plasmid-based system for virus generation



**Figure 4**  
**Neutralisation in human sera**



**Figure 5**

**Neutralisation in human sera from different geographic locations**




Figure 6: Total sequence of Ad35.

```

1  CATCATCAAT AATATACCTT ATAGATGGAA TGGTGCCAAT ATGTAAATGA GGTGATTTTA AAAAGTGTGC
71  GCCGTGTGGT GATTGGCTGT GGGGTAAACG GTTAAAAGGG GCGGCGCGGC CGTGGGAAAA TGACGTTTTA
141 TGGGGGTGGA GTTTTTTTGC AAGTTGTGCG GGGAAATGTT ACGCATAAAA AGGCTTCTTT TCTCACGGAA
211 CTACTTAGTT TTCCACCGGT ATTTAACAGC AAATGACGTA GTTTTGACCG GATGCAAGTG AAAATTGCTG
281 ATTTTCGCGC GAAAACTGAA TGAGGAAGTG TTTTCTGAA TAATGTGGTA TTTATGCCAG GGTGGAGTAT
351 TTGTTACGGG CCAGGTAGAC TTTGACCCAT TACGTGGAGG TTTGATTAC CGTGTTTTTT ACCTGAATTT
421 CCGCGTACCG TGTCAAAGTC TTCTGTTTTT ACGTAGGTGT CAGCTGATCG CTAGGGTATT TATACCTCAG
491 GGTGTGTGTC AAGAGGCCAC TCTTGAGTGC CAGCGAGAAG AGTTTTCTCC TCTGCGCCGG CAGTTTAATA
561 ATAAAAAAT GAGAGATTTG CGATTTCTGC CTCAGGAAAT AATCTCTGCT GAGACTGGAA ATGAAATATT
631 GGAGCTTGTG GTGCACGCCC TGATGGGAGA CGATCCGGAG CCACCTGTGC AGCTTTTGA GCCTCCTACG
701 CTTCAGGAAC TGTATGATTT AGAGGTAGAG GGATCGGAGG ATTCTAATGA GGAAGCTGTG AATGGCTTTT
771 TTACCGATTG TATGCTTTTA GCTGCTAATG AAGGATTAGA ATTAGATCCG CTTTGGACA CTTCAATAC
841 TCCAGGGGTG ATTGTGGAAG GCGGTACAGG TGTAAGAAAA TTACCTGATT TGAGTTCCTG GGACTGTGAT
911 TTGCACTGCT ATGAAGACGG GTTCTCTCCG AGTGATGAGG AGGACCATGA AAAGGAGCAG TCCATGCAGA
981 CTGCAGCGGG TGAGGGAGTG AAGGCTGCCA ATGTTGGTTT TCAGTTGGAT TGCCCGGAGC TTCTGGACA
1051 TGGCTGTAA G TCTTGTGAAT TTCACAGGAA AAATACTGGA GTAAAGGAAC TGTTATGTTT GCTTTGTTAT
1121 ATGAGAACGC ACTGCCACTT TATTTACAGT AAGTGTGTTT AAGTTAAAT TTAAAGGAAT ATGCTGTTTT
1191 TCACATGTAT ATTGAGTGTG AGTTTTGTGC TTCTATTAT AGGTCTGTG TCTGATGCTG ATGAATCACC
1261 ATCTCCTGAT TCTACTACCT CACCTCCTGA TATTCAAGCA CCTGTTCTTG TGACGTGCG CAAGCCCATT
1331 CCGTGAAGC TTAAGCCTGG GAAACGTCCA GCAGTGGAGA AACTTGAGGA CTTGTTACAG GGTGGGGACG
1401 GACCTTTGGA CTTGAGTACA CGGAAACGTC CAAGACAATA AGTGTTCAT ATCCGTGTT ACTTAAGGTG
1471 ACGTCAATAT TTGTTGAGA GTGCAATGTA ATAAAAATAT GTTAALIGTT CACTGGTTTT TATGCTTTTT
1541 TGGGCGGGGA CTCAGGTATA TAAGTAGAAG CAGACCTGTG TGGTTAGCTC ATAGGAGCTG GCTTTCATCC
1611 ATGGAGGTTT GGGCCATTTT GGAAGACCTT AGGAAGACTA GGCAACTGTT AGAGAGCGCT TCGGACGGAG
1681 TCTCCGTTT TTGGAGATTC TGGTTCGCTA GTGAATTAGC TAGGGTAGTT TTTAGGATAA AACAGGACTA
1751 TAAACAAGAA TTTGAAAAGT TGTTGGTAGA TTGCCAGGA CTTTTTGAAG CTCTTAATTT GGGCCATCAG
1821 GTTCACTTTA AAGAAAAAGT TTTATCAGTT TTAGACTTTT CAACCCAGG TAGAACTGCT GCTGCTGTGG
1891 CTTTTCTTAC TTTTATATTA GATAAATGGA TCCCGCAGAC TCATTTTCAGC AGGGGATACG TTTTGGATTT
1961 CATAGCCACA GCATTGTGGA GAACATGGAA GGTTCGCAAG ATGAGGACAA TCTTAGGTTA CTGGCCAGTG
2031 AAGCCTTTGG GTGTAGCGGG AATCCTGAGG CATCCACCGG TCATGCCAGC GGTTCTGGAG GAGGAACAGC
2101 AAGAGACAA CCGGAGAGCC GGCTGGACCT CTCCAGTGGG GGAGGCGGAG TAGCTGACTT GTCTCTGAA
2171 CTGCAACGGG TGCTTACTGG ATCTACGCTC ACTGGACGGG ATAGGGCGGT TAAGAGGGAG AGGGCATCCA
2241 GTGGTACTGA TGCTAGATCT GAGTTGGCTT TAAGTTTAAT GAGTCGAGA CGTCTGAAA CCATTTGGTG
2311 GCATGAGGTT CAGAAAGAGG GAAGGGATGA AGTTTCTGTA TTGCAGGAGA AATATTCACT GGAACAGGTG
2381 AAAACATGTT GGTGGAGCC AGAGGATGAT TGGGCGGTGG CCATTAAAAA TTATGCCAAG ATAGCTTTGA
2451 GGCCTGATAA ACAGTATAAG ATCAGTAGAC GGATTAATAT CCGGAATGCT TGTTACATAT CTGGAATGG
2521 GGCTGAGGTG GTAATAGATA CTCAAGACAA GACAGTTATT AGATGCTGCA TGATGGATAT GTGGCCTGGA
2591 GTAGTCGGTA TGGAAGCAGT CACTTTTGTA AATGTTAAGT TTAGGGGAGA TGGTTATAAT GGAATAGTGT
2661 TTATGGCCAA TACCAAACCT ATATTGCATG GTTGTAGCTT TTTTGGTTT CACAATACCT GTGTAGATGC
2731 CTGGGGACAG GTTAGTGATC GGGGGTGTAG TTTCTATGCG TGTGGATTG CCACAGCTGG CAGAACCAAG
2801 AGTCAATTGT CTCTGAAGAA ATGCATATTC CAAAGATGTA ACCTGGGCAT TCTGAATGAA GGCGAAGCAA
2871 GGGTCCGTCA CTGCGCTTCT ACAGATACGT GATGTTTTAT TTTAATTAAG GGAAATGCCA GCGTAAAGCA
2941 TAACATGATT TGTGGTGCTT CCGATGAGAG GCCTTCACTA AATGCTCTGT TTTTGATCAC AATGTGTTGA
3011 ATGCTGGCTA CTGTGCATAT TGTTCCTCAT CAACGCAAAA AATGGCCTGT TTTTGATCAC AATGTGTTGA
3081 CCAAGTGCAC CATGCATGCA GGTGGGCGTA GAGGAATGTT TATGCCTTAC CAGTGTAACA TGAATCATGT
3151 GAAAGTGTG TTGGAACCA GATGCTTTTC CAGAATGAGC CTAACAGGAA TCTTTGACAT GAACACGCAA
3221 ATCTGGAAGA TCCTGAGGTA TGATGATACG AGATCGAGGG TGCGCGCATG CGAATGCGGA GGCAAGCATG
3291 CCAGGTTCCA GCCGGTGTGT GTAGATGTGA CCGAAGATCT CAGACCGGAT CATTTGGTTA TTGCCCGCAC
3361 TGGAGCAGAG TTCGGATCCA GTGGAGAAGA AACTGACTAA GGTGAGTATT GGGAAAACCT TGGGGTGGGA
3431 TTTTCAGATG GACAGATTGA GTAAAAATTT GTTTTTCTG TCTTGCAGCT GACATGAGTG GAAATGCTTC
3501 TTTTAAGGGG GGAGTCTTCA GCCCTTATCT GACAGGGCGT CTCCCATCCT GGGCAGGAGT TCGTCAGAA
3571 GTTATGGGAT CTACTGTGGA TGGAAGACCC GTTCAACCCG CCAATTTCTT AACGCTGACC TATGCTACTT
3641 TAAGTTCTTC ACCTTTGGAC GCAGCTGCAG CCGCTGCCGC CGCTCTGTC GCGCTAACA CTGTGCTTGG
3711 AATGGGTTAC TATGGAAGCA TCGTGGCTAA TTCCACTTCC TCTAATAACC CTTCTACACT GACTCAGGAC
3781 AAGTTACTTG TCCTTTTGGC CCAGCTGGAG GCTTTGACCC AACGTCTGGG TGAACCTTCT CAGCAGGTGG
3851 CCGAGTTGCG AGTACAAACT GAGTCTGCTG TCGGCACGGC AAAGTCTAAA TAAAAAAAT TCCAGAATCA
3921 ATGAATAAAT AAACGAGCTT GTTGTGTGAT TAAATCAAG TGTTTTTATT TCATTTTTCG CGCACGGTAT
3991 GCCCTGGACC ACCGATCTCG ATCATTGAGA ACTCGGTGGA TTTTTTCCAG AATCCTATAG AGGTGGGATT
4061 GAATGTTTAG ATACATGGGC ATTAGCCCGT CTTTGGGGTG GAGATAGCTC CATTGAAGGG ATTCATGCTC
4131 CGGGGTAGTG TTGTAATCA CCCAGTCATA ACAAGGTGCG AGTGCATGGT GTTGACAAAT ATCTTTTAGA
4201 AGTAGGCTGA TTGCCACAGA TAAGCCCTTG GTGTAGGTGT TTACAAACCG GTTGAGCTGG GAGGGGTGCA
4271 TTCGAGGTGA AATTATGTGC ATTTTGGATT GGATTTTAA GTTGCAATA TTGCCGCAA GATCCCGTCT

```



Figure 6, contd.

```

4341 TGGGTTTCATG TTATGAAGGA CTACCAAGAC GGTGTATCCG GTACATTTAG GAAATTTATC GTGCAGCTTG
4411 GATGGAAAAG CGTGGAAAAA TTTGGAGACA CCTTGTGTG CTCCGAGATT TTCCATGCAC TCATCCATGA
4481 TAATAGCAAT GGGGCCGTGG GCAGCGGCGC GGGCAAACAC GTTCCGTGGG TCTGACACAT CATAGTTATG
4551 TTCCTGAGTT AAATCATCAT AAGCCATTTT AATGAATTTG GGGCGGAGCG TACCAGATTG GGGTATGAAT
4621 GTTCCTTCGG GCCCCGGAGC ATAGTTCCCC TCACAGATTT GCATTTCCCA AGCTTTCAGT TCTGAGGGTG
4691 GAATCATGTC CACCTGGGGG GCTATGAAGA ACACCGTTTC GGGGGCGGGG GTGATTAGTT GGGATGATAG
4761 CAAGTTTCTG AGCAATTGAG ATTTGCCACA TCCGGTGGGG CCATAAATAA TTCCGATTAC AGGTTGCAGG
4831 TGGTAGTTTA GGAACGGCA ACTGCCGTCT TCTCGAAGCA AGGGGGCCAC CTCGTTTCATC ATTTCCCTTA
4901 CATGCATATT TTCCCGCACC AAATCCATTA GGAGGCGCTC TCCTCCTAGT GATAGAAAGTT CTTGTAGTGA
4971 GGAAAAGTTT TTCAGCGGTT TTAGACCGTC AGCCATGGGC ATTTTGGAAA GAGTTTGCTG CAAAAGTTCT
5041 AGTCTGTTCC ACAGTTCAGT GATGTGTTCT ATGGCATCTC GATCCAGCAG ACCTCCTCGT TTCGCGGGTT
5111 TGGACGGCTC CTGGAGTAGG GTATGAGACG ATGGGCGTCC AGCGCTGCCA GGGTTCGGTC CTTCCAGGGT
5181 CTCAGTGTTT GAGTCAGGGT TGTTTCCGTC ACAGTGAAGG GGTGTGCGCC TGCTTGGGCG CTTGCCAGGG
5251 TCGCGTTTCA ACTCATTCTG CTGGTGGAGA ACTTCTGTCT CTTGGCGCCC TGTATGTCGG CCAAGTAGCA
5321 GTTTACCATG AGTTCGTAGT TGAGCGCCTC GGCTGCGTGG CCTTTGGGCG GGAGCTTACC TTTGGAAGTT
5391 TTCTTGCTA CCGGGCAGTA TAGGCATTTT AGCGCATACA GCTTGGGCGC AAGGAAAATG GATTCTGGGG
5461 AGTATGCATC CGCGCCGCG GAGGCGCAAA CAGTTTCACA TTCCACCAGC CAGGTAAAT CCGGTTTATT
5531 GGGGTCAAAA ACAAGTTTTC CGCCATATTT TTTGATGCGT TTCTTACCTT TGGTCTCCAT AAGTTCGTGT
5601 CCTCGTTGAG TGACAAACAG GCTGTCCGTA TCTCCGTA GA CTGATTTTAC AGGCCTCTTC TCCAGTGGAG
5671 TGCTCGGTC TTCTTCGTAC AGGAACCTCT ACCACTCTGA TACAAAGGCG CGCGTCCAGG CCAGCACAAA
5741 GGAGGCTATG TGGGAGGGGT AGCGATCGTT GTCAACCAGG GGGTCCACCT TTTCCAAAGT ATGCAACAC
5811 ATGTCACCTT CTTCAACATC CAGGAATGTT ATTGGCTTGT AGGTGTATTT CACGTGACCT GGGGTCCCCG
5881 CTGGGGGGGT ATAAAAGGGG GCGGTTCTTT GCTCTCTCT ACTGTCTTCC GGATCGCTGT CCAGGAACGT
5951 CAGCTGTGAG GTAGGTTAT CCTCTCGAA GCGGGGATG ACCTCTGCA TCAAGTTGTC AATTCTTAAG
6021 AACGAGGAGG ATTTGATATT GACAGTGCCG GTTGAGATGC CTTTCATGAG GTTTTCGTCC ATTTGGTCAG
6091 AAAACACAAT TTTTTTATTG TCAAGTTTGG TGGCAAATGA TCCATACAGG GCGTTGGATA AAAGTTTGGC
6161 AATGGATCGC ATGGTTTGGT TCTTTTCTT GTCCGCGCGC TCTTTGGCGG CGATGTTGAG TTGGACATAC
6231 TCGCGTGCCA GGCACCTCCA TTCGGGGAAG ATAGTTGTTA ATTCATCTGG CACGATTCTC ACTTGCCACC
6301 CTCGATTATG CAAGGTAATT AAATCCACAC TGGTGGCCAC CTCGCCTCGA AGGGGTTCAT TGGTCCAACA
6371 GAGCCTACCT CCTTTCTAG AACAGAAAGG GGAAGTGGG TCTAGCATAA GTTCATCGGG AGGGTCTGCA
6441 TCCATGGTAA AGATTTCCCG AAGTAAATCC TTATCAAAAT AGCTGATGGG AGTGGGGTCA TCTAAGGCCA
6511 TTTGCCATTG TCGAGCTGCC AGTGCGCGCT CATATGGGT AAGGGGACTG CCCCAGGGCA TGGGATGGGT
6581 GAGAGCAGAG GCATACATGC CACAGATGTC ATAGACGTAG ATGGGATCCT CAAAGATGCC TATGTAGGTT
6651 GGATAGCATC GCCCCCTCT GATACCTGCT CGCACATAGT CATATAGTTC ATGTGATGGC GCTAGCAGCC
6721 CCGGACCCAA GTTGGTGCGA TTGGGTTTTT CTGTTCTGTA GACGATCTGG CGAAAGATGG CGTGAGAATT
6791 GGAAGAGATG GTGGGTCTTT GAAAAATGTT GAAATGGGCA TGAGGTAGAC CTACAGAGTC TCTGACAAAG
6861 TGGGCATAAG ATTCTTGAAG CTTGGTTACC AGTTCGGCGG TGACAAGTAC GTCTAGGGCG CAGTAGTCAA
6931 GTGTTTCTTG AATGATGTCA TAACCTGGTT GGTTTTCTT TTCCCACAGT TCGCGGTTGA GAAGGTATTC
7001 TTCGCGATCC TTCCAGTACT CTTCTAGCGG AAACCCGTCT TTGTCTGCAC GGTAAAGATCC TAGCATGTAG
7071 AACTGATTAA CTGCTTGTG AGGGCAGCAG CCTTCTCTA CGGGTAGAGA GTATGCTTGA GCAGCTTTTC
7141 GTAGCGAAGC GTGAGTAAGG GCAAAGGTGT CTCTGACCAT GACTTTGAGA AATTGGTATT TGAAGTCCAT
7211 GTCGTCACAG GCTCCCTGTT CCCAGAGTTG GAAGTCTACC CGTTTCTTGT AGGCGGGGTT GGGCAAAGCG
7281 AAAGTAACAT CATTGAAGAG AATCTTACCG GCTCTGGGCA TAAAATTGCG AGTGATGCGG AAAGGCTGTG
7351 GTACTTCCGC TCGATTGTTG ATCACCCTGG CAGCTAGGAC GATTTCTGTC AAACCGTTGA TGTGTGTCTC
7421 TACGATGTAT AATTCTATGA AACGCGGCGT GCCTCTGACG TGAGGTAGCT TACTGAGCTC ATCAAAGGTT
7491 AGGTCTGTGG GGTGAGATAA GCGGTAGTGT TCGAGAGCCC ATTCGTGAG GTGAGGATTT GCATGTAGGA
7561 ATGATGACCA AAGATCTACC GCCAGTGCTG TTTGTAAGT GTCCCGATAC TGACGAAAAT GCCGGCCAAT
7631 TGCCATTTT TCTGGAGTGA CACAGTAGAA GGTTCTGGGG TCTTGTGCG ATCGATCCCA CTTGAGTTTA
7701 ATGGCTAGAT CGTGGGCCAT GTTGACGAGA CGCTCTTCTC CTGAGAGTTT CATGACCAGC ATGAAAGGAA
7771 CTAGTTGTTT GCCAAAGGAT CCCATCCAGG TGTAAGTTTC CACATCGTAG GTCAGGAAGA GTCTTTCTGT
7841 GCGAGGATGA GAGCCGATCG GGAAGAAGT GATTTCTGTC CACCAGTTGG AGGATTGGCT GTTGATGTGA
7911 TGGAAAGTAG AGTTTCTGCG GCGCGCCGAG CATTCTGTGT TGTGCTTGT CAGACGGCCG CAGTAGTCGC
7981 AGCGTTGCAC GGGTTGTATC TCGTGAATGA GCTGTACCTG GCTTCCCTTG ACGAGAAATT TCAGTGGGAA
8051 GCCGAGGCTT GGCGATTGTA TCTCGTGCTC TTCTATATTC GCTGTATCGG CCTGTTTCATC TTCTGTTTCG
8121 ATGGTGGTCA TGCTGACGAG CCCCCGCGGG AGGCAAGTCC AGACCTCGGC GCGGGAGGGG CCGAGCTGAA
8191 GGACGAGAGC GCGCAGGCTG GAGCTGTCCA GAGTCTGAG ACGCTGCGGA CTCAGGTTAG TAGGTAGGGA
8261 CAGAAAGATTA ACTTGATGA TCTTTTCCAG GCGTGTGCGG AGGTTGAGT GGTACTTGAT TTCCACAGGT
8331 TCGTTTGTAG AGACGTCAAT GGCTTGCAAG GTTCCGTGTC CTTTGGGCGC CACTACCGTA CCTTTGTTTT
8401 TTCTTTTGAT CGGTGGTGGC TCTCTTGCTT CTTGATGCT CAGAAGCGGT GACGGGGACG CGCGCCGGGC
8471 GGCAGCGGTT GTTCCGGACC CGGGGGCATG GCTGGTAGTG GCACGTGCGC GCGCGCACG GGCAGGTTCT
8541 GGTATTGCGC TCTGAGAAGA CTTGCGTGCG CCACCACGCG TCGATTGACG TCTGTATCT GACGTCTCTG
8611 GGTGAAAGCT ACCGGCCCCG TGAGCTTGAA CCTGAAAGAG AGTTCAACAG AATCAATTTT GGTATCGTTA
8681 ACGGCAGCTT GTCTCAGTAT TTCTTGATCG TCACCAGAGT TGTCTTGGTA GCGATCTCC GCCATGAAGT
8751 GCTCGATTTT TTCTCTCTGA AGATCTCCGC GACCCGCTCT TTCGACGGTG GCGCGAGGT CATTGGAGAT

```



Figure 6, contd.

```

8821 ACGGCCCATG AGTTGGGAGA ATGCATTCAT GCCCGCCTCG TTCCAGACGC GGCTGTAAAC CACGGCCCCC
8891 TCGGAGTCTC TTGCGCGCAT CACCACCTGA GCGAGGTAA GCTCCACGTG TCTGGTGAAG ACCGCATAGT
8961 TGCATAGGCG CTGAAAAAGG TAGTTGAGTG TGGTGGCAAT GTGTTCGGCG ACGAAGAAAT ACATGATCCA
9031 TCGTCTCAGC GGCATTTTCG TAACATCGCC CAGAGCTTCC AAGCGCTCCA TGGCCTCGTA GAAGTCCACG
9101 GCAAAATTAA AAAACTGGGA GTTTCGCGCG GACACGGTCA ATTCTCTCTC GAGAAGACGG ATGAGTTCGG
9171 CTATGGTGGC CCGTACTTCG CGTTCGAAGG CTCCCGGGAT CTCTTCTTCC TCTTCTATCT CTTCTTCCAC
9241 TAACATCTCT TCTTCGTCTT CAGGCGGGGG CGGAGGGGGC ACGCGGCGAC GTCGACGGCG CACGGGCAAA
9311 CCGTCGATGA ATCGTTCAAT GACCTCTCCG CGGCGGCGGC GCATGGTTTC AGTGACGGCG CGGCCGTTCT
9381 CGCGCGGTGC CAGAGTAAAA ACACCGCCGC GCATCTCTTT AAAGTGGTGA CTGGGAGGTT CTCCGTTTGG
9451 GAGGGAGAGG GCGCTGATTA TACATTTTAT TAATTGGCCC GTAGGGACTG CGCGCAGAGA TCTGATCGTG
9521 TCAAGATCCA CGGATCTGA AAACCTTTTC ACGAAAGCGT CTAACCAAGT ACAGTCACAA GGTAGGCTGA
9591 GTACGGCTTC TTGTGGGCGG GGGTGGTTAT GTGTTCGGTC TGGGTCTTCT GTTTCTTCTT CATCTCGGGA
9661 AGGTGAGACG ATGCTGCTGG TGATGAAATT AAAGTAGGCA GTTCTAAGAC GCGCGATGGT GCGGAGGAGC
9731 ACCAGGTCTT TGGGTCCGGC TTGCTGGATA CGCAGGCGAT TGGCCATTCC CCAAGCATTG TCCTGACATC
9801 TAGCAAGATC TTTGTAGTAG TCTTGCATGA GCCGTTCTAC GGGCACTTCT TCCTCACCCTG TTCTGCCATG
9871 CATACGTGTG AGTCCAAATC CGCGCATTTG TTGTACCAGT GCCAAGTCAG CTACGACTCT TTCGGCGAGG
9941 ATGGCTTGCT GTACTTGGGT AAGGGTGGCT TGAAAGTCAT CAAAATCCAC AAAGCGGTGG TAAGCCCCTG
10011 TATTAATGGT GTAAGCACAG TTGGCCATGA CTGACCAGTT AACTGTCTGG TGACCAGGGC GCACGAGCTC
10081 GGTGTATTTA AGGCGCGAAT AGGCGCGGGT GTCAAAGATG TAATCGTTGC AGGTGCGCAC CAGATACTGG
10151 TACCCTATAA GAAAATGCGG CCGTGGTTGG CCGTAGAGAG GCCATCGTTC TGTAGCTGGA GCGCCAGGGG
10221 CGAGGTCTTC CAACATAAGG CCGTGATAGC CGTAGATGTA CCTGGACATC CAGGTGATTG CTGCGGCGGT
10291 AGTAGAAGCC CGAGGAAACT CGCGTACGCG GTTCCAAATG TTGCGTAGCG GCATGAAGTA GTTCATTGTA
10361 GGCACGGTTT GACCAGTGAG GCGCGCGCAG TCATTGATGC TCTATAGACA CGGAGAAAAT GAAAGCGTTC
10431 AGCGACTCGA CTCCGTAGCC TGGAGGAAAG TAAAGGGTTC GGTTCGCGGT GTACCCCGGT TCGAGACTTG
10501 TACTCGAGCC GGCCGGAGCC GCGGCTAACG TGGTATTGGC ACTCCCGTCT CGACCCAGCC TACAAAAATC
10571 CAGGATACGG AATCGAGTCG TTTTGCTGGT TTCCGAATGG CAGGGAAGTG AGTCCTATTT TTTTTTTTTT
10641 TTTGCCGCTC AGATGCATCC CGTGCTGCGA CAGATGCGCC CCCAACAACA GCCCCCTCG CAGCAGCAGC
10711 AGCAGCAACC ACAAAGGCT GTCCCTGCAA CTACTGCAAC TGCCGCGGTG AGCGGTGCGG GACAGCCCGC
10781 CTATGATCTG GACTTGGAAG AGGGCGAAGG ACTGGCACGT CTAGGTGCGC CTTGCCCCGA GCGGCATCCG
10851 CGAGTTCAAC TGAAAAAAGA TTCTCGCGAG GCGTATGTGC CCCAACAGAA CCTATTTAGA GACAGAAGCG
10921 GGAGGAGCC GGAGGAGATG CGAGCTTCCC GCTTTAAGCG CTGCGTGAG CTGCGTCACG GTTTGGACCG
10991 AAGACGAGTG TTGCGAGACG AGGATTTCGA AGTTGATGAA GTGACAGGGA TCAGTCCCTG CAGGGCACAC
11061 GTGGCTGCAG CCAACCTTGT ATCGGCTTAC GAGCAGACAG TAAAGGAAGA GCGTAACCTC CAAAAGTCTT
11131 TTAATAATCA TGTGCGAACC CTGATTGCCC GCGAAGAAGT TACCCTTGGT TTGATGCATT TGTGGGATTT
11201 GATGGAAGCT ATCATTGAGA ACCCTACTAG CAAACCTCTG ACCGCCAGC TGTTTCTGGT GGTGCAACAC
11271 AGCAGAGACA ATGAGGCTTT CAGAGAGGCG CTGCTGAACA TCACCGAACC CGAGGGGAGA TGGTTGTATG
11341 ATCTTATCAA CATTCTACAG AGTATCATAG TGCAGGAGCG GAGCCTGGGC CTGGCCGAGA AGGTAGCTGC
11411 CATCAATTAC TCGGTTTTGA GCTTGGGAAA ATATTACGCT CGCAAAATCT ACAAGACTCC ATACGTTCCC
11481 ATAGACAAGG AGGTGAAGAT AGATGGGTTT TACATGCGCA TGACGCTCAA GGTCTTGACC CTGAGCGATG
11551 ATCTGGGGT GTATCGCAAT GACAGAATGC ATCGCGCGT TAGCGCCAGC AGGAGGCGCG AGTTAAGCGA
11621 CAGGGAAGTG ATGCACAGTT TGCAAGAGAG TCTGACTGGA GCTGGAACCG AGGGTGAGAA TTACTTCGAC
11691 ATGGGAGCTG ACTTGCAAGT GCAGCCTAGT CGCAGGGCTC TGAGCGCCGC GACGGCAGGA TGTGAGCTTC
11761 CTTACATAGA AGAGGCGGAT GAAGGCGAGG AGGAAGAGGG CGAGTACTTG GAAGACTGAT GGCACAACCC
11831 GTGTTTTTTG CTAGATGGAA CAGCAAGCAC CGGATCCCGC AATGCGGGCG GCGCTGCAGA GCCAGCCGTC
11901 CGGCATTAACT CCTCGGACG ATTGGACCCA GGCCATGCAA CGTATCATGG CGTTGACGAC TCGCAACCCC
11971 GAAGCCTTTA GACAGCAACC CCAGGCCAAC CGTCTATCGG CCATCATGGA AGCTGTAGTG CCTTCCCGAT
12041 CTAATCCAC TCATGAGAAG GTCCTGGCCA TCGTGAACGC GTTGGTGGAG AACAAAGCTA TTCGTCCAGA
12111 TGAGGCCGGA CTGGTATACA ACGCTCTCTT AGAACGCGTG GCTCGCTACA ACAGTAGCAA TGTGCAAAAC
12181 AATTTGGACC GTATGATAAC AGATGTACGC GAAGCCGTGT CTCAGCGCGA AAGGTTCAG CGTGATGCCA
12251 ACCTGGGTT GCTGGTGGCG TTAATGCTT TCTTGAGTAC TCAGCCTGCT AATGTGCCG GTGGTCAACA
12321 GGATTATACT AACTTTTTAA GTGCTTTGAG ACTGATGGTA TCAGAAGTAC CTCAGAGCGA AGTGTATCAG
12391 TCCGGTCTTG ATTACTTCTT TCAGACTAGC AGACAGGGCT TGCAGACGGT AAATCTGAGC CAAGCTTTTA
12461 AAAACCTTAA AGGTTTGTGG GGAGTGCATG CCCCGGTAGG AGAAAGAGCA ACCGTGTCTA GCTTGTTAAC
12531 TCCGAAGTCC CGCCTGTTAT TACTGTGGT AGCTCCTTTC ACCGACAGCG GTAGCATCGA CCGTAATTCC
12601 TATTTGGGTT ACCTACTAAA CCTGTATCGC GAAGCCATAG GGCAAGTCA GGTGGACGAG CAGACCTATC
12671 AAGAAATTAC CCAAGTCAGT CGCGCTTTGG GACAGGAAGA CACTGGCAGT TTGGAAGCCA CTCTGAACCT
12741 CTTGCTTACC AATCGGTCTC AAAAGATCCC TCCTCAATAT GCTCTTACTG CGGAGGAGGA GAGGATCCTT
12811 AGATATGTGC AGCAGAGCGT GGGATTGTTT CTGATGCAAG AGGGGGCAAC TCCGACTGCA GCACTGGACA
12881 TGACAGCGCG AAATATGGAG CCCAGCATGT ATGCCAGTAA CCGACCTTTC ATTAACAAAC TGCTGGACTA
12951 CTTGCACAGA GCTGCCGCTA TGAATCTGAC TTATTTTACC AATGCCATCT TAAACCCGCA CTGGCTGCCG
13021 CCACCTGGTT TCTACACGGG CGAATATGAC ATGCCGACC CTAATGACCG ATTTCTGTGG GACGACGTGG
13091 ACAGCGATGT TTTTTCACCT CTTTCTGATC ATCGCACGTG GAAAAAGGAA GGCGGTGATA GAATGCATTG
13161 TTCTGCATCG CTGTCCGGGG TCATGGGTGC TACCGCGGCT GAGCCCGAGT CTGCAAGTCC TTTTCTAGT
13231 CTACCCCTTT CTCTACACAG TGTACGTAGC AGCGAAGTGG GTAGAATAAG TCGCCCGAGT TTAATGGGCG

```

Figure 6, contd.

```

13301 AAGAGGAGTA CCTAAACGAT TCCTTGCTCA GACCGGCAAG AGAAAAAAT TTCCCAAACA ATGGAATAGA
13371 AAGTTTGGTG GATAAAATGA GTAGATGGAA GACTTATGCT CAGGATCACA GAGACGAGCC TGGGATCATG
13441 GGGACTACAA GTAGAGCGAG CCGTAGACGC CAGCGCCATG ACAGACAGAG GGGTCTTGTG TGGGACGATG
13511 AGGATTCCGC CGATGATAGC AGCGTGTTGG ACTTGGGTGG GAGAGGAAGG GGCAACCCGT TTGCTCATTT
13581 GCGCCCTCGC TTGGGTGGTA TGTGTGAAA AAAAATAAAA AAGAAAAACT CACCAAGGCC ATGGCGACGA
13651 GCGTACGTTT GTTCTTCTTT ATTATCTGTG TCTAGTATAA TGAGGCGAGT CGTGCTAGGC GGAGCGGTGG
13721 TGTATCCGGA GGGTCCTCCT CCTTCGTACG AGAGCGTGAT GCAGCAGCAG CAGGCGACGG CGGTGATGCA
13791 ATCCCCACTG GAGGCTCCCT TTGTGCCTCC GCGATACCTG GCACCTACGG AGGGCAGAAA CAGCATTCTG
13861 TACTCGGAAC TGGCACCTCA GTACGATACC ACCAGGTTGT ATCTGGTGGA CAACAAGTCG GCGGACATTG
13931 CTTCTCTGAA CTATCAGAAT GACCACAGCA ACTTCTTGAC CACGGTGGTG CAGAACAATG ACTTTACCCC
14001 TACGGAAGCC AGCACCCAGA CCATTAACCT TGATGAACGA TCGCGGTGGG GCGGTACGCT AAAGACCATC
14071 ATGCATACTA ACATGCCAAA CGTGAACGAG TATATGTTTA GTAACAAGTT CAAAGCGCGT GTGATGGTGT
14141 CCAGAAAACC TCCCGACGGT GCTGCAGTTG GGGATACTTA TGATCACAAG CAGGATATTT TGGAAATATG
14211 GTGGTTCCGAG TTTACTTTGC CAGAAGGCAA CTTTTAGTT ACTATGACTA TTGATTTGAT GAACAATGCC
14281 ATCATAGATA ATTACTTGAA AGTGGGTAGA CAGAATGGAG TGCTTGAAAG TGACATTGGT GTTAAGTTCC
14351 ACACCAGGAA CTTCAAGCTG GGATGGGATC CCGAAACCAA GTTGATCATG CCTGGAGTGT ATACGTATGA
14421 AGCCTTCCAT CCTGACATTG TCTTACTGCC TGGCTGCGGA GTGGATTTTA CCGAGAGTCG TTTGAGCAAC
14491 CTTCTTGGTA TCAGAAAAAA ACAGCCATTT CAAGAGGGTT TTAAGATTTT GTATGAAGAT TTAGAAGGTG
14561 GTAATATTCC GGCCCTCTTG GATGTAGATG CCTATGAGAA CAGTAAGAAA GAACAAAAAG CCAAAATAGA
14631 AGCTGCTACA GCTGCTGCAG AAGCTAAGG AAACATAGTT GCCAGCGACT CTACAAGGGT TGCTAACGCT
14701 GGAGAGGTCA GAGGAGACAA TTTGCGCCA ACACCTGTTT CGACTGCAGA ATCATTATTG CCCGATGTGT
14771 CTGAAGGAAC GGACGTGAAA CTCACTATTC AACCTGTAGA AAAAGATAGT AAGAATAGAA GCTATAATGT
14841 GTTGAAGAC AAAATCAACA CAGCCTATCG CAGTTGGTAT CTTTCGTACA ATTATGGCGA TCCCGAAAAA
14911 GAGGTGCGT CCTGACAT TCTACCCAC TTAGATGTCA CTTGCGGAG AGAGCGGTT TACTGGTCCG
14981 TTCCAGACAT GATGAAGGAT CCTGTCACTT TCCGCTCCAC TAGACAAGTC AGTAACTACC CTGTGGTGGG
15051 TGCAGAGCTT ATGCCCGTCT TCTCAAAGAG CTTCTACAAC GAACAAGCTG TGTACTCCCA GCAGCTCCGC
15121 CAGTCCACCT CGCTTACGCA CGTCTTCAAC CGCTTTCTTG AGAACCAGAT TTTAATCCGT CCGCCGGCGC
15191 CCACCATTAC CACCGTCAGT GAAAACGTTT CTGCTCTCAC AGATCACGGG ACCCTGCCGT TGCBCAGCAG
15261 TATCCGGGGA GTCCAACGTG TGACCGTTAC TGACGCCAGA CGCCGCACCT GTCCCTACGT GTACAAGGCA
15331 CTGGGCATAG TCGCACCGCG CGTCCTTTCA AGCCGCACTT TCTAAAAAAA AAAAATGTCC ATTCTTATCT
15401 CGCCAGTAA TAACACCGGT TGGGGTCTGC GCGCTCCAAG CAAGATGTAC GGAGGCGCAC GCAAACGTTT
15471 TACCCAACAT CCCGTGCGTG TTCGCGGACA TTTTCGCGCT CCATGGGGTG CCCTCAAGGG CCGCACTCCG
15541 GTTCGAACCA CCGTCGATGA TGTAATCGAT CAGGTGGTTG CCGACGCCCC TAATTATACT CCTACTGCGC
15611 CTACATCTAC TGTGGATGCA GTTATTGACA GTGTAGTGGC TGACGCTCGC AACTATGCTC GACGTAAGAG
15681 CCGGCGAAGG CGCATTGCCA GACGCCACCG AGCTACCCTT GCCATGCGAG CCGCAAGAGC TCTGCTACGA
15751 AGAGCTAGAC GCGTGGGGCG AAGAGCCATG CTTAGGGCGG CCAGACGTGC AGCTTCGGGC GCCAGCGCCG
15821 GCAGGTCCCG CAGGCAAGCA GCGGCTGTCC CAGCGGCGAC TATTGCCGAC ATGGCCCAAT CCGGAAGAGG
15891 CAATGTATAC TGGGTGCGTG ACGCTGCCAC CCGTCAACGT GTACCCGTGC GCACCCGTCC CCTCGCACT
15961 TAGAAGATAC TGAGCAGTCT CCGATGTTGT GTCCCGCGG CGAGGATGTC CAAGCGCAA TACAAGGAAG
16031 AAATGCTGCA GGTATCGCA CCTGAAGTCT ACGGCCAACG GTTGAAGGAT GAAAAAAAC CCCGCAAAT
16101 CAAGCGGGTT AAAAAGGACA AAAAAGAAGA GGAAGATGGC GATGATGGG TGGCGGAGTT TGTGCCGAG
16171 TTTGCCCCAC GCGCAGCGT GCAATGGCGT GGGCGCAAAG TTCGACATGT GTTGAGACCT GGAACCTCGG
16241 TGGTCTTTAC ACCCGCGAG CCGTCAAGCG CTACTTTTAA GCGTTCTTAT GATGAGGTGT ACGGGGATGA
16311 TGATATTCTT GAGCAGGCGG CTGACCGATT AGGCGAGTTT GCTTATGGCA AGCGTAGTAG AATAACTTCC
16381 AAGGATGAGA CAGTGTCAT ACCCTTGGAT CATGGAAATC CCACCCCTAG TCTTAAACCG GTCACTTTGC
16451 AGCAAGTGTT ACCCGTAACT CCGCGAACAG GTGTTAAACG CGAAGGTGAA GATTTGTATC CCACTATGCA
16521 ACTGATGGTA CCCAAACGCC AGAAGTTGGA GGACGTTTTG GAGAAAGTAA AAGTGGATCC AGATATTCAA
16591 CCTGAGGTTA AAGTGAGACC CATTAAAGCAG GTAGCGCCTG GTCTGGGGGT ACAAACGTGA GACATTAAGA
16661 TTCCCACTGA AAGTATGGAA GTGCAAACCTG AACCCGCAAA GCCTACTGCC ACCTCCACTG AAGTGCAAAC
16731 GGATCCATGG ATGCCCATGC CTATTACAAC TGACGCCGCC GGTCCCACTC GAAGATCCCG ACGAAAGTAC
16801 GGTCCAGCAA GTCTGTTGAT GCCCAATTAT GTTGACACC CATCTATTAT TCCTACTCCT GGTACCAGG
16871 GCACTCGCTA CTATCGCAGC CGAAACAGTA CCTCCCGCCG TCGCCGCAAG ACACCTGCAA ATCGCAGTCG
16941 TCGCCGTAGA CGCACAAAGCA AACCGACTCC CGCGCCCTTG GTGCGGCAAG TGTACCGCAA TGGTAGTGCG
17011 GAACCTTTGA CACTGCCGCG GTGCGGTTAC CATCCGAGTA TCATCACTTA ATCAATGTTG CCGCTGCCTC
17081 CTTGCAGATA TGGCCCTCAC TTGTGCGCTT CGCGTTCCCA TCACTGGTTA CCGAGGAAGA AACTCGCGCC
17151 GTAGAAGAGG GATGTTGGGA CGCGGAATGC GACGCTACAG GCGACGGCGT GCTATCCGCA AGCAATTGCG
17221 GGGTGGTTTT TTACCAGCCT TAATTCCAAT TATCGCTGCT GCAATTGGCG CGATACCAGG CATAGCTTCC
17291 GTGGCGGTTT AGGCCTCGCA ACGACATTGA CATTGGAAAA AAAACGTATA AATAAAAAAA AATACAATGG
17361 ACTCTGACAC TCCTGGTCCT GTGACTATGT TTTCTTAGAG ATGGAAGACA TCAATTTTTT ATCCTTGGCT
17431 CCGCGACACG GCACGAAGCC GTACATGGGC ACCTGGAGCG ACATCGGCAC GAGCCAACCT AACGGGGGCG
17501 CCTTCAATTG GAGCAGTATC TGGAGCGGGC TTAATAATTT TGGCTCAACC ATAAAAACAT ACGGGAACAA
17571 AGCTTGGAAC AGCAGTACAG GACAGGCGCT TAGAAATAAA CTTAAAGACC AGAACTTCCA AAAAAAGTA
17641 GTCGATGGGA TAGCTTCCGG CATCAATGGA GTGGTAGATT TGGCTAACCA GGCTGTGCAG AAAAAGATAA
17711 ACAGTCGTTT GGACCCGCCG CCAGCAACCC CAGGTGAAAT GCAAGTGGAG GAAGAAATTC CTCCGCCAGA

```

Figure 6, contd.

```

17781 AAAACGAGGC GACAAGCGTC CGCGTCCCGA TTTGGAAGAG ACGCTGGTGA CGCGCGTAGA TGAACCGCCT
17851 TCTTATGAGG AAGCAACGAA GCTTGGGAATG CCCACCACTA GACCGATAGC CCCAATGGCC ACCGGGGTGA
17921 TGAAACCTTC TCAGTTGCAT CGACCCGTCA CCTTGGATTT GCCCCTCCC CCTGCTGCTA CTGCTGTACC
17991 CGCTTCTAAG CCTGTGCTG CCCCAGAAC AGTCGCCGTA GCCAGGTCAC GTCCCGGGGG CGCTCCTCGT
18061 CCAAATGCGC ACTGGCAAAA TACTCTGAAC AGCATCGTGG GTCTAGGCGT GCAAAGTGTA AAACGCCGTC
18131 GCTGCTTTTA ATTAATATG GAGTAGCGCT TAACTTGCCT ATCTGTGTAT ATGTGTCATT ACACGCCGTC
18201 ACAGCAGCAG AGGAAAAAAG GAAGAGGTG TGCCTGACG CTGAGTTACT TTCAAGATGG CCACCCCATC
18271 GATGCTGCCC CAATGGGCAT ACATGCACAT CGCCGGACAG GATGCTTCGG AGTACCTGAG TCCGGGTCTG
18341 GTGCAGTTCG CCGCGCCAC AGACACCTAC TTCAATCTGG GAAATAAGTT TAGAAATCCC ACCGTAGCGC
18411 CGACCCACGA TGTGACCACC GACCGTAGCC AGCGGCTCAT GTTGCCTTC GTGCCCCTTG ACCGGGAGGA
18481 CAATACATAC TCTTACAAAG TCGGTACAC CCTGGCCGTG GCGACAACA GAGTGTGGA TATGGCCAGC
18551 ACGTTCTTTG ACATTAGGGG CGTGTGGAC AGAGGTCCCA GTTTCAAACC CTATTCTGGT ACGGCTTACA
18621 ACTCTCTGGC TCCTAAAGGC GCTCCAAATG CATCTCAATG GATTGCAAAA GCGGTACCAA CTGCAGCAGC
18691 CGCAGGCAAT GGTGAAGAAG AACATGAAAC AGAGGAGAAA ACTGCTACTT ACACCTTTGC CAATGCTCCT
18761 GTAAAAGCCG AGGCTCAAAT TACAAAAGAG GGCTTACCAA TAGGTTTGGG GATTTGAGCT GAAAACGAAT
18831 CTAAACCCAT CTATGCAGAT AAACTTTATC AGCCAGAACC TCAAGTGGGA GATGAAACTT GGACTGACCT
18901 AGACGGAAAA ACCGAAGAGT ATGGAGGCAG GGCTCTAAAG CCTACTACTA ACATGAAACC CTGTTACGGG
18971 TCCTATGCGA AGCCTACTAA TTTAAAAGGT GGTGAGGCAA AACCAGAAAA CTCGGAACCG TCGAGTGAAA
19041 AAATTGAATA TGATATTGAC ATGGAATTTT TTGATAACTC ATCGCAAAGA ACAAACCTCA GTCCTAAAT
19111 TGTATGTAT GCAGAAAATG TAGGTTTGGG AACGCCAGAC ACTCATGTAG TGTACAAACC TGGAACAGAA
19181 GACACAAGTT CCGAAGCTAA TTTGGGACAA CAGTCTATGC CCAACAGACC CAACTACATT GGCTTCAGAG
19251 ATAACCTTAT TGGACTCATG TACTATAACA GTACTGGTAA CATGGGGGTG CTGGCTGGTC AAGCGTCTCA
19321 GTTAAATGCA GTGGTTGACT TGCAGGACAG AAACACAGAA CTTTCTTACC AACTCTTGCT TGACTCTCTG
19391 GCGACAGAA CAGATACTT TAGCTGTGG AATCAGGCTG TGGACGTTA TGATCTGAT GATCTGTAT
19461 TTGAAAATCA TGGTGTGGAA GATGAACCTC CCAACTATTG TTTTCCACTG GACGGCATAG GTGTTCCAAC
19531 AACCAGTTAC AAATCAATAG TTCCAAATGG AGAAGATAAT AATAATTGGA AAGAACCTGA AGTAAATGGA
19601 ACAAGTGAGA TCGGACAGGG TAATTTGTTT GCCATGGAAA TTAACCTTCA AGCCAATCTA TGGCGAAGTT
19671 TCCTTTATTC CAATGTGGCT CTGTATCTCC CAGACTCGTA CAAATACACC CCGTCCAATG TCACTCTTCC
19741 AGAAAACAAA AACACCTACG ACTACATGAA CCGGCGGGTG GTGCCGCCAT CTCTAGTAGA CACCTATGTG
19811 AACATTGGTG CCAGGTGGTC TCTGGATGCC ATGGACAATG TCAACCCATT CAACCCACCAC CGTAACGCTG
19881 GCTTGCCTTA CCGATCTATG CTTCTGGGTA ACGGACGTTA TGTGCTTTT CACATACAAG TGCCTCAAAA
19951 ATTCTTCGCT GTTAAAAACC TGCTGCTTCT CCCAGGCTCC TACACTTATG AGTGGAACTT TAGGAAGGAT
20021 GTGAACATGG TTCTACAGAG TTCCCTCGGT AACGACCTGC GGGTAGATGG CGCCAGCATC AGTTTCACGA
20091 GCATCAACCT CTATGCTACT TTTTCCCA TGCTCACAA CACCGCTTCC ACCCTGAAG CCATGCTGCG
20161 GAATGACACC AATGATCAGT CATTCAACGA CTACCTATCT GCAGCTAACA TGCTCTACCC CATTCTGCC
20231 AATGCAACCA ATATTCCCAT TTCCATTCTT TCTCGCAACT GGGCGGCTTT CAGAGGCTGG TCATTTACCA
20301 GACTGAAAAC CAAAGAAACT CCCTCTTTGG GGTCTGGATT TGACCCCTAC TTTGTCTATT CTGGTTCTAT
20371 TCCCTACCTG GATGGTACCT TCTACCTGAA CCACACTTTT AAGAAGGTTT CCATCATGTT TGACTCTTCA
20441 GTGAGCTGGC CTGGAATGA CAGGTTACTA TCTCCTAACG AATTTGAAAT AAAGCGCACT GTGGATGGCG
20511 AAGGCTACAA CGTAGCCCAA TGCAACATGA CCAAAGACTG GTTCTTGGTA CAGATGCTCG CCAACTACAA
20581 CATCGCTAT CAGGCTTCT ACATCCAGA AGGATACAAA GATCGCATGT ATTCAATTTT CAGAACTTC
20651 CAGCCCATGA GCAGGCAGGT GGTGATGAG GTCAATTACA AAGACTTCAA GGCCGTCGCC ATACCCTACC
20721 AACACAACAA CTCTGGCTTT GTGGGTTACA TGGCTCCGAC CATGCGCCAA GGTCAACCTT ATCCCGCTAA
20791 CTATCCCTAT CCACTCATTG GAACAACGTC CGTAAATAGT GTTACGCAGA AAAAGTTCTT GTGTGACAGA
20861 ACCATGTGGC GCATACCGTT CTCGAGCAAC TTCATGTCTA TGGGGGCCCT TACAGACTTG GGACAGAATA
20931 TGCTCTATGC CAACTCAGCT CATGCTCTGG ACATGACCTT TGAGGTGGAT CCCATGGATG AGCCACCCCT
21001 GCTTTATCTT CTCTTCGAAG TTTTCGACGT GGTGAGAGTG CATCAGCCAC ACCGCGGCAT CATCGAGGCA
21071 GTCTACCTGC GTACACCGTT CTCGGCCGGT AACGCTACCA CGTAAGAAGC TTCTTGCTTC TTGCAATAG
21141 CAGCTGCAAC CATGGCCTGC GGATCCCAA ACGGCTCCAG CGAGCAAGAG CTCAGAGCCA TTGTCCAAGA
21211 CCTGGGTTGC GGACCTATT TTTTGGGAAC CTACGATAAG CGCTTCCCGG GGTTTATGGC CCCCAGTAAG
21281 CTCGCTGTG CCATTGTAAA TACGGCCGGA CGTGAGACGG GGGGAGAGCA CTGGTTGGCT TTCGGTTGGA
21351 ACCACGTTT TAACACCTGC TACCTTTTGG ATCTTTTGG ATTCTCGGAT GATCGTCTCA AACAGATTTA
21421 CCAGTTTGAA TATGAGGGTC TCCTGCGCCG CAGCGCTCTT GCTACCAAGG ACCGCTGTAT TACGCTGGAA
21491 AAATCTACCC AGACCGTGCA GGGCCCCCGT TCTGCCGCTT GCGGACTTTT CTGCTGCATG TTCTTACAG
21561 CCTTTGTGCA CTGGCCTGAC CGTCCCATGG ACGGAAACCC CACCATGAAA TTGCTAACTG GAGTGCCAAA
21631 CAACATGCTT CATTCTCTA AAGTCCAGCC CACCCTGTGT GACAATCAAA AAGCACTCTA CCATTTTCTT
21701 AATACCCATT CGCCTTATT TCGCTCTCAT CGTACACACA TCGAAAGGGC CACTGCGTTC GACCGTATGG
21771 ATGTTCAATA ATGACTCATG TAAACAACGT GTTCAATAAA CATCACTTTA TTTTTTACA TGTATCAAGG
21841 CTCTGGATTA CTTATTTATT TACAAGTCGA ATGGGTTCTG ACGAGAATCA GAATGACCCG CAGGCAGTGA
21911 TACGTTGCGG AACTGATACT TGGGTTGCCA CTTGAATTGG GGAATCACCA ACTTGGGAAC CGGTATATCG
21981 GGCAGGATGT CACTCCACAG CTTTCTGGTC AGCTGCAAAG GTCCAAGCAG GTCAGGAGCC GAAATCTTG
22051 AATCACAATT AGGACCAAGT CTCTGAGCGC GAGAGTTGCG GTACACCGGA TTGCAGCACT GAAACACCAT
22121 CAGCGACGGA TGTCTACGC TTGCCAGCAC GGTGGGATCT GCAATCATGC CCACATCCAG ATCTTCAGCA
22191 TTGGCAATGC TGAACGGGGT CATCTTGCAG GTCTGCCCTAC CCATGGCGGG CACCCAATTA GGCTTGTTGGT

```



Figure 6, contd.

22261 TGCAATCGCA GTGCAGGGGG ATCAGTATCA TCTTGGCCTG ATCCTGTCTG ATTCCTGGAT ACACGGCTCT  
 22331 CATGAAAGCA TCATATTGCT TGAAAGCCTG CTGGGCTTTA CTACCCCTCG TATAAAACAT CCCGCAGGAC  
 22401 CTGCTCGAAA ACTGGTTAGC TGCACAGCCG GCATCATTCA CACAGCAGCG GCGGTCATTG TTGGCTATTT  
 22471 GCACCACACT TCTGCCCCAG CCGTTTTGGG TGATTTTGGT TCGCTCGGGA TTCTCCTTTA AGGCTCGTTG  
 22541 TCCGTTCTCG CTGGCCACAT CCATCTCGAT AATCTGCTCC TTCTGAATCA TAATATTGCC ATGCAGGCAC  
 22611 TTCAGCTTGC CCTCATAATC ATTGCAGCCA TGAGGCCACA ACGCACAGCC TGTACATTCC CAATTATGGT  
 22681 GGGCGATCTG AGAAAAAGAA TGTATCATTG CCTGCAGAAA TCTTCCCATC ATCGTGCTCA GTGTCTTGTG  
 22751 ACTAGTGAAA GTTAACTGGA TGCCCTGGTG CTCTTCGTTT ACGTACTGGT GACAGATGCG CTTGTATTGT  
 22821 TCGTGTGTCT CAGGCATTAG TTTAAAAACAG GTTCTAAGTT CGTTATCCAG CCTGTACTTC TCCATCAGCA  
 22891 GACACATCAC TTCCATGCCT TTCTCCCAAG CAGACACCAG GGGCAAGCTA ATCGGATTCT TAACAGTGCA  
 22961 GGCAGCAGCT CCTTTAGCCA GAGGGTCATC TTTAGCGATC TTCTCAATGC TTCTTTTGCC ATCCTTCTCA  
 23031 ACGATGCGCA CGGGCGGGTA GCTGAAACCC ACTGCTACAA GTTGCGCCTC TTCTCTTTCT TCTTCGCTGT  
 23101 CTTGACTGAT GTCTTGCAAT GGGATATGTT TGGTCTTCTT TGGCTTCTTT TTGGGGGGTA TCGGAGGAGG  
 23171 AGGACTGTCG CTCGTTCCG GAGACAGGGA GGATTGTGAC GTTTCTGCTA CCATTACCAA CTGACTGTCTG  
 23241 GTAGAAGAAC CTGACCCAC ACGGCGACAG GTGTTTTTCT TCGGGGGCAG AGGTGGAGGC GATTGCGAAG  
 23311 GGCTGCGGTC CGACCTGGAA GCGGATGAC TGGCAGAACC CCTTCCGCGT TCGGGGGTGT GCTCCCTGTG  
 23381 GCGGTCGCTT AACTGATTTC CTTCGCGGCT GGCCATTGTG TTCTCCTAGG CAGAGAAACA ACAGACATGG  
 23451 AAACCTCAGC ATTGCTGTCA ACATCGCCAC GAGTGCCATC ACATCTCGTC CTCAGCGACG AGGAAAAGGA  
 23521 GCAGAGCTTA AGCATTCCAC CGCCAGTCC TGCCACCACC TCTACCCTAG AAGATAAGGA GGTGACGCA  
 23591 TCTCATGACA TGCAGAATAA AAAAGCGAAA GAGTCTGAGA CAGACATCGA GCAAGACCCG GGCTATGTGA  
 23661 CACCGGTGGA ACACGAGGAA GAGTTGAAAC GCTTTCTAGA GAGAGAGGAT GAAAACTGCC CAAAACAGCG  
 23731 AGCAGATAAC TATCACCAG ATGCTGGAAA TAGGGATCAG AACACCGACT ACCTCATAGG GCTTGACGGG  
 23801 GAAGACGCGC TCCTTAAACA TCTAGCAAGA CAGTCGCTCA TAGTCAAGGA TGCATTATTG GACAGAACTG  
 23941 CAAACGTCAG CCAAACGGCA CCTGCGAGCC AAATCCTCGC TTAAACTTTT ATCCAGCTTT TGCTGTGCCA  
 24011 GAAGTACTGG CTACCTATCA CATCTTTTTT AAAAATCAAA AAATTCCAGT CTCTGCGCGC GCTAATCGCA  
 24081 CCCGCGCCGA TGCCCTACTC AATCTGGGAC CTGGTTACAG CTTACCTGAT ATAGCTTCTT TGGAGAGGT  
 24151 TCCAAAGATC TTCCAGGGTC TGGGCAATAA TGAGACTCGG GCCGCAATG CTCTGCAAAA GGGAGAAAAT  
 24221 GGCATGGATG AGCATCACAG CGTCTGGTG GAATTGGAAG GCGATAATGC CAGACTCGCA GTACTCAAGC  
 24291 GAAGCGTCGA GGTACACAC TTGCGATATC CCGCTGTCAA CCTGCCCCCT AAAGTCATGA CCGCGGTCAT  
 24361 GGACCAGTTA CTCATTAAGC GCGCAAGTCC CCTTTCAGAA GACATGCATG ACCCAGATGC CTGTGATGAG  
 24431 GGTAACCCAG TGGTCAGTGA TGAGCAGCTA ACCCGATGGC TGGGCACCGA CTCTCCCCGG GATTGGAAG  
 24501 AGCGTCGCAA GCTTATGATG GCCGTGGTGC TGGTTACCGT AGAAGTAGAG TGTCTCCGAC GTTTCTTTAC  
 24571 GCATTGAGAA ACCTTGCGCA AACTCGAAGA BAATCTGCAC TACACTTTTA GACACGGCTT TGTGCGGCAG  
 24641 GCATGCAAGA TATCTAACGT GGAATCACC AAACCTGGTT CCTACATGGG TATTCTGCAT GAGAATCGCC  
 24711 TAGGACAAAG CGTGTGCAC AGCACCTTA AGGGGGAAGC CCGCCGTGAT TACATCCGCG ATTGTGTCTA  
 24781 TCTCTACCTG TGCCACACGT GGCAAAACCG CATGGGTGTA TGGCAGCAAT GTTTAGAAGA ACAGAACTTG  
 24851 AAAGAGCTTG ACAAGCTCTT ACAGAAATCT CTTAAGGTTT TGTGGACAGG GTTCGACGAG CGCACCGTCG  
 24921 CTTCCGACCT GGCAGACCTC ATCTTCCAG AGCGTCTCAG GGTACTTTTG CGAAACGGAT TGCCGTGACTT  
 24991 TATGAGCCAG AGCATGCTTA ACAATTTTCG CTCTTTTCATC CTGGAACGCT CCGGTATCCT GCCCGCCACC  
 25061 TGCTGCGCAC TGCCCTCCGA CTTTGTGCC TCTACCTACC GCGAGTGCCC CCCGCCGCTA TGGAGTCACT  
 25131 GCTACCTGTT CCGTCTGGCC AACTATCTCT CCTACCACTC GGATGTGATC GAGGATGTGA GCGGAGACGG  
 25201 CTTGCTGGAG TGCCACTGCC GCTGCAATCT GTGCACGCC CACCGGTCCC TAGCTTGCAA CCCCCAGTTG  
 25271 ATGAGCGAAA CCCAGATAAT AGGCACCTTT GGTGACCAAG GCCCCAGCAG CCAAGGCGAT GGGTCTTCTC  
 25341 CTGGGCAAAG TTTAAACTG ACCCCGGGAC TGTGGACCTC CGCTACTTGG CGCAAGTTTG CTCCGGAAGA  
 25411 TTACCACCCC TATGAAATCA AGTTCTATGA GGACCAATCA CAGCCTCCAA AGGCCGAAC TTTGGCTTGC  
 25481 GTCATACCCC AGGGGGCAAT TCTGGCCCAA TTGCAAGCCA TCCAAAAATC CCGCCAAGAA TTTCTACTGA  
 25551 AAAAGGGTAA GGGGGTCTAC CTTGACCCCC AGACCGGCGA GGAATCAAC ACAAGGTTCC CTCAGGATGT  
 25621 CCCAACGACG AGAAAACAAG AAGTTGAAGG TGCAGCCGCC GCCCCAGAA GATATGGAGG AAGATTGGGA  
 25691 CAGTCAGGCA GAGGAGGCGG AGGAGGACAG TCTGGAGGAC AGTCTGGAGG AAGACAGTTT GGAGGAGGAA  
 25761 AACGAGGAGG CAGAGGAGGT GGAAGAAGTA ACCGCCGACA AACAGTTATC CTCGGCTGCG GAGACAAGCA  
 25831 ACAGCGCTAC CATCTCCGCT CCGAGTCGAG GAACCCGGCG GCGTCCCAGC AGTAGATGGG ACGAGACCGG  
 25901 ACGCTTCCCG AACCCAACCA GCGCTTCCAA GACCGGTAA GAGGATCGGC AGGGATACAA GTCTTGCGG  
 25971 GGGCATAAGA ATGCCATCAT CTCCTGCTTG CATGAGTGCG GGGGCAACAT ATCCTTCACG CCGCGCTACT  
 26041 TGCTATTCCA CCATGGGGTG AACTTTCCGC GCAATGTTTT GCATTACTAC CGTCACCTCC ACAGCCCCTA  
 26111 CTATAGCCAG CAAATCCCGA CAGTCTCGAC AGATAAAGAC AGCGGCGGCG ACCTCCAACA GAAAACACG  
 26181 AGCGGCAGTT AGAAAATACA CAACAAGTGC AGCAACAGGA GGATTAAAGA TTACAGCCAA CGAGCCAGCG  
 26251 CAAACCCGAG AGTTAAGAAA TCGGATCTTT CCAACCTGT ATGCCATCT CCAGCAGAGT CCGGGTCAAG  
 26321 AGCAGGAAC TAAAATAAAA AACCGATCTC TCGGTTCTGCT CACCAGAAGT TGTGTTGATC ACAAGAGCGA  
 26391 AGATCAACTT CAGCGCACTC TCGAGGACGC CGAGGCTCTC TTCAACAAGT ACTGCGCGCT GACTCTTAAA  
 26461 GAGTAGGCAG CGACCGCGCT TATTCAAAAA AGGCGGGAAT TACATCATCC TCGACATGAG TAAAGAAATT  
 26531 CCCACGCTT ACATGTGGAG TTATCAACCC CAAATGGGAT TGGCAGCAGG CGCCTCCCAG GACTACTCCA  
 26601 CCCGCATGAA TTGGCTCAGC GCCGGGCTT CTATGATTTT TCGAGTTAAT GATATACGCG CCTACCGAAA  
 26671 CCAAATACTT TTGGAACAGT CAGCTCTTAC CACCACGCCC CGCCAACACC TTAATCCCAG AAATTGGCCC

Figure 6, contd.

```

26741 GCCGCCCTAG TGTACCAGGA AAGTCCCGCT CCCACCACTG TATTACTTCC TCGAGACGCC CAGGCCGAAG
26811 TCCAAATGAC TAATGCAGGT GCGCAGTTAG CTGGCGGCTC CACCCTATGT CGTCACAGGC CTCGGCATAA
26881 TATAAAACGC CTGATGATCA GAGGCCGAGG TATCCAGCTC AACGACGAGT CCGTGAGCTC TCCGCTTGGT
26951 CTACGACCAG ACGGAATCTT TCAGATTGCC GGCTGCGGGA GATCTTCCTT CACCCCTCGT CAGGCTGTTC
27021 TGACTTTTGA AAGTTCGTCT TCGCAACCCC GCTCGGGCGG AATCGGGACC GTTCAATTTG TAGAGGAGTT
27091 TACTCCCTCT GTCTACTTCA ACCCCTTCTC CGGATCTCCT GGGCACTACC CGGACGAGTT CATACCGAAC
27161 TTCGACGCGA TTAGCGAGTC AGTGGACGGC TACGATTGAT GTCTGGTGAC GCGGCTGAGC TATCTCGGCT
27231 GCGACATCTA GACCACTGCC GCCGCTTTCG CTGCTTTGCC CGGGAACCTA TTGAGTTCAT CTACTTCGAA
27301 CTCCCCAAGG ATCACCCTCA AGGTCCGGCC CACGGAGTGC GGATTACTAT CGAAGGCAAA ATAGACTCTC
27371 GCCTGCAACG AATTTTCTCC CAGCGGCCCG TGCTGATCGA GCGAGACCAG GGAAACACCA CGGTTTCCAT
27441 CTACTGCATT TGTAATCACC CCGGATTGCA TGAAAGCCTT TGCTGTCTTA TGTGTACTGA GTTTAATAAA
27511 AACTGAATTA AGACTCTCCT ACGGACTGCC GCTTCTTCAA CCCGGATTTT ACAACCAGAA GAACAAAAC
27581 TTTCTGTCTG TCCAGGACTC TGTTAACTTC ACCTTTCCTA CTCACAAACT AGAAGCTCAA CGACTACACC
27651 GCTTTTCCAG AAGCATTTTC CCTACTAATA CTACTTTCAA AACCGGAGGT GAGCTCCACG GTCTCCCTAC
27721 AGAAAACCC TGGGTGGAAG CCGGCTTGT AGTACTAGGA ATTCTTGCGG GTGGGCTTGT GATTATTCTT
27791 TGCTACCTAT ACACACCTTG CTTACATTTT CTAGTGGTGT TGTGGTATTG GTTTAAAAAA TGGGGCCCAT
27861 ACTAGTCTTG CTTGTTTTAC TTTGCTTTT GGAACCGGGT TCTGCCAATT ACGATCCATG TCTAGACTTT
27931 GACCCAGAAA ACTGCACACT TACTTTTGCA CCCGACACAA GCCGCATCTG TGGAGTTCTT ATTAAGTGCG
28001 GATGGGAATG CAGGTCCGTT GAAATTACAC ACAATAACAA AACCTGGAAC AATACCTTAT CCACCACATG
28071 GGAGCCAGGA GTTCCCGAGT GGTACACTGT CTCTGTCCGA GGTCTGACG GTTCCATCCG CATTAGTAAC
28141 AACACTTTCA TTTTTTCTGA AATGTGCGAT CTGGCCATGT TCATGAGCAA ACAGTATTCT CTATGGCCTC
28211 CTAGCAAGGA CAACATCGTA ACGTCTTCCA TTGCTTATTG CTTGTGCGCT TGCTTCTTA CTGCTTTACT
28281 GTGCGTATGC ATACACCTGC TTGTAACCAC TCGCATCAAA AACGCCAATA ACAAGAAAAA AATGCCTTAA
28351 GCTCTTCTG TTTTCTGCTA TGGTCTCTT TGGTCTCTG AATCTGTACG GCTCTGTACG TGGGCTCTG
28421 GGACAAACAG TCGTCTCTAT CCCACTAGGA CATAATTACA CTCTCATAGG ACCCCCAATC ACTTCAGAGG
28491 TCATCTGGAC CAAACTGGGA AGCGTTGATT ACTTTGATAT AATCTGTAAC AAAACAAAAC CAATAATAGT
28561 AACTTGCAAC ATACAAAATC TTACATTGAT TAATGTTAGC AAAGTTTACA GCGGTTACTA TTATGGTTAT
28631 GACAGATACA GTAGTCAATA TAGAAATTAC TTGGTTCGTG TTACCCAGTT GAAAACCACG AAAATGCCAA
28701 ATATGGCAAA GATTGATCC GATGACAATT CTCTAGAAAC TTTTACATCT CCCACCACAC CCGACGAAAA
28771 AAACATCCCA GATTCAATGA TTGCAATTGT TGCAGCGGTG GCAGTGGTGA TGGCACTAAT AATAATATGC
28841 ATGCTTTTAT ATGCTTGTG ATACAAAAAG TTTTACTCCTA AAAAACAAGA TCTCTACTA AGGCTTAACA
28911 TTTAATTTCT TTTTATACAG CCATGGTTTC CACTACCACA TTCCTTATGC TTACTAGTCT CGCAACTCTG
28981 ACTTCTGCTC GCTCACACCT CACTGTAAT ATAGGCTCAA ACTGCACACT AAAAGGACCT CAAGGTGGTC
29051 ATGTCTTTTG GTGGAGAATA TATGACAATG GATGGTTTAC AAAACCATGT GACCAACCTG GTAGATTTTT
29121 CTGCAACGGC AGAGACCTAA CCATTATCAA CGTGACAGCA AATGACAAAG GCTTCTATTA TGGAACCGAC
29191 TATAAAAGTA GTTTAGATTA TAACATTATT GTACTGCCAT CTACCACTCC AGCACCCCGC ACAACTACTT
29261 TCTCTAGCAG CAGTGTGCTT AACAATACAA TTTCCAATCC AACCTTTGCC GCGCTTTTAA AACGCACTGT
29331 GAATAATTCT ACAACTTCAC ATACAACAAT TTCCACTTCA ACAATCAGCA TCATCGCTGC AGTGACAATT
29401 GGAATATCTA TTCTTGTTTT TACCATAACC TACTACGCTT GCTGCTATAG AAAAGACAAA CATAAAGGTG
29471 ATCCATTACT TAGATTTGAT ATTTAATTTG TTCTTTTTTT TTATTTACAG TATGGTGAAC ACCAATCATG
29541 GTACCTAGAA ATTTCTTCTT CACCATACTC ATCTGTGCTT TTAATGTTTG CGCTACTTTC ACAGCAGTAG
29611 CCACAGCAAC CCCAGACTGT ATAGGAGCAT TTGCTTCTTA TGCACTTTTT GCTTTTGTTA CTTGCATCTG
29681 CGTATGTAGC ATAGTCTGCC TGGTTATTAA TTTTTTCCAA CTCTAGACT GGATCCTTGT GCGAATTGCC
29751 TACCTGCGCC ACCATCCCGA ATACCGCAAC CAAAATATCG CCGCACTTCT TAGACTCATC TAAAACCATG
29821 CAGGCTATAC TACCAATATT TTTGCTTCTA TTGCTTCCCT ACGCTGTCTC AACCCAGCT GCCTATAGTA
29891 CTCCACCAGA ACACCTTAGA AAATGCAAT TCCAACAACC GTGGTCATTT CTGTCTTGCT ATCGAGAAAA
29961 ATCAGAAATC CCCCCAATT TAATAATGAT TGCTGGAATA ATTAATATAA TCTGTTGCAC CATAATTTC
30031 TTTTGTATAT ACCCCTATT TGATTTTGGC TGGAAATGCT CCAATGCACA TGATCATCCA CAAGACCCAG
30101 AGGAACACAT TCCCCACAA AACATGCAAC ATCCAATAGC GCTAATAGAT TACGAAAGTG AACCACAACC
30171 CCCACTACTC CCTGCTATTA GTTACTTCAA CCTAACCGGC GGAGATGACT GAAACACTCA CCACCTCCAA
30241 TTCCGCGAG GATCTGCTCG ATATGGACGG CCGCTCTCA GAACAACGAC TTGCCCAACT ACGCATCCGC
30311 CAGCAGCAGG AACGCGTGGC CAAAGAGCTC AGAGATGTCA TCCAAATTCA CCAATGCAAA AAAGGCATAT
30381 TCTGTTTGGT AAAACAAGCC AAGATATCCT ACGAGATCAC CGCTACTGAC CATCGCCTCT CTTACGAAC
30451 TGGCCCCCAA CGACAAAAAT TTACCTGCAT GGTGGGAATC AACCCCATAG TTATACCCA ACAAGTGGA
30521 GATACTAAGG GTTGCATTCA CTGCTCCTGC GATTCCATCG AGTGCACCTA CACCCTGCTG AAGACCTAT
30591 GCGGCCTAAG AGACCTGCTA CCAATGAATT AAAAAAAAT GATTAATAAA AAATCACTTA CTTGAAATCA
30661 GCAATAAGGT CTCTGTTGAA ATTTCTTCCC AGCAGCACCT CACTTCCCTC TTCCCAACTC TGGTATTCTA
30731 AACCCCGTTC AGCGGCATAC TTTCTCCATA CTTTAAAGGG GATGTCAAAT TTTAGCTCCT CTCCTGTACC
30801 CACAATCTTC ATGTCTTTCT TCCCAGATGA CCAAGAGAGT CCGGCTCAGT GACTCCTTCA ACCCTGTCTA
30871 CCCCATGAA GATGAAAGCA CCTCCCAACA CCCCTTTATA ACCCCAGGGT TTATTTCCC AAATGGCTTC
30941 ACCAAAGCC CAGACGGAGT TCTTACTTTA AAATGTTTAA CCCCATAAC AACCACAGGC GGATCTCTAC
31011 AGCTAAAAGT GGGAGGGGGA CTTACAGTGG ATGACACTGA TGGTACCTTA CAAGAAAACA TACGTGCTAC
31081 AGCACCCATT ACTAAAAATA ATCACTCTGT AGAACTATCC ATTGGAAATG GATTAGAAAC TCAAAACAAT
31151 AAACATATGT CCAAAATGGG AAATGGGTTA AAATTTAACA ACGGTGACAT TTGTATAAAG GATAGTATTA

```

Figure 6, contd.

```

31221 ACACCTTATG GACTGGAATA AACCTCCAC CTAAGTGTCA AATTGTGGAA AACACTAATA CAAATGATGG
31291 CAAACTTACT TTAGTATTAG TAAAAAATGG AGGGCTTGT AATGGCTACG TGTCTCTAGT TGGTGTATCA
31361 GACACTGTGA ACCAAATGTT CACACAAAAG ACAGCAAACA TCCAATTAAG ATTATATTTT GACTCTTCTG
31431 GAAATCTATT AACTGAGGAA TCAGACTTAA AAATTCCACT TAAAAATAAA TCTTCTACAG CGACCAGTGA
31501 AACTGTAGCC AGCAGCAAAG CCTTTATGCC AAGTACTACA GCTTATCCCT TCAACACCAC TACTAGGGAT
31571 AGTGAAAAC ACATTCATGG AATATGTTAC TACATGACTA GTTATGATAG AAGTCTATTT CCTTGAACA
31641 TTTCTATAAT GCTAAACAGC CGTATGATTT CTTCCAATGT TGCCTATGCC ATACAATTTG AATGGAATCT
31711 AAATGCAAGT GAATCTCCAG AAAGCAACAT AGCTACGCTG ACCACATCCC CCTTTTTCTT TTCTTACAT
31781 ACAGAAGACG ACAACTAAAA TAAAGTTTAA GTGTTTTTAT TTAAAAATCAC AAAATTTCGAG TAGTTATTTT
31851 GCCTCCACCT TCCCATTTGA CAGAATACAC CAATCTCTCC CCACGCACAG CTTTAAACAT TTGGATACCA
31921 TTAGAGATAG ACATTGTTTT AGATTCCACA TTCCAAACAG TTTCAGAGCG AGCCAATCTG GGGTCAGTGA
31991 TAGATAAAAA TCCATCGCGA TAGTCTTTTA AAGCGCTTTC ACAGTCCAAC TGCTGCGGAT GCGACTCCGG
32061 AGTTTGGATC ACGGTCACTT GGAAGAAGAA CGATGGGAAT CATAATCCGA AAACGGTATC GGACGATTGT
32131 GTCTCATCAA ACCCACAAGC AGCCGCTGTC TGCGTCGCTC CGTGGGACTG CTGTTTATGG GATCAGGGTC
32201 CACAGTTTCC TGAAGCATGA TTTTAATAGC CCTTAACATC AACTTTCTGG TCGGATGCGC GCAGCAACGC
32271 ATTCTGATTT CACTCAAATC TTTGCAGTAG GTACAACACA TTATTACAAT ATTGTTTAAAT AAACCATAAT
32341 TAAAAGCGCT CCAGCCAAAA CTCATATCTG ATATAATCGC CCCTGCATGA CCATCATACC AAAGTTTAAAT
32411 ATAAATTAAA TGACGTTCCC TCAAAAACAC ACTACCCACA TACATGATCT CTTTTGGCAT GTGCATATTA
32481 ACAATCTGTC TGTACCATGG ACAACGTTGG TTAATCATGC AACCCAATAT AACCTTCCGG AACCACTG
32551 CCAACACCGC TCCCCCAGCC ATGCATTGAA GTGAACCTG CTGATTACAA TGACAATGAA GAACCCAATT
32621 CTCTCGACCG TGAATCACTT GAGAATGAAA AATATCTATA GTGGCACAAC ATAGACATAA ATGCATGCAT
32691 CTTCTCATAA TTTTAACTC CTCAGGATTT AGAAACATAT CCCAGGGAAT AGGAAGCTCT TGCAGAACAG
32761 TAAAGCTGGC AGAACAAGGA AGACCACGAA CACAACCTAC ACTATGCATA GTCATAGTAT CACAATCTGG
32831 CAAACAGCGC TGGTCTTCAG TCAATGAAAG TCGGCTTCA TTTCTTCA CAACTGATTA CTGGCTCTG
32901 GTGTAAGGGT GATGTCTGGC GCATGATGTC GAGCGTGGCG GCAACCTTGT CATAATGGAG TTGCTTCCTG
32971 ACATTCTCGT ATTTTGTATA GCAAAACGCG GCCCTGGCAG AACACACTCT TCTTCGCCCT CTATCCTGCC
33041 GCTTAGCGTG TTCCGTGTGA TAGTTCAAGT ACAGCCACAC TCTTAAGTTG GTCAAAAGAA TGCTGGCTTC
33111 AGTTGTAATC AAAACTCCAT CGCATCTAAT TGTCTGAGG AAATCATCCA CGGTAGCATA TGCAAAATCCC
33181 AACCAAGCAA TGCAACTGGA TTGCGTTTCA AGCAGGAGAG GAGAGGGAAG AGACGGAAGA ACCATGTTAA
33251 TTTTTATTCC AAACGATCTC GCAGTACTTC AAATTGTAGA TCGCGCAGAT GGCATCTCTC GCCCCCACTG
33321 TGTGGTGAA AAAGCACAGC TAAATCAAAA GAAATGCGAT TTTCAAGGTG CTCAACGGTG GCTTCCAACA
33391 AAGCCTCCAC GCGCACATCC AAGAACAAAA GAATACCAAA AGAAGGAGCA TTTTCTAACT CCTCAATCAT
33461 CATATTACAT TCCTGCACCA TTCCCAGATA ATTTTCAGCT TTCCAGCCTT GAATTATTCTG TGTCAGTTCT
33531 TGTGGTAAAT CCAATCCACA CATTACAAAC AGGTCCCGGA GGGCGCCCTC CACCACCATT CTTAAACACA
33601 CCCTCATAAT GACAAAATAT CTTGCTCCTG TGTCACCTGT AGCGAATTGA GAATGGCAAC ATCAATTGAC
33671 ATGCCCTTGG CTCTAAGTTC TTCTTTAAGT TCTAGTTGTA AAAACTCTCT CATATTATCA CCAAACTGCT
33741 TAGCCAGAAG CCCCCCGGGA ACAAGAGCAG GGGACGCTAC AGTGCAGTAC AAGCGCAGAC CTCCCCAATT
33811 GGCTCCAGCA AAAACAAGAT TGAATAAGC ATATTGGGAA CCACCAGTAA TATCATCGAA GTTGCTGGAA
33881 ATATAATCAG GCAGAGTTTC TTGTAGAAAT TGAATAAAG AAAAATTTGC CAAAAAACA TTCAAAACCT
33951 CTGGGATGCA AATGCAATAG GTTACCGCGC TGCGCTCCAA CATTGTAGT TTTGAATTAG TCTGCAAAAA
34021 TAAAAAATAA ACAAGCGTCA TATCATAGTA GCCTGACGAA CAGGTGGATA AATCAGTCTT TCCATCACAA
34091 GACAAGCCAC AGGGTCTCCA GCTCGACCTT CGTAAAACCT GTCATCGTGA TTAAACAACA GCACCGAAAG
34161 TTCCTCGCGG TGACCAGCAT GAATAAGTCT TGATGAAGCA TACAATCCAG ACATGTTAGC ATCAGTTAAG
34231 GAGAAAAAAC AGCCAACATA GCCTTTGGGT ATAATTATGC TTAATCGTAA GTATAGCAAA GCCACCCCTC
34301 GCGGATACAA AGTAAAAGGC ACAGGAGAAT AAAAAATATA ATTATTTCTC TGCTGCTGTT TAGGCAACGT
34371 CGCCCCGGT CCCTCTAAAT ACACATACAA AGCCTCATCA GCCATGGCTT ACCAGAGAAA GTACAGCGGG
34441 CACACAAACC ACAAGCTCTA AAGTCACTCT CCAACCTSTC CACAATATAT ATACACAAGC CCTAACTGA
34511 CGTAATGGGA CTAAAGTGTA AAAAATCCCG CCAAACCCAA CACACACCCC GAAACTGCGT CACCAGGGAA
34581 AAGTACAGTT TCACTTCCGC AATCCCAACA AGCGTCACTT CCTCTTTCTC ACGGTACGTC ACATCCCATT
34651 AACTTACAAC GTCATTTTCC CACGGCCGCG CCGCCCCCTT TAACCGTTAA CCCACAGCC AATCACCACA
34721 CGGCCACAC TTTTAAAAAT CACCTCATTT ACATATTGGC ACCATTCCAT CTATAAGGTA TATTATTGAT
34791 GATG

```



Figure 7






Figure 8



143

Figure 9



144

Figure 10




Figure 11







Figure 12



## Construction of cosmid vector pWE.Ad35.pIX-rITR




Figure 13

Figure 14




Figure 15




Figure 16






Figure 17




Figure 18




Figure 19




Figure 20




Figure 21








Figure 22b





FIGURE 23

1	MLLQMKRRARPS	EDTFNPVYPY	DE	ETGPTVP	PFLTP	PFVSP	Fib 5 protein					
1	MLLQMKRRARPS	EDTFNPVYPY	ED	ESSSQH	PFINP	GFISG	Fib 16 protein					
1	MLLQMKRRARPS	EDTFNPVYPY	ED	ESTSQH	PFINP	GFISG	Fib 35 protein					
1	MLLQMKRRARPS	EDTFNPVYPY	ED	ESTSQH	PFINP	GFISG	Fib 51 protein					
41	NGFQES	PPGVLS	LRLSE	PLVTS	NGMLA	LKMG	GNGLS	SLDEAG	Fib 5 protein			
40	NGFAQ	SPD	GVLT	LKCVN	PLTT	TASG	PLQLK	VGS	SLTV	DTID	Fib 16 protein	
40	NGFTQ	SPD	GVLT	LKCLT	PLTT	TGGS	PLQLK	VGG	GLTV	DDTD	Fib 35 protein	
40	NGFTQ	SPD	GVLT	UNCLT	PLTT	TGGS	PLQLK	VGG	GLTV	DDTD	Fib 51 protein	
81	NLTSON	VTTVSP	PLKKT	KSIN	LEIS	APLTV	TSEAL	TVAA	Fib 5 protein			
80	GSLEEN	IT-AAA	PLTK	TNHS	IGLL	IGSGL	QTT	-	-	-	Fib 16 protein	
80	GTLOEN	IR-ATA	PITK	NNHS	VELS	IGNGL	LET	-	-	-	Fib 35 protein	
80	GTLOEN	IR-VTA	PITK	NNHS	VELS	IGNGL	LET	-	-	-	Fib 51 protein	
121	AAPLMV	AGNTLT	MOSQ	APLTV	HDSK	LSIAT	QG	PLTV	SE	GK	Fib 5 protein	
110	-	-	-	-	-	-	-	-	-	-	Fib 16 protein	
110	-	-	-	-	-	-	-	-	-	-	Fib 35 protein	
110	-	-	-	-	-	-	-	-	-	-	Fib 51 protein	
161	LALQTS	GPLTT	TTDS	STLT	ITAS	PPLTT	ATGS	LGID	LKEPI	Fib 5 protein		
129	LCL	-	-	-	-	-	-	-	-	-	Fib 16 protein	
116	-	-	-	-	-	-	-	-	-	-	Fib 35 protein	
116	-	-	-	-	-	-	-	-	-	-	Fib 51 protein	
201	YTONGK	LGLKY	GAPL	HVT	DD	NTLT	VATG	PGVT	INNT	SLQ	Fib 5 protein	
132	-	-	-	-	-	-	-	-	-	-	Fib 16 protein	
116	-	-	-	-	-	-	-	-	-	-	Fib 35 protein	
116	-	-	-	-	-	-	-	-	-	-	Fib 51 protein	
241	TKVTGA	LGFDS	QGNM	QNLN	VAGGL	RIDSON	RRLI	LDVSY	PP	Fib 5 protein		
132	-	-	-	-	-	-	-	-	-	-	Fib 16 protein	
116	-	-	-	-	-	-	-	-	-	-	Fib 35 protein	
116	-	-	-	-	-	-	-	-	-	-	Fib 51 protein	
281	DAQNOL	NLR	LG	QPLF	INSA	HNLD	INYN	KGLY	LFTAS	NNS	Fib 5 protein	
132	-	-	-	-	-	-	-	-	-	-	Fib 16 protein	
116	-	-	-	-	-	-	-	-	-	-	Fib 35 protein	
116	-	-	-	-	-	-	-	-	-	-	Fib 51 protein	
321	KKLEVN	LSTAK	GLM	FDATA	IAIN	AGDGL	EF	GSPN	APNT	NP	Fib 5 protein	
137	-	-	-	-	-	-	-	-	-	-	Fib 16 protein	
116	-	-	-	-	-	-	-	-	-	-	Fib 35 protein	
116	-	-	-	-	-	-	-	-	-	-	Fib 51 protein	
361	LKTKIG	HGLE	FDSN	KAMV	PKLGT	GLSP	DSTGA	ITVGN	KNN	Fib 5 protein		
137	LITK	-	-	-	-	-	-	-	-	-	Fib 16 protein	
116	-	-	-	-	-	-	-	-	-	-	Fib 35 protein	
116	-	-	-	-	-	-	-	-	-	-	Fib 51 protein	
401	DKLT	LWTT	PAPSP	NCRL	-	-	NAEK	DAKLT	LVLTK	CGSQ	IL	Fib 5 protein
166	-	-	-	-	-	-	-	-	-	-	-	Fib 16 protein
137	-	-	-	-	-	-	-	-	-	-	-	Fib 35 protein
137	-	-	-	-	-	-	-	-	-	-	-	Fib 51 protein
438	ATVSVL	AVKGS	LAPI	-	SGTV	QSAHL	IIRF	DENG	VLLN	NSF	Fib 5 protein	
203	GYIT	LMGAS	EYTN	TLFKN	NQVT	IDYN	LAFD	NTG	QYIT	YLS	Fib 16 protein	
174	GYVS	LVGV	SDTV	NQMFT	QKTANI	QLRL	YFDSS	GNNL	LTEES	Fib 35 protein		
174	GYVS	LVGV	SDTV	NQMFT	OKSATI	QLRL	YFDSS	GNNL	LTEES	Fib 51 protein		
477	LDPEY	WNFR	NGDL	TEGT	AYTNA	VGFMP	NLSAY	PKSH	GKTA	Fib 5 protein		
243	SLKSN	INF	KDNQ	NMAT	GTTT	SAK	GFM	PSTT	AYPFI	-	-	Fib 16 protein
214	DLKI	PLKN	KSSTA	-TSET	VASSK	AFMP	PSTT	AYPFN	-	-	-	Fib 35 protein
214	NLKI	PLKN	KSSTA	-TSE	ATSSK	AFMP	PSTT	AYPFN	-	-	-	Fib 51 protein
517	KSNIV	SQVY	LNGD	-	-	-	-	-	-	-	-	Fib 5 protein
280	ATET	LNE	YIYG	ECY	YKST	-NGT	LF	PLKV	TV	TLN	RRMLAS	Fib 16 protein
250	TRDS	-	-	-	-	-	-	-	-	-	-	Fib 35 protein
250	TRDS	-	-	-	-	-	-	-	-	-	-	Fib 51 protein
548	DTTP	SAYS	MSFS	WDWS	GHN	YINE	-	-	-	-	-	Fib 5 protein
319	GM	-	-	-	-	-	-	-	-	-	-	Fib 16 protein
288	NV	-	-	-	-	-	-	-	-	-	-	Fib 35 protein
288	NV	-	-	-	-	-	-	-	-	-	-	Fib 51 protein
585	E	-	-	-	-	-	-	-	-	-	-	Fib 5 protein
356	D	-	-	-	-	-	-	-	-	-	-	Fib 16 protein
325	D	-	-	-	-	-	-	-	-	-	-	Fib 35 protein
325	D	-	-	-	-	-	-	-	-	-	-	Fib 51 protein

Figure 24



**Figure 25**




Figure 26




Figure 27






Figure 28






Figure 29



**Figure 30 Temperature dependent growth of  
PER.C6**



**Figure 31** DBP levels in PER.C6 cells transfected with pcDNA3, pcDNA3wtE2A or pcDNA3ts125E2A






Figure 32 Suspension growth of PE3.C6ts125E2A C5-9




**Figure 33** Growth curve PER.C6 and PER.C6tsE2A





# Figure 34 Stability of PE3.C6ts125E2A







Figure 35 BtTa activity in Per/E2A/t A clones





European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 00 20 1738

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X,D	BASLER C. F. ET AL.: "Sequence of the immunoregulatory early region 3 and flanking sequences of adenovirus type 35" GENE, vol. 170, no. 2, 8 May 1996 (1996-05-08), page 249-254 XP004042835 ISSN: 0378-1119	8,9	C12N15/86 C12N15/34 C12N5/10 A61K48/00
Y	* the whole document *	1-3,5-7, 10-19, 21,22	
X	WO 96 12030 A (RHONE POULENC RORER SA ;LEE MARTIN (FR); PERRICAUDET MICHEL (FR)) 25 April 1996 (1996-04-25)	23	
Y	* the whole document *	1-3,5-7, 10-19, 21,22	
Y	FALLAUX F. J. ET AL.: "New helper cells and matched early region 1-deleted adenovirus vectors prevent generation of replication-competent adenoviruses" HUMAN GENE THERAPY, vol. 9, no. 13, 1 September 1998 (1998-09-01), pages 1909-1917, XP002111678 ISSN: 1043-0342	14	
	* the whole document *		
Y	WO 96 13598 A (UNIV PENNSYLVANIA ;WILSON JAMES M (US); KELLEY WILLIAM M (US); FIS) 9 May 1996 (1996-05-09)	18,19	
	* the whole document *		
	--- -/--		
The present search report has been drawn up for all claims			
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>7 September 2000</b>	Examiner <b>Mandl, B</b>
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ..... &amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.82 (P04C01)



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 00 20 1738

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X,D	FLOMENBERG P. R. ET AL.: "Sequence and genetic organization of Adenovirus Type 35 early region 3." JOURNAL OF VIROLOGY, vol. 62, no. 11, 1988, pages 4431-4437, XP002119500 * the whole document *	8,9	
X	BASLER C.F. ET AL.: "Subgroup B adenovirus type 35 early region 3 mRNAs differ from those of the subgroup C adenoviruses." VIROLOGY, vol. 215, 1996, pages 165-177, XP002119501 * the whole document *	8,9	
A	LEE M. G. ET AL.: "THE CONSTITUTIVE EXPRESSION OF THE IMMUNOMODULATORY GP 19K PROTEIN IN E1-, E3- ADENOVIRAL VECTORS STRONGLY REDUCES THE HOST CYTOTOXIC T CELL RESPONSE AGAINST THE VECTOR" GENE THERAPY, vol. 2, no. 4, 1 June 1995 (1995-06-01), pages 256-262, XP000561548 ISSN: 0969-7128 * the whole document *	1-23	
A	WO 97 12986 A (CORNELL RES FOUNDATION INC) 10 April 1997 (1997-04-10) * page 11, line 3 - line 35 * * page 13, line 1 - line 30 *	1-23	
A	WO 98 22609 A (ARMENTANO DONNA E ; GREGORY RICHARD J (US); GENZYME CORP (US); SMIT) 28 May 1998 (1998-05-28) * the whole document *	1-23	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
Place of search	Date of completion of the search	Examiner	
THE HAGUE	7 September 2000	Mandl, B	
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03 82 (P04C01)



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number  
EP 00 20 1738

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	GALL J. ET AL.: "ADENOVIRUS TYPE 5 AND 7 CAPSID CHIMERA: FIBER REPLACEMENT ALTERS RECEPTOR TROPISM WITHOUT AFFECTING PRIMARY IMMUNE NEUTRALIZATION EPITOPES" JOURNAL OF VIROLOGY, vol. 70, no. 4, 1 April 1996 (1996-04-01), pages 2116-2123, XP002050655 ISSN: 0022-538X * the whole document *	1-23	
P,X	SHAYAKHMETOV D. M. ET AL.: "Efficient gene transfer into human CD34+ cells by a retargeted adenovirus vector." JOURNAL OF VIROLOGY, vol. 74, no. 6, March 2000 (2000-03), pages 2567-2583, XP000938716 ISSN: 0022-538X * the whole document *	1,2,4-9, 12,13, 15-17, 20-23	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>7 September 2000</b>	Examiner <b>Mandl, B</b>
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ..... &amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03.82 (P04C01)



**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 00 20 1738

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

07-09-2000

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9612030 A	25-04-1996	FR 2725726 A	19-04-1996
		AU 712243 B	04-11-1999
		AU 3658495 A	06-05-1996
		CA 2201399 A	25-04-1996
		EP 0787198 A	06-08-1997
		FI 971613 A	16-04-1997
		JP 10507079 T	14-07-1998
		NO 971590 A	07-04-1997
		ZA 9508686 A	22-05-1996
WO 9613598 A	09-05-1996	US 5856152 A	05-01-1999
		AU 695811 B	20-08-1998
		AU 4405596 A	23-05-1996
		CA 2203808 A	09-05-1996
		EP 0797678 A	01-10-1997
		JP 10507928 T	04-08-1998
		US 5871982 A	16-02-1999
WO 9712986 A	10-04-1997	US 5837511 A	17-11-1998
		EP 0866873 A	30-09-1998
		JP 11514866 T	21-12-1999
WO 9822609 A	28-05-1998	US 5877011 A	02-03-1999
		AU 5455298 A	10-06-1998
		EP 0946742 A	06-10-1999

EPO FORM P0439

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82